

Assessment of adrenaline-induced DNA damage in whole blood cells with the comet assay

Dijana Topalović¹, Dragana Dekanski², Biljana Spremo-Potparević¹, Ninoslav Djelić³,
Vladan Bajić⁴, and Lada Živković¹

Department of Biology and Human Genetics, Faculty of Pharmacy, University of Belgrade¹, Biomedical Research, R&D Institute, Galenika a.d.², Department of Biology, Faculty of Veterinary Medicine, University of Belgrade³, Laboratory for Radiobiology and Molecular Genetics, Institute for Nuclear Research "Vinča", University of Belgrade⁴, Belgrade, Serbia

[Received in May 2018; Similarity Check in May 2018; Accepted in November 2018]

Harmful effects of elevated levels of catecholamines are mediated by various mechanisms, including gene transcription and formation of oxidation products. The aim of this study was to see whether the molecular mechanisms underlying the damaging action of adrenaline on DNA are mediated by reactive oxygen species (ROS). To do that, we exposed human whole blood cells to 10 $\mu\text{mol L}^{-1}$ adrenaline or 50 $\mu\text{mol L}^{-1}$ H_2O_2 (used as positive control) that were separately pre-treated or post-treated with 500 $\mu\text{mol L}^{-1}$ of quercetin, a scavenger of free radicals. Quercetin significantly reduced DNA damage in both pre- and post-treatment protocols, which suggests that adrenaline mainly acts via the production of ROS. This mechanism is also supported by gradual lowering of adrenaline and H_2O_2 -induced DNA damage 15, 30, 45, and 60 min after treatment. Our results clearly show that DNA repair mechanisms are rather effective against ROS-mediated DNA damage induced by adrenaline.

KEY WORDS: *DNA repair; hydrogen peroxide; genotoxicity; quercetin; ROS*

The primary role of adrenaline is to mobilise resources in response to emotional and physical stress. Apart from its primary effect, adrenaline regulates insulin secretion (1) and stimulates platelet aggregation (2). It is used to treat asthma, cardiac arrest, and anaphylaxis (3). It has also been reported for cardiotoxicity (4), oesophageal cancer cell proliferation, and resistance to chemotherapy in breast cancer (5, 6). Excess adrenaline in the circulation under stress can increase blood glucose levels, blood pressure, and heart rate (7, 8) and may induce myocardial cell damage (9).

Furthermore, there are indications that the catechol group of dopamine, noradrenaline, and adrenaline is involved in the redox cycle that leads to the formation of reactive oxygen species (ROS) (10-12). Recent experimental evidence has shown the ability of adrenaline to make changes in the DNA (13-15). Some studies (10, 16, 17) suggest that oxidation and cyclisation of adrenaline result in the formation of by-products such as ROS, which may cause DNA damage.

However, studies investigating possible mechanisms of adrenaline genotoxicity are scarce and their findings ambiguous. Some confirm adrenaline-induced DNA breaks (14) and significant cell-cycle delays needed for the cell to

repair genetic damage (18) but others report no association between adrenaline and chromosome aberrations in human or chicken peripheral blood lymphocytes (19, 20) or other cytogenetically detectable effects (18).

The aim of our study was to look deeper into the genotoxic action of adrenaline by focusing on the cell capacity to activate DNA repair mechanisms for damages caused by this stress hormone. To do that, we used the comet assay to quantify and analyse primary DNA damage and repair capacity in individual cells.

MATERIALS AND METHODS

Blood sampling

Whole blood samples were taken with a finger prick from three healthy female volunteers (aged between 20 and 40 years) and collected in heparinised containers. The volunteers did not smoke, drink alcohol, or receive any medicines or dietary supplements. They gave informed consent and the study was approved by the Ethics Committee of the Faculty of Pharmacy, Belgrade, Serbia (Approval No. 1103/2).

Study design

Adrenaline (CAS No. 51-43-4, epinephrine) was obtained from Sigma-Aldrich Chemie (St. Louis, MO,

Correspondence to: Dijana Topalović, MSc, Department of Biology and Human Genetics, Faculty of Pharmacy, University of Belgrade, Vojvode Stepe 450, 11000 Belgrade, Serbia,
E-mail: dijana.topalovic@pharmacy.bg.ac.rs

USA). Hydrogen peroxide (H_2O_2 , CAS No. 7722-84-1), known to induce chromosomal aberrations, gene mutations, and DNA damage (21, 22), was obtained from ZORKA Pharma (Šabac, Serbia) and used as positive control. For negative control, cells were incubated with phosphate buffered saline (PBS, Fisher Scientific, Pittsburgh, PA). The effects of adrenaline and H_2O_2 were investigated separately in our earlier research (14) at four concentrations (5, 10, 50, and 150 $\mu\text{mol L}^{-1}$ for adrenaline and 5, 10, 25, and 50 $\mu\text{mol L}^{-1}$ for H_2O_2) that corresponded to the ones used in other studies of adrenaline-induced DNA damage (12, 18). For this study we selected the concentrations of 10 $\mu\text{mol L}^{-1}$ of adrenaline and 50 $\mu\text{mol L}^{-1}$ of H_2O_2 , because they produced significant DNA damage in human whole blood cells without significantly affecting cell viability.

In order to ascertain the formation of ROS as the presumed mechanism underlying adrenaline-induced DNA damage, we also treated the cells with quercetin, a well-known scavenger of free radicals, in the concentration of 500 $\mu\text{mol L}^{-1}$ taken from literature data (19, 23). It was also the most effective concentration in our recent study (24).

Two protocols were used to evaluate the contribution of different mechanisms underlying adrenaline genotoxicity: In the pre-treatment protocol, the cells were incubated with quercetin at 37 °C for 30 min prior to their exposure to adrenaline (at 37 °C for 30 min). In the post-treatment protocol, the cells were treated with quercetin under the same conditions after their exposure to adrenaline. The same experiment was conducted with quercetin and H_2O_2 following the same protocols. In all experiments, cell preparations were also incubated with PBS or adrenaline/ H_2O_2 , which served as negative or positive control.

An additional set of experiments was conducted to identify the repair kinetics of adrenaline- and H_2O_2 -induced DNA damage. Following adrenaline (10 $\mu\text{mol L}^{-1}$) treatment at 37 °C for 30 min, the cells were treated with PBS at 37 °C for 15, 30, 45, and 60 min. Positive control cells were exposed to 50 $\mu\text{mol L}^{-1}$ H_2O_2 and then treated with PBS in the same way as described for adrenaline.

Single cell gel electrophoresis assay (comet assay)

The alkaline version of the comet assay was performed as described by Singh et al. (25). Before the treatments, 6 μL of freshly obtained whole blood samples were suspended in 0.67 % low melting point agarose (LMP, Sigma-Aldrich, St. Louis), pipetted onto slides precoated with a layer of 1 % normal melting point agarose (Sigma-Aldrich, St. Louis), and left at 4 °C for 5 min to solidify. The cell suspensions were then treated as described above. After treatment, the samples were covered with the 0.5 % LMP agarose and also left at 4 °C for 5 min, after which time they were submerged in freshly prepared and cooled lysing solution (2.5 mol L^{-1} NaCl, 100 mmol L^{-1} EDTA, 10 mmol L^{-1} Tris, 1 % Triton X 100, and 10 % dimethylsulfoxide, pH 10, adjusted with NaOH), and left

to stay at 4 °C overnight. The next day, the slides were electrophoresed at 24 V and 300 mA, washed with neutralisation buffer and distilled water, and then stained with ethidium bromide (20 $\mu\text{g L}^{-1}$). The comets were scored on an Olympus BX 50 microscope (Olympus Optical Co., GmbH, Hamburg, Germany) equipped with an HBO mercury lamp (50 W, 516-560 nm, Carl Zeiss Microscopy, Jena, Germany) at 100x magnification.

DNA damage was evaluated according to Anderson et al. (22). Comets were visually scored and classified into five categories corresponding to the extent of DNA migration: (A) no damage, <5 %; (B) low level damage, 5-20 %; (C) medium level damage, 20-40 %; (D) high level damage, 40-95 %; and (E) total damage, >95 %. One hundred randomly selected nucleoids were counted on each of two slides per treatment. The DNA damage was characterised as the total DNA migration over 5 % (B + C + D + E comet classes), and the mean value was calculated for 200 nucleoids (100 per slide) for each experiment in triplicate. Apoptotic and necrotic cells were set apart from normal cells and excluded from the analyses following the instructions given by Singh (26).

Statistical analysis

We used one-way analysis of variance (ANOVA) with Tukey's *post-hoc* test to compare treatments and respective controls. Data were expressed as mean \pm standard error of the mean (SEM) of three measurements. Differences at $p < 0.05$ were considered statistically significant.

RESULTS

Both adrenaline and H_2O_2 caused significant primary DNA damage (Figure 1). Adrenaline induced DNA damage in around 20 % and H_2O_2 in around 30 % of the cells. The latter also induced a higher level of DNA damage (54.1 % of the cells had category D+E damage) than adrenaline (20.1 % with D+E damage). Quercetin, in turn, attenuated their damage significantly with both protocols (Figure 1) and was more effective in post-treatment, but the difference between the protocols was not statistically significant.

Figure 2 shows DNA damage repair in cells over time. While adrenaline-induced DNA damage was reduced gradually, and the reduction reached the level of significance at 45 and 60 min, H_2O_2 -induced damage repair reached significance earlier, at 30 min, but then rose again, still maintaining the level of significance. In fact, DNA damage increased at 60 min in both treatments.

DISCUSSION

This study has confirmed the damaging effect of 10 $\mu\text{mol L}^{-1}$ adrenaline on DNA in human whole blood cells. Quercetin (500 $\mu\text{mol L}^{-1}$) added before or after treatment




Figure 1 a) Protective effect of quercetin (Q) against DNA damage induced by a) adrenaline (A) and b) by H₂O₂. Q+A – pre-treatment protocol; A+Q – post-treatment protocol; Q+H₂O₂ – pre-treatment protocol; H₂O₂+Q – post-treatment protocol. Bars represent the mean number of cells with DNA damage ±SEM; B+C – low and medium damage; D+E – high and total damage; * p<0.05 vs. adrenaline- and H₂O₂-treated control

with adrenaline (10 μmol L⁻¹) or H₂O₂ (50 μmol L⁻¹) significantly reduced DNA damage. It has already been reported to protect against DNA strand breaks by acting as a metal chelator (27) and by increasing the concentration of enzymatic and non-enzymatic components of the antioxidant cell system (28, 29). In our study it displayed higher efficiency in the post-treatment protocol against both oxidants. Since its most pronounced feature is the scavenging of superoxide anions, peroxy nitrite, and singlet oxygen and •OH radicals (23, 30, 31), this finding confirms that the main mechanism of adrenaline action is through free radicals. That ROS is the mechanism of adrenaline genotoxicity is also supported by the reports of antioxidant prevention of adrenaline-induced DNA damage (14, 15, 32).

As for DNA repair, it significantly reduced damage 45 and 60 min into adrenaline exposure, which is in accordance with Djelić et al. (18), where DNA damage significantly dropped after 1 h. These findings suggest that adrenaline genotoxicity decreases over time and that one hour is sufficient for cells to activate DNA repair mechanisms and compensate for the damage (33, 34). This repair was significantly effective even earlier with H₂O₂, probably

because peroxide caused greater damage. Cells tend to recover faster from the impact of a more potent oxidant by activating their repair mechanisms earlier (35). Another reason for less efficient repair with adrenaline is that oxidation caused by high concentrations of adrenaline takes the quinone pathway, which yields larger amounts of highly reactive intermediates, which, in turn, slow down DNA repair (36) and prolong cell recovery.

One interesting finding in our study is that DNA repair from the effects of adrenaline weakened at 60 min. Reasons may be several. First, the concentration of adrenaline (10 μmol L⁻¹) was high enough to trigger a redox cycle, multiplying the production of ROS. Under these conditions, antioxidative enzymes cannot protect the cells against excess ROS (15). This is also confirmed by our finding of the stronger effect of quercetin in the post-treatment protocol. Second, ROS dominantly causes single- and double-stranded DNA breaks, and it takes longer for repair to compensate for double-stranded breaks (37). The resulting imbalance can also cause other types of primary DNA lesions, and overcome cell repair capacity (37, 38). Third, stress hormones induce DNA damage that prevents cell cycle checkpoint activities and the onset of apoptosis.




Figure 2 DNA damage repair in cells exposed to a) adrenaline (A) and b) H₂O₂ at 15, 30, 45, and 60 min after treatment. Bars represent the mean number of cells with DNA damage (B+C+D+E comet classes) ±SEM; * p<0.05 vs. adrenaline- and H₂O₂-treated control

Stress increases ROS-generated damage and affects the DNA repair (39).

The similar responses of adrenaline and H₂O₂ to quercetin, similar DNA damage repair, as well as the stronger effect of quercetin when given after either of the oxidants, support our hypothesis that ROS production is the underlying mechanism of adrenaline genotoxicity. Highly reactive intermediates that arise from oxidised adrenaline, triggered by stress, weaken DNA repair. Even so, our results clearly show that DNA repair mechanisms are rather effective against ROS-mediated DNA damage induced by adrenaline.

Acknowledgments

This research was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grants OI 173034 and III46002) and the hCOMET COST (Action No. CA 15132).

REFERENCES

1. Verberne AJM, Korim WS, Sabetghadam A, Llewellyn-Smith IJ. Adrenaline: insights into its metabolic roles in hypoglycaemia and diabetes. *Br J Pharmacol* 2016;173:1425-37. doi: 10.1111/bph.13458
2. Spalding A, Vaitkevicius H, Dill S, MacKenzie S, Schmaier A, Lockette W. Mechanism of epinephrine-induced platelet aggregation. *Hypertension* 1998;31:603-7. doi: 10.1161/01.HYP.31.2.603
3. The American Society of Health-System Pharmacists (ASHSP). Epinephrine [displayed 15 August 2015]. Available at <https://www.drugs.com/monograph/epinephrine.html>
4. Adameova A, Abdellatif Y, Dhalla NS. Role of the excessive amounts of circulating catecholamines and glucocorticoids in stress-induced heart disease. *Can J Physiol Pharm* 2009;87:493-514. doi: 10.1139/y09-042
5. Drell IV TL, Joseph J, Lang K, Niggemann B, Zaenker KS, Entschladen F. Effects of neurotransmitters on the chemokinesis and chemotaxis of MDA-MB-468 human breast carcinoma cells. *Breast Cancer Res Tr* 2003;80:63-70. doi: 10.1023/A:1024491219366
6. Pu J, Bai D, Yang X, Lu X, Xu L, Lu J. Adrenaline promotes cell proliferation and increases chemoresistance in colon cancer HT29 cells through induction of miR-155. *Biochem Bioph Res Co* 2012;428:210-5. doi: 10.1016/j.bbrc.2012.09.126
7. Sherwood L, editor. *Human Physiology: From Cells to Systems*. 7th ed. Belmont (CA): Thomson Brooks/Cole; 2009.
8. Gavrilović Lj, Stojiljković V, Kasapović J, Pejić S, Todorović A, Pajović BS, Dronjak S: Chronic physical stress changes gene expression of catecholamine biosynthetic enzymes in the adrenal medulla of adult rats. *Acta Vet-Beograd* 2012;62:151-69. doi: 10.2298/AVB1203151G
9. Dhalla NS, Sasaki H, Mochizuki S, Dhalla KS, Liu X, Elimban V. Catecholamine-induced cardiomyopathy. In: D. Acosta, Jr., editor. *Cardiovascular toxicology*. 3rd ed. New York (NY): CRC Press; 2001. p. 263-318.
10. Miura T, Muraoka S, Fujimoto Y, Zhao K. DNA damage induced by catechol derivatives. *Chem-Biol Interact* 2000;126:125-36. doi: 10.1016/S0009-2797(00)00156-3
11. Djelić N, Anderson D. The effect of the antioxidant catalase on oestrogens, triiodothyronine, and noradrenaline in the Comet assay. *Teratogen Carcin Mut* 2003;23(Suppl 2):69-81. doi: 10.1002/tcm.10084
12. Dobrzyńska MM, Baumgartner A, Anderson D. Antioxidants modulate thyroid hormone and noradrenaline-induced DNA damage in human sperm. *Mutagenesis* 2004;19:325-30. doi: 10.1093/mutage/geh037
13. Flint MS, Baum A, Chambers WH, Jenkins FJ. Induction of DNA damage, alteration of DNA repair and transcriptional activation by stress hormones. *Psychoneuroendocrinology* 2007;32:470-9. doi: 10.1016/j.psyneuen.2007.02.013
14. Čabarkapa A, Živković L, Žukovec D, Djelić N, Bajić V, Dekanski D, Spremo-Potparević B. Protective effect of dry olive leaf extract in adrenaline induced DNA damage evaluated using *in vitro* comet assay with human peripheral leukocytes. *Toxicol in Vitro* 2014;28:451-6. doi: 10.1016/j.tiv.2013.12.014
15. Radaković M, Đelić N, Stevanović J, Andelković M, Kolarević S, Dačić S, Stanimirović Z. The investigation of DNA damage induced by adrenaline in human lymphocytes *in vitro*. *Acta Vet-Beograd* 2014;64:281-92. doi: 10.2478/acve-2014-0027
16. Behonick GS, Novak MJ, Nealley EW, Baskin SI. Toxicology update: the cardiotoxicity of the oxidative stress metabolites of catecholamines (aminochromes). *J Appl Toxicol* 2001;21(Suppl 1):S15-22. doi: 10.1002/jat.793
17. Genova ML, Abd-Elsalam NM, Mahdy el SME, Bernacchia A, Lucarini M, Pedulli GF, Lenaz G. Redox cycling of adrenaline and adrenochrome catalysed by mitochondrial Complex I. *Arch Biochem Biophys* 2006;447:167-73. doi: 10.1016/j.abb.2006.01.010
18. Djelić N, Radaković M, Spremo-Potparević B, Živković L, Bajić V, Stevanović J, Stanimirović Z. Evaluation of cytogenetic and DNA damage in human lymphocytes treated with adrenaline *in vitro*. *Toxicol in Vitro* 2015;29:27-33. doi: 10.1016/j.tiv.2014.08.001
19. Djelić N, Djelić D, Spremo-Potparević B, Marković B, Živković L. Cytogenetic analysis of the effects of epinephrine on cultured human lymphocytes. *Acta Vet-Beograd* 2003;53:113-20.
20. Lankoff A, Carmichael WW, Grasman KA, Yuan M. The uptake kinetics and immunotoxic effects of microcystin-LR in human and chicken peripheral blood lymphocytes *in vitro*. *Toxicology* 2004;204:23-40. doi: 10.1016/j.tox.2004.05.016
21. Willcox KJ, Ash SL, Catignani GL. Antioxidants and prevention of chronic disease. *Crit Rev Food Sci Nutr* 2004;44:275-95. doi: 10.1080/10408690490468489
22. Anderson D, Yu TW, Phillips BJ, Schmezer P. The effect of various antioxidants and other modifying agents on oxygen-radical-generated DNA damage in human lymphocytes in the Comet assay. *Mutat Res-Fund Mol M* 1994;307:261-71. doi: 10.1016/0027-5107(94)90300-X
23. Cemeli E, Schmid TE, Anderson D. Modulation by flavonoids of DNA damage induced by estrogen-like compounds. *Environ Mol Mutagen* 2004;44:420-6. doi: 10.1002/em.20071
24. Živković L, Borozan S, Čabarkapa A, Topalović D, Ciptasari U, Bajić V, Spremo-Potparević B. Antigenotoxic properties

- of *Agaricus blazei* against hydrogen peroxide in human peripheral blood cells. *Oxid Med Cell Longev* 2017;2017:8759764. doi: 10.1155/2017/8759764
25. Singh NP, McCoy MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp Cell Res* 1988;175:184-91. doi: 10.1016/0014-4827(88)90265-0
26. Singh NP. Apoptosis assessment by the DNA diffusion assay. In: Blumenthal RD, editors. *Chemosensitivity: Volume II. In vivo models, imaging, and molecular regulators*. Totowa (NJ): Humana Press Inc.; 2005. p. 55-67.
27. Aherne SA, O'Brien NM. Mechanism of protection by the flavonoids, quercetin and rutin, against *tert*-butylhydroperoxide- and menadione-induced DNA single strand breaks in Caco-2 cells. *Free Radical Bio Med* 2000;29:507-14. doi: 10.1016/S0891-5849(00)00360-9
28. Kadrabova J, Krajcovicova-Kudlackova M, Madaric A, Valachovicova M, Mislanova C, Korenovska M. Protective properties of complex of quercetin, selenium, catechins and curcumin against DNA damage. *Oxid Antioxid Med Sci* 2012;1:179-84. doi: 10.5455/oams.180912.or.018
29. Myhrstad MCW, Carlsen H, Nordström O, Blomhoff R, Moskaug JØ. Flavonoids increase the intracellular glutathione level by transactivation of the γ -glutamylcysteine synthetase catalytical subunit promoter. *Free Radical Bio Med* 2002;32:386-93. doi: 10.1016/S0891-5849(01)00812-7
30. Wilms LC, Hollman PC, Boots AW, Kleinjans JC. Protection by quercetin and quercetin-rich fruit juice against induction of oxidative DNA damage and formation of BPDE-DNA adducts in human lymphocytes. *Mutat Res* 2005;582:155-62. doi: 10.1016/j.mrgentox.2005.01.006
31. Cemeli E, Baumgartner A, Anderson D. Antioxidants and the Comet assay. *Mutat Res* 2009;681:51-67. doi: 10.1016/j.mrrrev.2008.05.002
32. Persad S, Takeda S, Panagia V, Dhalla NS. β -adrenoceptor-linked signal transduction in ischemic-reperfused heart and scavenging of oxyradicals. *J Mol Cell Cardiol* 1997;29:545-58. doi: 10.1006/jmcc.1996.0298
33. Chiamonte R, Bartolini E, Riso P, Calzavara E, Erba D, Testolin G, Comi P, Sherbet GV. Oxidative stress signalling in the apoptosis of Jurkat T-lymphocytes. *J Cell Biochem* 2001;82:437-44. doi: 10.1002/jcb.1158
34. Benhusein GM, Mutch E, Aburawi S, Williams FM. Genotoxic effect induced by hydrogen peroxide in human hepatoma cells using comet assay. *Libyan J Med* 2010;5:1-6. doi: 10.3402/ljm.v5i0.4637
35. Imlay JA, Linn S. DNA damage and oxygen radical toxicity. *Science* 1988;240:1302-9. doi: 10.1126/science.3287616
36. Đelić N. Mechanisms of genotoxic effects of hormones. *Genetika* 2002;34:59-71.
37. Collins A, Ai-guo M, Duthie S. The kinetics of repair of oxidative DNA damage (strand breaks and oxidised pyrimidines) in human cells. *Mutat Res* 1995;336:69-77. doi: 10.1016/0921-8777(94)00043-6
38. Parshad R, Price FM, Bohr VA, Cowans KH, Zujewski JA, Sanford KK. Deficient DNA repair capacity, a predisposing factor in breast cancer. *Brit J Cancer* 1996;74:1-5. PMID: PMC2074608
39. Forlenza MJ, Latimer JJ, Baum A. The effects of stress on DNA repair capacity. *Psychol Health* 2000;15:881-91. doi: 10.1080/08870440008405589

Ocjena oštećenja DNA uzrokovanog adrenalinom u stanicama pune krvi pomoću komet-testa

Štetni učinci povišenih razina katekolamina posredovani su različitim mehanizmima, uključujući transkripciju gena i formiranje produkata oksidacije. Svrha je ovoga istraživanja bila utvrditi jesu li molekularni mehanizmi koji stoje u osnovi štetnoga djelovanja adrenalina na DNA posredovani reaktivnim vrstama kisika (ROS). Da bismo to postigli, izložili smo humane stanice pune krvi adrenalinu ($10 \mu\text{mol L}^{-1}$) i vodikovu peroksidu ($50 \mu\text{mol L}^{-1}$) (pozitivna kontrola), koje su odvojeno predtretirane i post-tretirane kvercetinom ($500 \mu\text{mol L}^{-1}$), „hvatačem“ slobodnih radikala. Kvercetin je značajno smanjio broj stanica s DNA oštećenjima i u predtretmanu i u post-tretmanu, što je sugeriralo da adrenalin djeluje uglavnom stvaranjem ROSa. Taj je mehanizam podržan i postupnim smanjenjem broja stanica s oštećenjima DNA izazvanim adrenalinom i H_2O_2 u vremenskim intervalima od 15, 30, 45 i 60 minuta nakon tretmana. Naši rezultati jasno pokazuju da su mehanizmi popravka prilično učinkoviti u odnosu na ROS-posredovana oštećenja DNA izazvana adrenalinom.

KLJUČNE RIJEČI: *genotoksičnost; kvercetin; popravak DNA; ROS; vodikov peroksid*