

International Journal of DAGENE

# Danubian Animal Genetic Resources

**Volume 8, Issue 1 (2023)**

DAGENE  
International Association for the Conservation  
of Animal Breeds in the Danube Region  
1078 Budapest, István street 2.  
Hungary



Danubian Animal Genetic Resources  
Volume 8, Issue 1 (2023)

Person responsible for edition:  
Editor-in-Chief: András GÁSPÁRDY (Hungary)

Editorial board:

President: Ante IVANKOVIĆ (Croatia)

Pál HAJAS (Hungary)

Beate BERGER (Austria)

Marcel MATIUTI (Romania)

Zsolt BECSKEI (Serbia)

Peter CHRENEK (Slovakia)

János POSTA (Hungary)

Technical editor: Jelena RAMLJAK (Croatia)

Editorial office: DAGENE - International Association for the Conservation of Animal Breeds in the Danube Region, 1078 Budapest, István street 2.  
[office@dagene.eu](mailto:office@dagene.eu) - [www.dagene.eu](http://www.dagene.eu)

Publisher: DAGENE - International Association for the Conservation of Animal Breeds in the Danube Region, 1078 Budapest, István street 2. [office@dagene.eu](mailto:office@dagene.eu) - [www.dagene.eu](http://www.dagene.eu)

Person responsible for publishing: András GÁSPÁRDY president of DAGENE

HU ISSN: 2498-5910

Printed by CENTER-PRINT Nyomda Ltd., Debrecen

This journal, founded 2016 is the official publication medium for the yearly activity of DAGENE members. The profile of the journal, reflecting the Association's mission, is to research, evaluate, present and document the conservation of endangered domestic animal breeds and their frontier areas in a scientific manner. The journal is mainly waiting for manuscripts on breed preservation in the Danube Valley from members of the Association, but also welcomes manuscripts from other dedicated authors and PhD students from other regions of the world, in English. Manuscripts should be submitted to [aivankovic@agr.hr](mailto:aivankovic@agr.hr).

Our journal is freely distributed in hard copy among the members of DAGENE and cooperating associations, however it is available from URL address of the journal: <http://ojs3.mtak.hu/index.php/dagr>

Supporting and advertising are possible at the office. Persons interested in becoming members of the DAGENE should contact the office.



This publication was supported by the Hungarian Ministry of Agriculture (BGMF/264-1/2023)

DOI: <https://doi.org/10.59913/dagr.2023.12257>

## Looking for added value of milk products of autochthonous sheep breeds

BECSKEI, Zsolt<sup>1\*</sup> – PASKAŠ, Snežana<sup>2</sup> – SAVIĆ, Mila<sup>2</sup> – TARIĆ, Elmin<sup>2</sup> – GÁSPÁRDY, András<sup>3</sup> – PAJIC, Marija<sup>4</sup> – RAŠETA, Mladen<sup>5</sup> – LAZAROVA, Mariya<sup>6</sup>

<sup>1</sup>Department of Animal Breeding and Genetics, Faculty of Veterinary Medicine University of Belgrade, Belgrade, Serbia

<sup>2</sup>Department of Animal Science, Faculty of Agriculture University of Novi Sad, Novi Sad, Serbia

<sup>3</sup>Department of Animal Breeding and Genetics, University of Veterinary Medicine, Budapest, Hungary

<sup>4</sup>Department of Veterinary Medicine, Faculty of Agriculture, University of Novi Sad, Novi Sad, Serbia

<sup>5</sup>Institute of Meat Hygiene and Technology, Belgrade, Serbia

<sup>6</sup>Department of Animal Husbandry - Ruminants and Dairy Farming, Faculty of Agriculture, Trakia University, Stara Zagora, Bulgaria

\*corresponding author: [beckezolt@gmail.com](mailto:beckezolt@gmail.com)

### Abstract

In order to determine the added value of the product, characterization of sheep milk and cheese samples from Sjenica sheep were assessed and compared to British milk sheep. The analyses were carried out according to standard chemical procedures. A comparative study showed that the milk from the Sjenica sheep was higher in fat and protein and had a lower casein/fat ratio. It also had higher total and somatic cell counts than British milk sheep. Sjenica cheese had significantly higher dry matter and fat content. However, both cheeses were classified as soft full-fat cheese with brine. The study showed that sheep's milk contains many essential components for cheesemaking and distinguishes sheep's cheese from other types of cheese, but standardisation of cheese production is still needed. To preserve the autochthonous sheep breeds of Serbia, promoting cheese production with added value is crucial and together with increased production could contribute to the sustainable development of rural areas.

Keywords: sheep, autochthonous breed, added value product, sheep cheese

## Introduction

There is a growing awareness of the need for high levels of biodiversity and its important role in improving the sustainability of low-input dairy production systems (POPOVIĆ-VRANJEŠ et al., 2018). Protecting local products means not only preserving the different local ecosystems at different levels: animals and plants (breeds and local varieties), but also microbial ecosystems, landscapes and dairy products (BERARD and MARCHENAY, 2006). Although many laws and measures have been adopted in Serbia for the protection of autochthonous breeds and genetic resources, many of them are still at risk of complete disappearance (POPOVIĆ-VRANJEŠ, 2018). Due to its compositional characteristics, sheep's milk is well suited for cheese production (POPOVIĆ-VRANJEŠ et al., 2017), and in Serbia sheep's milk is mainly used for cheese production. Autochthonous breeds do not produce large volumes of milk, but they have good quality milk with a desirable protein-fat ratio and small, naturally homogenised fat globules (POPOVIĆ-VRANJEŠ, 2018). The average size of fat globules is therefore the smallest ( $< 3.5 \mu\text{m}$ ) in sheep milk (PARK et al., 2007). Milk quality depends on many factors, and each component of milk influences cheese yield and quality (POPOVIĆ-VRANJEŠ et al., 2018). SOBRINO et al. (2018) found that, except for the casein/fat ratio, the season was the greatest source of variation in bulk tank milk characteristics of Manchega sheep. The relative stability of the C/F ratio made the sheep's milk more standardised, which is necessary for cheese production. Conversely, the farming system (semi-intensive and intensive) did not affect the chemical composition and physicochemical properties of sheep's milk in the study by KASAPIDOU et al. (2021).

Production of sheep's cheese in Serbia is usually at the farm level or in small local dairy units, with only about 370 tonnes of sheep's cheese produced in 2016 (FILIPOVIĆ, 2019). Traditionally associated with high-quality products such as lamb and Sjenica cheese, the Sjenica Pester plateau is characterised by a specific floristic composition of pastures (SAVIĆ et al., 2017). Sjenica sheep cheese is the best-known cheese from this area and is a protected designation of origin (PDO) product (FILIPOVIĆ, 2019). Sjenica cheese is made from either cow's or sheep's milk and belongs to the group of white cheeses. Producers of this cheese face a number of problems. The main challenges for the export of Sjenica cheese are the insufficient certification of this food product for the European Union market and the lack of a standardised production process, which results in variations in the quality of the product. Although white brine cheese is the most consumed type of cheese in Serbia (PASKAŠ et al., 2020), there is a lack of awareness of Sjenica sheep's cheese among regional consumers due to insufficient communication efforts (FILIPOVIĆ, 2019).

DOI: <https://doi.org/10.59913/dagr.2023.12257>

This study aimed to investigate the influence of different sheep breeds on the chemical composition and hygienic quality of milk, as well as its suitability for cheese production. A study of the chemical composition of white brine cheese was also carried out.

## **Material and methods**

### *Milk sampling and analysis*

From March to October 2022, samples of raw sheep's milk (n=16; n1=8; n2=8;) were collected from local farms. The farms differed in the breed of sheep reared. On the first farm Sjenica sheep was reared and on the second farm British milk sheep. On each farm, bulk milk samples were collected once a month (at regular intervals) and analysed on the same day. Both flocks were milked by hand twice a day and samples were taken after the morning milking. The standard analysis of the chemical composition of the milk was carried out with the MilkoScan FT+ analyser using the FTIR technique. The MilkoScan™ FT+ techniques comply with the ISO 9622: ISO 9622/IDF 141:2020 standard and the official AOAC method 972.16. FossomaticFT FC (ISO 13366-2:2008) and BactoScan™ FC+ (ISO 16297:2020) were used to determine the somatic cell count and total bacterial count of raw milk.

### *Cheese sampling and analysis*

Cheese samples (n=10; n1=5; n2=5) were collected from the same producers and analysed for chemical composition. Dry matter was determined using a standard method for measuring weight loss after drying (AOAC 926.08-1927). Protein was determined by the Kjeldahl-Van Slyke method for total N (AOAC 2001.14) using a Kjeltec Auto Analyzer and multiplying by 6.38. Fat content was determined according to IDF (2008) and ash was determined by dry ashing at 550°C (AOAC 935.42). Phosphorus was determined using a spectrophotometer (PG Instruments, type T80+) (AOAC 995.11), while flame photometry (Sherwood, type M410) was used for Ca and Na. The NaCl content was determined by the Volhard method (FIL-IDF, 1988).

### *Statistical analysis*

Results were statistically analysed and presented as mean, standard deviation, range and coefficient of variation. A t-test at  $p < 0.05$  was used to determine a significant difference between the means. Statistical analysis was performed using Statistica 10 (StatSoft STATISTICA 10.0).

## Results and discussion

Table 1 shows the chemical characteristics and hygiene parameters of the sheep milk samples. There was a statistically significant difference in the average fat content between milk samples from two different sheep breeds. The total protein content of Sjenica sheep milk was 5.58 and the fat content was 7.23g/100g. These results are in line with the range previously reported for Sjenica sheep milk (SAVIĆ et al., 2017). Compared to the results of SAVIĆ et al. (2014), the fat content was higher, but the protein and dry matter content were lower. There were no significant differences ( $p < 0.05$ ) in the casein/fat ratio between milk samples from the two different breeds. Both breeds showed a higher variation in the ash content and in the C/F ratio, which can negatively affect cheese production. The results related to the hygienic parameters of milk SCC and TBC showed that the milk from both breeds was of good quality. No difference was found in the hygienic parameters (SCC and TBC), but the results of the milk samples varied greatly in milk from both breeds. Monitoring SCC is a good tool for assessing the hygienic and sanitary quality of milk, but in sheep's milk, it is necessary to consider non-pathological factors that cause large variations in SCC and are different from those in cow's milk (RAYNAL-LJUTOVAC et al., 2007). This study shows the opposite results to those of TONAMO et al. (2020) for the SCC count of British milk sheep. They also differed from the SCC and TBC results in Lacaune sheep milk (MERILIN JUNIOR et al. 2015). SCC in milk affects cheese yield and cheesemaking performance. Milk with a low SCC can yield more than 4% more protein in cheese production than milk with a high SCC (RAYNAL-LJUTOVAC et al., 2007).

Table 1. Physicochemical composition of milk from two different breeds of sheep

Parameters (g/100g)	SJENICA SHEEP MILK			BRITISH MILK SHEEP MILK		
	$\bar{x} \pm SD$	range	CV (%)	$\bar{x} \pm SD$	range	CV (%)
Dry matter	17.89±1.06	16.54-19.77	5.93	17.26±0.73	16.09-18.66	4.26
SNF	10.64±0.58	9.79-11.5	5.42	10.99±0.61	10.21-12.20	5.58
Fat	7.23 <sup>a</sup> ±0.69	6.57-8.5	9.49	6.34 <sup>b</sup> ±0.34	5.88-7.0	5.31
Protein	5.58±0.49	5.13-6.7	8.80	5.50±0.61	4.33-6.38	11.09
Lactose	4.05±0.37	3.48-4.52	9.03	4.28±0.43	3.6-4.88	10.11
Casein	4.44±0.41	4.06-5.37	9.22	4.47±0.57	3.40-5.16	12.78

DOI: <https://doi.org/10.59913/dagr.2023.12257>

Ash	1.03±0.22	0.82-1.46	21.56	1.10±0.19	0.83-1.38	17.56
C/F	0.62±0.07	0.49-0.72	11.71	0.70±0.10	0.46-0.82	14.73
SCC (*1000/ml)	1499.62±45 7.82	872-2031	30.53	1384.62±484.59	465-2065	34.78
TBC (CFU*1000/ml)	224.87±101 .15	124-392	44.98	129.12±76.09	35-240	58.93

SNF-solids non-fat; C/F-casein/fat; SCC-somatic cells count; TBC-total bacteria count;  $\bar{x} \pm \text{sd}$ -arithmetic mean + standard deviation; cv-coefficient of variation; Means with different letters are significantly different ( $p < 0.05$ )

Table 2. Chemical composition of white-brined cheese from two different breeds of sheep

Parameters	SJENICA SHEEP CHEESE			BRITISH MILK SHEEP CHEESE		
	$\bar{x} \pm \text{SD}$	range	CV (%)	$\bar{x} \pm \text{SD}$	range	CV (%)
Dry matter (g/100g)	45.77a±1.10	44.80-47.06	2.41	43.77±0.83	42.80b-44.90	1.89
Fat (%)	24.85 <sup>a</sup> ±0.73	23.89-25.74	2.93	23.71 <sup>b</sup> ±0.74	22.98-24.68	3.10
Protein (%)	15.45±0.77	14.63-16.25	4.98	14.45±0.86	13.38-15.32	5.95
Total ash (%)	2.35±0.11	2.21-2.51	4.76	2.42±0.12	2.23-2.56	5.18
FDM	54.28±0.72	53.33-55.23	1.33	54.18±1.25	52.62-55.80	2.30
MFFB	72.15 <sup>a</sup> ±0.90	71.10-73.08	1.25	73.71 <sup>b</sup> ±0.80	72.75-74.80	1.08
NaCl (%)	3.34 <sup>a</sup> ±0.40	2.85-3.74	11.88	2.75 <sup>b</sup> ±0.32	2.22-3.01	11.83
S/M (%)	6.17 <sup>a</sup> ±0.80	5.18-7.06	12.95	4.88 <sup>b</sup> ±0.55	4.03-5.34	11.24
Ca (%)	0.38±0.07	0.31-0.48	18.27	0.43±0.06	0.37-0.51	13.34
P (%)	0.24±0.04	0.19-0.28	15.45	0.26±0.03	0.22-0.29	10.24

FDM-fat on a dry matter basis; PDM- a protein on a dry matter basis; MFFB-moisture on a fat-free basis; S/M (%)-salt in moisture;  $\bar{x} \pm \text{SD}$ -arithmetic mean + standard deviation; cv-coefficient of variation; Means with different letters are significantly different ( $p < 0.05$ ).

Table 2 shows the chemical composition of white-brined cheese from the two different breeds. It was found that dry matter and fat were significantly lower in British sheep's cheese. In a study by JANDRIĆ and SAVIĆ (2019), after 30 days of ripening, Sjenica sheep's milk cheese contained an average of 28.31 g/100 g of milk

DOI: <https://doi.org/10.59913/dagr.2023.12257>

fat, while at the beginning of the process, it contained 23.43 g/100 g. The research results for Sjenica cheese were within the mentioned ranges. A statistically significant difference was also found for NaCl content, MFFB and S/M ratio. The salt content of the cheese depends on the salt content of the brine. As the salt concentration in the brine increased, the salt content in the cheese also increased, as did the total ash, fat and protein content (RAHIMI et al., 2013). No difference was found in protein and FDM content, although the milk was not standardised. The dynamics of fat in dry matter showed a tendency of a balanced increase throughout the ripening period, from 49.44% to 52.72% in Sjenica sheep cheese (JANDRIĆ and SAVIĆ, 2019). The values obtained in this study were higher than those mentioned for both cheeses. According to the FDM content, the cheeses were classified as full-fat cheeses, and according to the MFFB, they belonged to the group of soft cheeses (SERBIAN REGULATIONS, 2014). The Ca and P contents of the two sheep cheeses were the most variable components. Ca and P directly influence cheese texture by participating in the protein network that forms the structural matrix of cheese (RAHIMI et al., 2013). Therefore, it was expected that the effect of different Ca and P levels on cheese texture would also vary, which could have an impact on cheese sensory properties.

## **Conclusion and recommendation**

The results of this study showed the favourable composition and properties of Sjenica sheep's milk, with significantly higher fat and dry matter contents compared to British milk sheep's milk. The results also confirmed the good quality of Sjenica white cheese. By adhering to higher standards and good manufacturing practices, sheep's milk and cheese can be produced safely without compromising quality. In particular, the production of Sjenica cheese could help to preserve the genetic diversity of domestic animals. To conserve local species and breeds, more breed-level data is needed and a knowledge base on livestock biodiversity needs to be developed.

To preserve the autochthonous sheep breeds of Serbia, promoting cheese production with added value is crucial and together with increased production could contribute to the sustainable development of rural areas.

## **Acknowledgement:**

The study was supported by the Ministry of Science, Technological Development and Innovation of the Republic of Serbia (Contract number 451-03-47/2023-01/200143).



DOI: <https://doi.org/10.59913/dagr.2023.12257>

## References

- BERARD, L. – MARCHENAY, P. (2006): Local products and geographical indications: taking into account of local knowledge and biodiversity. *Cultural Diversity and Biodiversity. International Social Science Journal*, 187:109-116. <https://doi.org/10.1111/j.1468-2451.2006.00592.x>
- FILIPOVIĆ, J. (2019): Market-Oriented Sustainability of Sjenica Sheep Cheese. *Sustainability* 11, 834 <http://doi.org/10.3390/su11030834>
- IDF (2008). International IDF Standard 222:2008 (Cheese and processed cheese products. Determination of Fat Content-Van Gulik method). IDF, Brussels, Belgium.
- JANDRIĆ, M. – SAVIĆ, Ž. (2019): Influence of milk on the content and changes of milk fat during production of Sjenički white cheese. *Agriculture and Forestry* 65(4), 175-182. <https://doi.org/10.17707/AgricultForest.65.4.15>
- KASAPIDOU, E. – BASDAGIANNI, Z. – PAPADOPOULOS, V. – KARAIKOU, C. – KESIDIS, A. – TSIOTSIAS, A. (2021): Effects of Intensive and Semi-Intensive Production on Sheep Milk Chemical Composition, Physicochemical Characteristics, Fatty Acid Profile, and Nutritional Indices. *Animals* 11, 2578. <https://doi.org/10.3390/ani11092578>
- MERLIN JUNIOR, A.I. – COSTA, G.R. – DOS SANTOS, S.J. – LUDOVICO, A. (2015): Sheep milk: Physical-chemical characteristics and microbiological quality. *Archivoslatino americanos de nutricion*. 65, 3, 193-198.
- PARK, Y.W.M. – JUÁREZ, M. – RAMOS, M. – HAENLEIN, G.F.W. (2007): Physico-chemical characteristics of goat and sheep milk, *Small Ruminant Research* 68, (1-2), 88-113. <https://doi.org/10.1016/j.smallrumres.2006.09.013>.
- PASKAŠ, S. – MIOČINOVIĆ, J. – LOPIČIĆ-VASIĆ, T. – MUGOŠA, I. – PAJIĆ, M. – BECSKEI, Z. (2020): Consumer attitudes towards goat milk and milk products in Vojvodina. *Mjekarstvo* 70 (3), 171-183. <https://doi.org/10.15567/mljekarstvo.2020.0304>
- POPOVIĆ-VRANJEŠ, A. – PASKAŠ, S. – KRSTOVIĆ, S. – JURAKIĆ, Ž. – JEŠIĆ, G. – STRUGAR, K. (2017): The production of hard cheese with added value from raw ewe's milk. *International Symposium on Animal Science (ISAS) 2017*. 05th - 10th June 2017, Herceg Novi, Montenegro. 211-219.
- POPOVIĆ-VRANJEŠ, A. – PASKAŠ, S. – BECSKEI, Z. – JEVTIĆ, M. – KRSTOVIĆ, S. (2018): The impact of organic agriculture on biodiversity conservation and milk quality for traditional cheese production. *Contemporary Agriculture* 67 (1), 58-64. <https://doi.org/10.2478/contagri-2018-0009>
- RAHIMI, J. – KHOSROU SHAHI, A. – MORADI, M.M. – MOHAMADI, H. – ABBASI, H. – MADADLOU, A. (2013): Texture and chemistry of Iranian white cheese as influenced by brine treatments. *Journal of Food Processing and Technology* 4:4. <https://doi.org/10.4172/2157-7110.1000219>
- RAYNAL-LJUTOVAC, K. – PIRISI, A. – de CREMOUX, R.C. – GONZALO.C. (2007): Somatic cells of goat and sheep milk: Analytical, sanitary, productive and

DOI: <https://doi.org/10.59913/dagr.2023.12257>

technological aspects Small Ruminant Research 68, 126–144.

<https://doi.org/10.1016/j.smallrumres.2006.09.012> x

SAVIĆ, Ž. – RADOVIĆ, B. – MILENKOVIĆ, M. – LALIĆ, N. (2014): Kvalitet mleka i osobine surutke dobijene pri proizvodnji Sjeničkog sira u industrijskim uslovima. XIX SAVETOVANJE O BIOTEHNOLOGIJI” Zbornikradova, 19 (21).

SAVIĆ, M. – POPOVIĆ-VRANJEŠ A. – VUČKOVIĆ, S. – BECSKEI, Zs. – RELIĆ, R. – DIMITRIJEVIĆ, B. – TRAILOVIĆ, R. – PRODANOVIĆ, S. – DIMITRIJEVIĆ, V. (2017): Importance of Sjenica Zackel sheep for sustainable development at Sjenica Pester plateau. International Symposium on Animal Science (ISAS) 2017. 05th - 10th June 2017, Herceg Novi, Montenegro. 115-121.

SERBIAN REGULATION (2014): Ordinance on quality of dairy products and starter cultures. Official Gazette of the Republic of Serbia, No. 34/2014

SOBRINO, J. L. – COLADO, P.M.J. – SIGLER, G.I.A. – MARÍN, M.L.A. – SÁNCHEZ, N.N. – ASENSIO, R.J. – PALOMARES, P.G.D.M. – SÁNCHEZ, A.R. (2018) Composition and colour indices of sheep’s bulk-tank milk are influenced by production practices, Italian Journal of Animal Science 17:2, 477-488, <https://doi.org/10.1080/1828051X.2017.1383860>

TONAMO, A. – KOMLÓSI, I. – VARGA, L. – CZEGLÉDI, L. – PELES, F. (2020): Bacteriological Quality of Raw Ovine Milk from Different Sheep Farms. Animals 10,1163. <https://doi.org/10.3390/ani10071163>