

Changes in *Pseudomonas* sp. CY growth in the presence of atrazine

Suzana Đedović¹, Monika Stojanova², Jovan Bojkovski³, Igor Kljujev⁴, Vera Karličić⁴, Blažo Lalević^{4*}, Vera Raičević⁴

¹ Institute of Pesticides and Environmental Protection, Banatska 31b, 11080 Zemun, Serbia

² Open Science, Association for scientific research, educational and cultural activities, Jane Sandanski 61/16, 6000 Ohrid, North Macedonia

³ University of Belgrade, Faculty of Veterinary Medicine, 11000 Belgrade, Serbia

⁴ University of Belgrade, Faculty of Agriculture, Nemanjina 6, 11080 Zemun, Serbia

* Corresponding author: blazol@agrif.bg.ac.rs

Abstract

Microbial degradation, compared with many other degradation processes, is the most important pathway for the depletion of triazine herbicides in soil. The aim of this study was to determine the growth potential of *Pseudomonas* sp. CY in the presence of atrazine and additional carbon (sodium citrate) and nitrogen (ammonium-nitrate) sources. The experiment was performed with five treatments: i) 100 mg/L atrazine (control); ii) One hundred mg/L atrazine + sodium citrate (0.3 %, w/v); iii) One hundred mg/L atrazine + sodium citrate (0.3 %, w/v) + ammonium nitrate (0.6 %, w/v); iv) Atrazine (300 mg/L) + sodium citrate (0.3 %, w/v) and v) Atrazine (500 mg/L) + sodium citrate (0.3 %, w/v). The bacterial count was determined after incubation (7 days at 30°C) using the agar plate method, while atrazine degradation was determined by measuring the optical density at 221 nm. *Pseudomonas* sp. CY can partially utilize atrazine as the sole source of carbon and energy. The highest values of the bacterial count were determined at the highest initial atrazine concentrations; however, bacterial growth was not detected in these treatments. A significant impact of citrate on bacterial growth and atrazine degradation was observed, while the addition of nitrate decreased the atrazine degradation rate. This study confirmed that *Pseudomonas* sp. CY can be used as a prominent candidate for the remediation of atrazine-affected environments.

Key words: ammonium-nitrate, atrazine, bacterial growth, *Pseudomonas* sp., sodium-citrate

Introduction

The use of pesticides is strongly recommended in contemporary agricultural production. One of the pesticides that has been frequently applied during the last decades of the 20th century is atrazine (C₈H₁₄ClN₅). However, their widespread use has led to soil contamination (Duan et al., 2016). Atrazine is one of the most frequently detected pesticides in various agricultural soils. Furthermore, atrazine has phytotoxic (Wang et al., 2015), carcinogenic, and teratogenic (Sun et al., 2020) effects. Atrazine was also detected in surface and subsurface waters, although its use was prohibited by the EU in 2003, and Amadori et al. (2016) reported the widespread use of atrazine worldwide. Considering the half-life of atrazine, which varies from one month to approximately a year (Diana et

al., 2000), it is important to find an environmentally friendly and low-cost approach for the depletion of atrazine in the environment.

The ability of several bacterial genera to degrade atrazine has been well described (Bazhanov et al., 2016; Zhu et al., 2019). Nevertheless, in most cases, the low efficiency of microbial strains to degrade atrazine has been observed (Solomon et al., 2013). In addition, owing to the lack of microbe-based products for atrazine removal from the environment (Zhu et al., 2019), it is necessary to find more efficient microbial strains capable of cleaning atrazine-affected ecosystems. In this study, a pure culture of *Pseudomonas* sp. strain CY was used to estimate growth estimation in the presence of atrazine. This strain showed the ability to grow on benzene (Lalević et al., 2006).

Apart from degradation *via* direct metabolism, microbial strains are capable of the co-metabolic degradation of atrazine (Cupul et al., 2014). *In vitro* experiments have shown that the addition of other energy and nutrient source(s) may stimulate microbial activity (Ngigi et al., 2013). Although additional C sources stimulate atrazine mineralization (Dehghani et al., 2013), inhibition of this process after the addition of external N sources has been observed in other studies (Clausen et al., 2002). In these studies, various external C and N sources were used: acetate, pectin (Rhine et al., 2003), and sucrose (Wang et al., 2011) as C sources, whereas NH_4NO_3 , cyanuric acid (Rhine et al., 2003), nitrate, urea (Bichat et al., 1999) and organic amendments (Forouzangohar et al., 2005) were used as N sources. Hence, it is evident that additional nutrient sources may play a crucial role in the microbial mineralization of atrazine (Struthers et al., 1998).

The objective of this study was to determine the effect of secondary nutrient sources (sodium citrate and ammonium nitrate) on the growth of *Pseudomonas* sp. CY in the presence of atrazine to estimate the efficiency of atrazine biodegradation under laboratory conditions.

Material and methods

Gram-negative *Pseudomonas* sp. CY, previously isolated from kerosene, was maintained in the Laboratory of Environmental Microbiology, Faculty of Agriculture, University of Belgrade, Serbia.

The liquid mineral salt medium (Mandelbaum et al., 1993) was inoculated with a suspension of pure cultures of *Pseudomonas* sp. CY (0.5 %, v/v). The initial population density of *Pseudomonas* sp. CY were determined on nutrient agar (Torlak, Serbia). Atrazine (diluted in methanol) and external C and N sources were added as nutrient sources for bacterial growth. The experiment was performed within five treatments: i. $100 \text{ mg} \times \text{l}^{-1}$ of atrazine (100 A); ii. One hundred $\text{mg} \times \text{l}^{-1}$ atrazine + 0.3 % (w/v) sodium citrate (100 AS); iii. One hundred $\text{mg} \times \text{l}^{-1}$ atrazine + 0.3 % (w/v) sodium citrate + 0.6 % (w/v) ammonium-nitrate (100 ASA); iv. $300 \text{ mg} \times \text{l}^{-1}$ atrazine + 0.3 % (w/v) sodium citrate (300 AS) and $500 \text{ mg} \times \text{l}^{-1}$ atrazine + 0.3 % (w/v) sodium citrate (500 AS). Incubation was performed in an

orbital shaker (Biosan ES-20, Latvia) for seven days at 30°C. All experiments were performed in triplicates.

Samples for the determination of bacterial count were taken after 24, 48, 72, and 96 h and at the end of the incubation period. Bacterial counts were determined using nutrient agar (Torlak, Serbia) and expressed as colony-forming units (CFU) per mL. The atrazine degradation was estimated using a spectrophotometer (T70 UV/VIS Spectrometer, PG Instruments Ltd., UK) at 221 nm (Moreira et al., 2017). Optical density (OD₂₂₁) readings were used to determine the atrazine concentration in the liquid cultures at the beginning of the experiment, and after 24, 48, 72, 96, 120 and 168 h of incubation.

The results were statistically analyzed using the software package SPSS 20. To determine the statistically significant differences in the obtained values, one-way analysis of variance (ANOVA) was performed, followed by post-hoc Tukey's test ($p = 0.05$) for the values between different treatments at the same time interval, as well as between the same treatments on different time intervals.

Results and discussion

The results showed that the bacterial count, optical density of liquid cultures, and atrazine degradation were affected by the concentrations of atrazine, external nutrient sources, and incubation time. In this study, the initial population density of *Pseudomonas* sp. CY was 3.8×10^7 CFU x mL⁻¹.

As can be seen from Table 1, in all samples a decrease in the bacterial count compared with the initial population count was observed. During incubation, the presence of atrazine in all treatments hindered bacterial growth during the first 48 h of incubation. Zhang et al. (2012) revealed that pollutants might be accumulated in the microbial cells and cause metabolic disorders. Furthermore, they reported that atrazine inhibits bacterial growth. In the control, the lowest bacterial count was observed compared to other treatments, which suggests that the addition of external nutrient sources accelerated bacterial growth, particularly in the 100 AS and 100 ASA treatments. The addition of external nutrients and energy sources does not inhibit the microbial performance (Kannika et al., 2010). Moreover, the complex structure of atrazine contributes to its low bacterial growth (Sharma et al., 2019).

Table 1. Bacterial count ($\times 10^6$ CFU/mL) in the liquid medium supplemented with atrazine and external nutrient sources

Treatment	Time (hours)					
	24	48	72	96	120	168
	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$
100 A	8.03 ^{aA} \pm	2.92 ^{bA} \pm	3.65 ^{cA} \pm	6.22 ^{dA} \pm	2.31 ^{eA} \pm	6.51 ^{fA} \pm
(control)	0.67	0.65	0.52	1.39	0.10	1.97
100 AS	6.37 ^{aB} \pm	3.07 ^{bB} \pm	4.28 ^{cB} \pm	5.15 ^{dB} \pm	12.38 ^{eB} \pm	5.42 ^{fB} \pm
	1.14	0.42	0.30	0.70	0.10	0.98
100 ASA	6.29 ^{aB} \pm	4.36 ^{bC} \pm	5.28 ^{cC} \pm	5.98 ^{dC} \pm	9.18 ^{eC} \pm	7.77 ^{fC} \pm
	0.56	0.12	0.18	0.89	0.15	2.19
300 AS	10.99 ^{aC} \pm	10.03 ^{bD} \pm	9.30 ^{cD} \pm	9.94 ^{dD} \pm	9.80 ^{eD} \pm	9.40 ^{fD} \pm
	1.22	1.48	0.10	0.20	2.62	1.99
500 AS	13.11 ^{aD} \pm	13.11 ^{aE} \pm	11.97 ^{bE} \pm	9.39 ^{cE} \pm	11.51 ^{dE} \pm	12.49 ^{eE} \pm
	1.11	1.02	0.96	3.58	0.12	0.16

a, b, c - Values of the same treatment at different time intervals marked with different letters, are significantly different ($p < 0.05$).

A, B, C - Values of different treatments at the same time interval, marked with different letters, are significantly different ($P < 0.05$).

Although the highest *Pseudomonas* sp. CY counts were detected at 300 AS and 500 AS, a negligible bacterial growth was observed after 72 and 96 h, respectively. According to these results, higher atrazine concentrations remarkably decreased bacterial growth, which has been previously confirmed (Zhu et al., 2019). On the other hand, treatments with citrate (100 AS), and citrate + nitrate (100 ASA) showed more pronounced bacterial growth compared to the control; in both treatments, *Pseudomonas* sp. CY stationary phase was reached between 120 and 168 h of incubation. The addition of nitrate (100 ASA) resulted in a significantly higher cell number compared to 100 AS.

Ramadan et al. (1990) pointed out that some microbes are more likely to use nutrients and energy sources than pollutants. Sharma et al. (2019) found that several N sources are a better substrates for bacterial growth than atrazine. However, remarkable shifts in *Pseudomonas* sp. counts were observed during incubation. These changes may be linked to atrazine degradation (Yale et al., 2017). In most samples, the highest OD₂₂₁ was recorded at the start of the incubation period (Table 2), which indicated that *Pseudomonas* sp. strain CY was able to use atrazine. Nevertheless, an irregular degradation trend was observed in all treatments, as previously reported (Alattas et al., 2023).

Table 2. Optical density (OD₂₂₁) of the liquid cultures in the presence of atrazine and external nutrient sources

Treatment	Time (hours)						
	0	24	48	72	96	120	168
	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$	$\bar{x} \pm SD$
100 A (control)	1.41 ^{aA} ± 0.08	1.38 ^{aA} ± 0.16	0.38 ^{bA} ± 0.06	0.51 ^{cA} ± 0.13	1.02 ^{dA} ± 0.34	0.35 ^{bA} ± 0.19	1.08 ^{eA} ± 0.50
100 AS	1.33 ^{aB} ± 0.10	1.05 ^{bB} ± 0.27	0.40 ^{cA} ± 0.10	0.64 ^{dB} ± 0.07	0.81 ^{eB} ± 0.17	2.13 ^{fB} ± 0.13	0.87 ^{gB} ± 0.24
100 ASA	1.34 ^{aB} ± 0.06	1.03 ^{bB} ± 0.13	0.65 ^{cB} ± 0.02	0.83 ^{dC} ± 0.04	0.97 ^{eC} ± 0.22	0.34 ^{fA} ± 0.25	1.33 ^{aC} ± 0.53
300 AS	2.33 ^{aC} ± 0.08	1.96 ^{bC} ± 0.29	1.28 ^{cC} ± 0.50	1.30 ^{dD} ± 0.59	1.75 ^{dD} ± 0.05	2.44 ^{eC} ± 0.03	1.65 ^{fD} ± 0.48
500 AS	2.46 ^{aD} ± 0.05	2.37 ^{bD} ± 0.27	2.32 ^{cD} ± 0.28	2.15 ^{dE} ± 0.23	1.64 ^{eE} ± 0.86	2.47 ^{aC} ± 0.04	2.25 ^{fE} ± 0.04

a, b, c - Values of the same treatment at different time intervals marked with different letters, are significantly different ($p < 0.05$).

A, B, C - Values of different treatments at the same time interval, marked with different letters, are significantly different ($P < 0.05$).

Significantly higher ($p < 0.05$) OD₂₂₁ values were observed in the 300 AS and 500 AS treatments than in the control and other treatments. At 300 AS and 500 AS, OD₂₂₁ declined by 72 h, and 96 h of incubation was followed by a rapid increase up to 120 h. According to the obtained OD values and comparison with other treatments, *Pseudomonas* sp. CY was unable to degrade atrazine at 300 and 500 AS. The prolonged lag phase and adaptation period detected in these treatments may be crucial for low bacterial growth and inefficient atrazine degradation (Zhao et al., 2018). In the other treatments, including the control, the CY strain showed a faster acclimation period. However, complete degradation of atrazine was not observed in these treatments. In our study, an incubation period of 168 h was used to determine the atrazine degradation. Mandelbaum et al. (1993) revealed that atrazine mineralization is a relatively slow process, whereas Silva et al. (2004) suggested that atrazine degradation should be estimated after an adaptation period of 28 days. In addition, Khatoun and Rai (2020) pointed out the importance of some environmental factors during atrazine biodegradation by microbes, which possess enzymes and/or genes responsible for the formation of degradation products (Liang et al., 2022).

The results presented in Table 2 show that in the control and 100 ASA, after 120 h of incubation, the lowest OD₂₂₁ values were observed, while in 100 AS the lowest values were detected after 48 h of incubation. Significant differences in OD₂₂₁ values among the control, 100 AS, and 100 ASA groups were observed during incubation. Although these data suggest that the addition of citrate plays a significant role in atrazine biodegradation (Mandelbaum et al. 1993), our results showed that, in most samples, the OD₂₂₁ values were significantly lower in the control. This observation was previously

reported (Kannika et al., 2010), pointing out that the addition of inorganic N decreased the atrazine degradation process, which is similar to our data. The increase in OD₂₂₁ values at the end of the incubation period and the irregular curve of atrazine biodegradation could be attributed to the degradation of metabolites, which may serve as an additional nutrient sources and accelerators of atrazine degradation (Piutti et al., 2002).

Our results show the absence of a correlation between bacterial growth and atrazine degradation. Zhang et al. (2019) reported that *Klebsiella variicola* FH-1 reached log phase after 12-16 h of incubation, but the atrazine degradation rate was very low. Furthermore, they suggested that atrazine is not an optimal nutrient source for bacterial growth, but its degradation rate depends on the presence of external N-source(s). Taking into account that the selection of external C and N sources (Deghani et al., 2013) and environmental factors (Qu et al., 2020) plays a crucial role in atrazine biodegradation, optimization of environmental factors, such as temperature and pH, and addition of various C and N sources, coupled with detection of degradation products, will be involved in future research of atrazine biodegradation by *Pseudomonas* sp. CY.

Conclusion

The present study demonstrated that atrazine can be degraded by *Pseudomonas* sp. CY. Significant shifts in the bacterial count and optical density were observed during incubation. Shifts in cell number were mainly caused by external C and N sources rather by the initial atrazine concentrations. The addition of citrate increased bacterial growth and atrazine degradation, suggesting that *Pseudomonas* sp. CY has promising potential for applications in atrazine-affected environments. Further research will be focused on the optimization of critical parameters in *in vitro* conditions and on the determination of the role of various C and N sources during *Pseudomonas* sp. CY-assisted atrazine degradation.

Acknowledgement: This research was partially supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant Nos. 451-03-47/2023-01/200214 and 451-03-47/2023-01/200116).

References

Alattas, SG, Zabermawi NM, El Bestawy E. 2023: Biodegradation of atrazine using selected marine bacteria: possibilities for treating pesticide-contaminated wastewater. *Journal of King Saud University – Science* 35:102721.

- Amadori MF, Rodrigues MB, Rebouças CC, Peralta-Zamora PG, Grassi MT, Abate G. 2016: Behavior of atrazine and its degradation products deethylatrazine and deisopropylatrazine in oxisol samples. *Water Air Soil Pollution* 227:380.
- Bazhanov DP, Li C, Li H, Li J, Zhang X, Chen X, Yang H. 2016: Occurrence, diversity and community structure of culturable atrazine degraders in industrial and agricultural soils exposed to the herbicide in Shandong Province, P.R. China. *BMC Microbiology* 16:265.
- Bichat F, Sims GK, Mulvaney RL. 1999: Microbial utilization of heterocyclic nitrogen from atrazine. *Soil Science Society of America Journal* 63:100-110.
- Clausen GB, Larsen L, Johnsen K, Radnoti de Liphay J, Aamand J. 2002: Quantification of the atrazine-degrading *Pseudomonas* sp. strain ADP in aquifer sediment by quantitative competitive polymerase chain reaction. *FEMS Microbiology Ecology* 41:221-229.
- Cupul WC, Abarca GH, Vázquez RR, Salmones D, Hernández RG, Gutiérrez EA. 2014: Response of ligninolytic macrofungi to the herbicide atrazine: doseresponse bioassays. *Revista Argentina de Microbiologia* 46:348-357.
- Dehghani M, Nasseri S, Hashemi H. 2013: Study of the bioremediation of atrazine under variable carbon and nitrogen sources by mixed bacterial consortium isolated from corn field soil in Fars province of Iran. *Journal of Environmental and Public Health* 2013:973165.
- Diana SRW, Schaeffer D, Beckmen K, Beasley V. 2000: Effects of atrazine on amphibian growth and survival in artificial aquatic communities. *Environmental Toxicology and Chemistry* 19:2961-2967.
- Duan Q, Lee J, Liu Y, Chen H, Hu H. 2016: Distribution of heavy metal pollution in surface soil samples in China: A graphical review. *Bulletin of Environmental Contamination and Toxicology* 97:303-309.
- Forouzangohar M, Haghnia G, Kocheiki A, Tabatabaie Yazdi F. 2005: The effect of organic matter and soil texture on degradation of atrazine and metamilon. *Journal of Science, Agriculture and Natural Resources* 9:131-141.
- Kannika S, Heepngoen P, Sadowsky MJ, Boonkerd N. 2010: *Arthrobacter* sp. strain KU001 isolated from a Thai soil degrades atrazine in the presence of inorganic nitrogen sources. *Journal of Microbiology and Biotechnology* 20:602-608.
- Khatoun H, Rai JPN. 2020. Optimization studies on biodegradation of atrazine by *Bacillus badius* ABP6 strain using response surface methodology. *Biotechnological Reports (Amst)* 26:00459.
- Lalević B, Jovanović Lj, Raičević V, Nikšić M, Kiković D, Marinković N. 2006: *Pseudomonas* sp. growth in the presence of benzene. *Zemljište i biljka* 55:179-185.
- Liang Y, Ding L, Song Q, Zhao B, Wang S, Liu S. 2022. Biodegradation of atrazine by three strains: identification, enzymes activities, and biodegradation mechanism. *Environmental Pollutants and Bioavailability* 34:549-563.

- Mandelbaum RT, Wackett LP, Allan DL. 1993: Mineralization of the *s*-triazine ring of atrazine by stable bacterial mixed cultures. *Applied and Environmental Microbiology* 59:1695-1701.
- Moreira AJ, Borges AC, Gouvea LFC, MacLeod TCO, Freschi GPG. 2017: The process of atrazine degradation, its mechanism, and the formation of metabolites using UV and UV/MW photolysis. *Journal of Photochemistry and Photobiology A: Chemistry*, 347:160-167.
- Ngigi AN, Getenga ZM, Dorfler U, Boga HI, Kuria B, Ndalut P, Schroll R. 2013: Effects of carbon amendment on in situ atrazine degradation and total microbial biomass. *Journal of Environmental Science and Health, Part B* 48:40-48.
- Piutti S, Hallet S, Rousseaux S, Philippot L, Soulas G, Martinlaurent F. 2002: Accelerated mineralisation of atrazine in maize rhizosphere soil. *Biology and Fertility of Soils* 36:434-441.
- Ramadan MA, El-Tayeb OM, Alexander M. 1990: Inoculum size as a factor limiting success of inoculation for biodegradation. *Applied and Environmental Microbiology* 56:1392-1396.
- Rhine ED, Fuhrmann JJ, Radosevich M. 2003: Microbial community responses to atrazine exposure and nutrient availability: linking degradation capacity to community structure. *Microbial Ecology* 46:145-160.
- Qu M, Liu G, Zhao J, Li H, Liu W, Yan Y, Feng X, Zhu D. 2020. Fate of atrazine and its relationship with environmental factors in distinctly different lake sediments associated with hydrophytes. *Environmental Pollution* 256:113371.
- Sharma A, Kalyani P, Trivedi VD, Kapley A, Phale PS. 2019: Nitrogen-dependent induction of atrazine degradation pathway in *Pseudomonas* sp. strain AKN5. *FEMS Microbiology Letters* 366:fny277.
- Silva E, Fialho AM, Sa-Correia I, Burns RG, Shaw LJ. 2004: Combined bioaugmentation and biostimulation to cleanup soil contaminated with high concentrations of atrazine. *Environmental Science and Technology* 38:632-637.
- Solomon RDJ, Kumar A, Santhi VS. 2013: Atrazine biodegradation efficiency, metabolite detection, and *trzD* gene expression by enrichment bacterial cultures from agricultural soil. *Journal of Zhejiang University-Science B* 14:1162-1172.
- Struthers JK, Jauachandran K, Moorman TB. 1998: Biodegradation of atrazine by *Agrobacterium radiobacter* JI4a and use of this strain in bioremediation of contaminated soil. *Applied and Environmental Microbiology* 64:3368-3375.
- Sun J, Ma XL, Wang W, Zhang J, Zhang H, Wang YJ. 2020: Adsorption characteristics of atrazine on different soils in the presence of Cd(II). *Adsorption Science and Technology* 38:225-239.
- Wang Q, Que X, Zheng R, Pang Z, Li C, Xiao B. 2015: Phytotoxicity assessment of atrazine on growth and physiology of three emergent plants. *Environmental Science and Pollution Research* 22:9646-9657.

- Wang J, Zhu L, Liu A, Ma T, Wang Q, Xie H, Wang J, Jiang T, Zhao R. 2011: Isolation and characterization of an *Arthrobacter* sp. strain HB-5 that transforms atrazine. *Environmental Geochemistry and Health* 33:259-266.
- Yale RL, Sapp M, Sinclair CJ, Moir JWB. 2017: Microbial changes linked to the accelerated degradation of the herbicide atrazine in a range of temperate soils. *Environmental Science and Pollution Research* 24:7359-7374.
- Zhang J, Liang S, Wang X, Lu Z, Sun P, Zhang H, Sun F. 2019: Biodegradation of atrazine by the novel *Klebsiella variicola* strain FH-1. *Biomed Research International* 2019:4756579.
- Zhang Y, Meng D, Wang Z, Guo H, Wang Y. 2012: Oxidative stress response in two representative bacteria exposed to atrazine. *FEMS Microbiology Letters* 334:95-101.
- Zhao X, Wang L, Ma F, Yang J. 2018: Characterisation of an efficient atrazine-degrading bacterium, *Arthrobacter* sp. ZXY-2: an attempt to lay the foundation for potential bioaugmentation applications. *Biotechnology for Biofuels* 11:113.
- Zhu J, Fu L, Jin C, Meng Z, Yang N. 2019: Study on the isolation of two atrazine-degrading bacteria and the development of a microbial agent. *Microorganisms* 7:80.

Promene u rastu *Pseudomonas* sp. CY u prisustvu atrazina

Suzana Đedović¹, Monika Stojanova², Jovan Bojkovski³, Igor Kljujev⁴, Vera Karličić⁴, Blažo Lalević^{4*}, Vera Raičević⁴

¹ Institute za pesticide i zaštitu životne sredine, Banatska 31b, 11080 Zemun, Srbija

² Open Science, Udruženje za naučno istraživanje, edukativne i kulturne aktivnosti, Jane Sandanski 61/16, 6000 Ohrid, Severna Makedonija

³ Univerzitet u Beogradu, Fakultet veterinarske medicine, 11000 Beograd, Srbija

⁴ Univerzitet u Beogradu, Poljoprivredni fakultet, Nemanjina 6, 11080 Zemun, Srbija

* Autor za korespondenciju: B. Lalević, Email: blazol@agrif.bg.ac.rs

SAŽETAK

Mikrobijalna degradacija je, u poređenju sa drugim degradacionim procesima, najvažniji način razgradnje triazinskih herbicida u zemljištu. Cilj ovog istraživanja je bio da se determinišu potencijal rasta *Pseudomonas* sp. CY u prisustvu atrazina i dopunskih izvora ugljenika (natrijum citrat) i azota (amonijum nitrat). Eksperiment je obavljen u pet tretmana: 1. 100 mg/l atrazina (kontrola); 2. 100 mg/l atrazina + natrijum citrat (0,3 %, w/v); 3. 100 mg/l atrazina + natrijum citrat (0,3 %, w/v) + amonijum nitrat (0,6 %, w/v); 4. 300 mg/l atrazina + natrijum citrat (0,3 %, w/v) i 5. 500 mg/l atrazina + natrijum citrat (0,3 %, w/v). Broj bakterija je određen nakon inkubacije (7 dana na 30°C) metodom agarnih ploča, dok je koncentracija atrazina određena merenjem optičke gustine na talasnoj dužini od 221 nm. *Pseudomonas* sp. CY parcijalno može da koristi atrazin kao jedinstveni izvor ugljenika i energije. Najveća brojnost bakterija je ustanovljena pri najvećim početnim koncentracijama atrazina; međutim, rast bakterija u ovim tretmanima nije detektovan. Konstatovan je značajan uticaj citrata na rast bakterija i degradaciju atrazina, dok je dodavanje nitrata rezultiralo smanjenjem stepena degradacije atrazina. Ova istraživanja potvrđuju da se *Pseudomonas* sp. CY može koristiti kao značajan činilac remedijacije životne sredine kontaminirane atrazinom.

Ključne reči: amonijum-nitrat, atrazin, rast bakterija, *Pseudomonas* sp., natrijum-citrat

Received 21.11.2023

Revised 10.12.2023

Accepted 13.12.2023