

NATURAL AND ANTHROPOGENIC RADIOACTIVITY OF FEEDSTUFFS, MOSSES AND SOIL IN THE BELGRADE ENVIRONMENT, SERBIA

SVETLANA GRDOVIĆ¹, GORDANA VITOROVIĆ¹, BRANISLAVA MITROVIĆ¹,
V. ANDRIĆ², B. PETRUJKIĆ¹ and M. OBRADOVIĆ¹

¹ Faculty of Veterinary Medicine, 11000 Belgrade, Serbia

² Vinča Institute for Nuclear Sciences, 11000 Belgrade, Serbia

Abstract – By gamma spectroscopic measurement a content of natural radio-nuclides (⁴⁰K, ²³⁸U, ²²⁶Ra, ²³²Th) and ¹³⁷Cs was determined in samples of soil, alfalfa, maize and moss on six sites in the surroundings of Belgrade. Natural radio-nuclides in the soil were at the level characteristic for Serbia, whereas a relatively high level of activity of ¹³⁷Cs (around 30 Bq kg⁻¹) was determined. On the other hand, in plant samples mostly used as feed (such as alfalfa and maize) the concentration of natural radio-nuclide activity and ¹³⁷Cs was relatively low, i.e. below the range of detection. The content of natural radio-nuclides in moss was within the standard range of values specific for Serbia. However, the activity level of ¹³⁷Cs in moss gathered from the wider area around Belgrade, was high, the highest measured level being in the Avala-Zuce area (158-221 Bq kg⁻¹). Our results show that this radio-nuclide is still present in the living environment of Belgrade even 20 years after the Chernobyl disaster, and that moss is a good indicator of living environment ¹³⁷Cs contamination.

Keywords: Belgrade environment, feedstuffs, mosses, radio-nuclides, soil

UDC 504.5(497.111)

INTRODUCTION

The increased use of nuclear energy worldwide, along with nuclear weapon probes, coal combustion, the production and use of phosphorus fertilizers, the mining industry and the formation of radioactive waste dumps, all contribute to the distribution of natural radioactivity. The quantity of radio-nuclides from the cited sources has significant impact on the environment (Pietrzak-Flis et al., 2000; Pantelić et al., 2004; Bikit et al., 2005). One of the important anthropogenic sources of environmental pollution by the primordial radio-nuclides ²²⁶Ra, ²³²Th, and ⁴⁰K are phosphorus fertilizers (Adel et al., 2005). Uranium, which is of anthropogenic origin, is often present in soil in a form which is easily accessible to plants, which offers a greater possibility for its incorporation in a food chain (Bolca et al., 2007).

The partial meltdown of the reactor at Chernobyl in April 1986 released high amounts of radionuclides

into environment. ¹³¹I, ¹³⁴Cs and ¹³⁷Cs were the most important and most dangerous radionuclides released and spread over a large part of Europe.

After the Chernobyl nuclear accident, Hecht (1988), Nifontova (1995), Delfanti et al., (1999), Dragović et al., (2004), Dragović et al., (2007) concluded that mosses, lichens and mushrooms were typical representatives of bioindicator plants, for both radiocesium and natural radio-nuclides presented in the environment.

Physical decay, weathering and other major factors have contributed to the significant decline of radionuclides in the environment. The level of the long-lived fission product, ¹³⁷Cs, has been reduced to very low values (Pantelić et al., 2007).

Mosses have some advantages compared to other bioindicator species. The accumulating capacity of mosses is higher than that of other plants

Table 1. Species of mosses collected in the area of six regions around Belgrade classified according to their individual growth substrate

Regions	Moosees species	
	Epilitic (stone)	Epixilic (tree bark)
Opovo	<i>Grimmia pulvinata</i> (Hedw.) Sm.	<i>Hypnum cupressiforme</i> Hedw.
Vinča-Boleč	<i>Tortula muralis</i> Hedw.	<i>Pseudoleskeella nervosa</i> (Bird.) Nyholm
Kosmaj-Nemenikuće	<i>Pylaisia polyantha</i> (Hedw.) Schimp.	<i>Dicranella heteromalla</i> (Hedw.) Schimp
Avala-Zuce	<i>Ditrichum pusillum</i> (Hedw.) Hampe	<i>Hypnum cupressiforme</i> Hedw.
Barajevo	<i>Didymodon spadiceus</i> (Mitt.) Limpr.	<i>Anomodon attenuatus</i> (Hedw.) Huebener
Jakovo-Bojčinska šuma	<i>Brachythecium rutabulum</i> (Hedw.) Schimp	<i>Brachytheciastrum velutinum</i> (Hedw.) Ignatov & Hutten

(Elstner et al., 1987; Mishev et al., 1996). They capture airborne particulates by both passive and active processes through an extra cellular ion-exchange process (Knight et al., 1961).

The radiological investigation of a certain area involves the collecting of samples which represent potential danger for the health of the population. The goal of this paper was to determine the level of activity and presence of natural radionuclides (^{40}K , ^{238}U , ^{226}Ra , ^{232}Th) and the anthropogenic radionuclide ^{137}Cs in the samples of soil (arable and uncultivated), feedstuffs (maize and alfalfa) and mosses on the territory of six regions around Belgrade from which most food for human consumption originates.

The goal of this paper was to determine the level of activity of radio-nuclides ^{40}K , ^{238}U , ^{226}Ra , ^{232}Th and ^{137}Cs in the soil, feedstuffs and mosses (bioindicator species) in the Belgrade environment, Serbia.

MATERIALS AND METHODS

The samples of soil, feedstuffs and mosses were collected during the period of 2007 – 2008 in six regions around Belgrade: Avala-Zuce (a village near the mountain Avala), Kosmaj-Nemenikuće (a village

near the mountain Kosmaj), Barajevo, Grocka - Vinča (a village near the river Danube), Opovo (a village near the town of Pančevo and the river Tamiš) and Jakovo-Bojčin forest (a village near the Bojčin forest and the Sava river). Each sample was taken in triplicate from each sampling site.

These sampling sites were selected because they represent significant arable land for the production of food (and feed) for the Belgrade population.

On each of the six sampling sites two types of soil samples were taken: cultivated and uncultivated. From the arable land (cultivated soil) samples of maize and alfalfa were collected, from uncultivated soil samples of mosses were taken.

Soil samples in amounts of 3-5 kg were collected from a depth of 10-20 cm, homogenized, dried at a temperature of 105°C and put in 1l Marineli beakers. Alfalfa and maize samples were collected in amounts of 2-5 kg, ground and burned to mineral ash. Plastic 200 ml vessels were used for measurements. Moss samples collected in an amount of 2 kg (in fresh form) from uncultivated soil were sorted by their individual growth substrate, into two groups: epilithic (stone) and epixilic (tree bark). Radioactivity was measured in the mixture of moss

Table 2. Average activity concentration of radionuclides in the soil and plants (Bq kg⁻¹) from the environment of Belgrade

Samples	⁴⁰ K	²³⁸ U	²²⁶ Ra	²³² Th	¹³⁷ Cs
Soil cultivated	608 ± 15	47 ± 7	51 ± 5	50 ± 4	25 ± 2
Soil uncultivated	560 ± 16	41 ± 5	48 ± 2	47 ± 2	39 ± 1
Alfaalfa	538 ± 19	< 5	< 9.6	< 1.6	< 0.4
Maize	95 ± 5	< 1.9	< 1.5	< 0.3	< 0.1

mean ± standard deviation

species. The moss samples were first cleaned of soil and other impurities and dried at a temperature of 80°C until they became a constant mass, were ground and packed in 200 ml plastic vessels.

All samples were kept in the laboratory (at room temperature) for 40 days prior to measurement in order to achieve a radioactive balance. The radioactivity of the samples was determined by gamma spectrometric measurements on an HPGe detector (Ortec, USA) with a relative efficiency of 30% and energy resolution 1.85 keV (1332.5 ⁶⁰Co). Radioactive measurements were performed according to the procedure described by Mitrović et al., (2009).

All obtained results were expressed as means ± standard deviation (MEAN ± SD) and are shown in tables (Snedecor and Cochrain, 1971).

RESULTS AND DISCUSSION

Species of mosses collected in each region are shown in Table 1.

Average values of activity of natural and artificial radio-nuclides in samples of cultivated and uncultivated soils, alfalfa and maize, gathered from six sites around Belgrade are shown in Table 2.

In the environment radionuclides of natural (primordial) and artificial (anthropogenic) origin are present. Primordial radionuclides are formed as a result of cosmic and terrestrial radiance and have relatively low concentration. The most significant representatives of natural radionuclides are ⁴⁰K, ²³⁸U, ²²⁶Ra and ²³²Th. The level of uranium in arable land

can be higher as a result of the increased use of phosphorous fertilizers, whose production is directly dependent on the use of phosphate minerals which can often carry significant levels of radioactive uranium. This implies the importance of control of the raw materials used in phosphorous fertilizer production. The presence of anthropogenic radionuclides in the environment is generally a result of nuclear probes and accidents in nuclear plants.

Pantelić et al., (2007) state that after the Chernobyl accident (1986) a high average activity concentration of ¹³⁷Cs was measured in alfalfa (791 Bq/kg) and in mosses (600-2000 Bq/kg) but not in maize (2.1 Bq/kg). Ten years later the activity concentration of ¹³⁷Cs has decreased to an average level of 0.5 Bq/kg in alfalfa and 0.1 Bq/kg in maize. Activity in mosses has decreased but was still relatively high, ranging between 90 and 210 Bq/kg. These results are in accordance with results obtained in our trial.

Even 20 years after the Chernobyl accident, the presence of ¹³⁷Cs can be determined in some samples due to its long half-life (27.7 years). Therefore, it is of great importance to investigate bioindicators because they present the most reliable indicator of whether the living environment was contaminated with ¹³⁷Cs as a result of the Chernobyl accident.

The ability of mosses to show the quality of air, water and soil, based on their presence, absence or floristic composition, presents an important trait and it enables their use as bioindicators of radioactive contamination. Mosses are, due to their anatomical, morphological and physiological characteristics,

Table 3. Average activity concentration of radionuclides in samples of epilithic mosses (Bq kg⁻¹) originating from the Belgrade environment

Regions	⁴⁰ K	²³⁸ U	²²⁶ Ra	²³² Th	¹³⁷ Cs
Opovo	407 ± 17	< 20	18 ± 3	22 ± 4	41 ± 2
Vinča - Boleč	384 ± 12	53 ± 9	70 ± 8	18 ± 1.8	89 ± 3
Kosmaj - Nemenikuće	370 ± 17	56 ± 10	75 ± 10	18.4 ± 3.8	99 ± 3
Avala - Zuce	178 ± 9	< 12.6	6.1 ± 1.3	8.4 ± 2.1	158 ± 5
Barajevo	154 ± 7	< 16	< 26	< 4.7	9 ± 0.4
Jakovo – Bojčin forest	270 ± 16	39 ± 9	11 ± 2	15 ± 3	14 ± 1

mean ± standard deviation

capable of accumulating pollutants from aero sediments, radionuclides and pollutants from the soil in their tissue. They represent good bio-indicators because they: have wide ecological and geological distribution, are present during whole year (and are in most cases perennial plants). Also, they exhibit a high degree of tolerance to pollutants, absorption of mineral matter is performed by the whole body surface and on leaves they do not have either cuticle or stomas. Thus, a clear correlation between the speed of accumulation of radioactive contaminants and moss growth can be observed (Grdović, 2005). Lately, and especially after the Chernobyl accident, mosses are used as reliable indicators of increased activity concentration of radionuclides in the environment (Stanković et al., 1997).

⁴⁰K causes the most dominant natural radioactivity. The activity level of this radionuclide was high in both soil and alfalfa (538-608 Bq/kg), while in maize it was significantly lower (95 Bq/kg). The activity levels of the other natural radionuclides (²³⁸U, ²²⁶Ra and ²³²Th) were lower and within the limits of average values for Serbia (Pantelić et al., 2004).

The activity level of ¹³⁷Cs in the soil was in a range of 25-39 Bq/kg. These results indicate that even after 20 years since the Chernobyl nuclear accident, this radionuclide is present in the natural environment of Belgrade. In the cultivated soil lower levels of ¹³⁷Cs were found than in the non-cultivated soil, but the differences were not significant and was within the limits of the average values for Serbia (Pantelić et al., 2004). These results are in accordance with those of Bikit et al., (2005) and Bolca et al., (2007).

Measured specific activities of natural radionuclides in the plant samples, except for ⁴⁰K, were below detection limits for the applied measurement conditions (geometry of measurement, quantity of the sample and time of measurement). These plants are commonly used as feedstuffs (alfalfa, maize) and ⁴⁰K is determined as a major contribution to the total activity of this kind of sample. Also, all measured and reported specific activities for artificial radionuclides (just ¹³⁷Cs is reported) are below the minimum detectable activity (MDA) values for performed measurements. These results are in accordance with results presented earlier for some regions of Serbia (Pantelić et al., 2004 and 2007).

Table 4. Average activity concentration of radionuclides in samples of epixilic mosses (Bq kg⁻¹) originating from Belgrade environment

Regions	⁴⁰ K	²³⁸ U	²²⁶ Ra	²³² Th	¹³⁷ Cs
Opovo	451 ± 17	38 ± 0.9	41 ± 3	37 ± 4	68 ± 5
Vinča - Boleč	150 ± 9	35 ± 4	7.4 ± 1.1	7.2 ± 1.3	43 ± 5
Kosmaj - Nemenikuće	110 ± 7	32 ± 9	7.1 ± 1.4	7.7 ± 1.9	32 ± 9
Avala - Zuce	490 ± 19	80 ± 10	38 ± 7	45 ± 4	221 ± 13
Barajevo	322 ± 24	< 113	< 189	< 23	63 ± 3
Jakovo – Bojčin forest	247 ± 10	20 ± 3	7.7 ± 1.1	16 ± 2	16 ± 1

mean ± standard deviation

Tables 3 and 4 show the average values of activity levels of natural and artificial radio-nuclides in moss samples gathered from the investigated regions around Belgrade.

There were no significant differences in the activities of natural radio-nuclides between the moss species regarding growth substrate. For all measured samples, the arithmetic mean of ²³⁸U activity was 43 Bq kg⁻¹, for ²²⁶Ra it was 36 Bq kg⁻¹, for ⁴⁰K it was 294 Bq kg⁻¹, and for ²³²Th it was 19 Bq kg⁻¹. Similar results were also found by Dragović et al., (2007),

¹³⁷Cs presence in the environment is due to atmospheric fallout from the Chernobyl accident which occurred in 1986. In the epilithic moss samples collected in the environment around Belgrade, ¹³⁷Cs activity ranged from 9 Bq kg⁻¹ (Barajevo) and 14 Bq kg⁻¹ (Jakovo) to 158 Bq kg⁻¹ d.m. (Avala-Zuce). In the epixilic moss samples, ¹³⁷Cs activity ranged from 16 Bq kg⁻¹ (Jakovo) and 32 Bq kg⁻¹ (Kosmaj) to 221 Bq kg⁻¹ d.m. (Avala-Zuce). Similar results were also found by Delfanti et al., (1999) but Dragovic et al., (2004) reported higher ¹³⁷Cs activities in the mosses collected in the environment of mountain regions of Serbia and Montenegro. Even 20 years after the accident the pre-

sence of ¹³⁷Cs can be detected in the environment of Belgrade. For that purpose mosses are a useful bio-indicator.

Based on the findings shown in Tables 2, 3 and 4 it can be seen that the activity concentration of measured radionuclides, except for ⁴⁰K, was significantly higher in mosses compared to maize and alfalfa. This was the case for both natural radionuclides (²³⁸U, ²²⁶Ra, ²³²Th) and for ¹³⁷Cs, originating from the Chernobyl accident. It can be concluded that maize and alfalfa are plants that do not possess the ability to accumulate radionuclides and are safe as animal feed. The mosses present a more reliable indicator for radioactive contamination of the environment.

Acknowledgment - This work was supported by grant no. 19022 of the Ministry of Science and Technology of Serbia.

REFERENCES

- Adel, A., Uosif, M., and A. El-taher, (2005). Natural radioactivity and dose assessment for phosphate rock from Wadi El-Mashash and El-Mahamid Mines, Egypt. *Journal of Environmental radioactivity*, **84** 1, 65-78.

- Bikit, I., Slivka, J., Čonkić, Lj., Krmar, M., Vesković, M., Žikić-Todorović, N., Varga, E., Ćurčić, S., and D. Mrdja, (2005). Radioactivity of the soil in Vojvodina (northern province of Serbia and Montenegro). *Journal of Environmental Radioactivity*, **78**, 11-19.
- Bolca, M., Sac, M., Cokuysal, B., Karali, T., and E. Ekdal, (2007). Radioactivity in soils and various foodstuffs from the Gediz river basin of Turkey. *Radiation Measurements*, **42**, 263-270.
- Delfanti, R., Papucci, C., and C. Benco, (1999). Mosses as indicators of radioactivity deposition around a coal-fired power station. *The Science of the Total Environment*, **227**, 49-56.
- Dragović, S., Nedić, O., Stanković S., and G. Bačić, (2004). Radiocesium accumulation in mosses from highlands of Serbia and Montenegro: chemical and physiological aspects. *Journal of Environmental radioactivity*, **77**, 381-388.
- Dragović, S., Onjia, A., Dragović, R., and G. Bačić, (2007). Implementaion of neural networks for classification of moss and lichen samples on the basis of gamma-ray spectrometric analysis. *Environmental Monitoring Assess*, **130**, 245-253.
- Elstner, E.F., Fink, R., Holl, W., Lengfelder, E., and H. Ziegler, (1987). Natural and Chernobyl-caused radioactivity in mushrooms, mosses and soil-samples of defined biotops in SW Bavaria. *Oecologia*, **73**, 553-558.
- Grdović, S. (2005). *Mahovine šireg područja Beograda i njihov bioindikatorski značaj*. Monografija, Zadužbina Andrejević, Beograd 2005.
- Hecht, H. (1988). Radioaktive Belastung von wild und nutztieren nach dem Unfall von Tschernobyl. *Fleischwirtsch.*, **68**, 4, 508-513.
- Knight, A.H., Crooke, W.M., and R.M. Inkson, (1961). Cation retain capacities of tissues of higher and lower plants and their related uronic-acid contents. *Nature*, **192**, 142-143.
- Mitrović B., Vitorović G., Vitorović D., Pantelić G., and I. Adamović (2009). Natural and athropogenic radioactivity in the environment of mountain region of Serbia. *J. Environ. Monitor.*, **11**, 383-388.
- Mishev, P., Damynova, A., and L. Yurukova, (1996). Mosses as biomonitors of airborne pollution in the northern part of Rila mountain. *Observatorie de montagne de Moussala OM2*, 137-141.
- Nifontova, M. (1995). Radionuclides in the moss/lichen cover of tundra communities in the Yamal Peninsula. *The Science of Total Environment*, **160/161**, 749-752.
- Pantelić, G., Tanasković, I., Javorina, LJ., Vitorović, G., and V. Vuletić, (2004). Aktivnost ¹³⁷Cs u uzorcima iz životne sredine regiona Užica. *Veterinarski glasnik*, **58**, 3-4, 233-239.
- Pantelić G., Javorina LJ., Vitorović G., Vuletić V., Tanasković I., and M. Eremić-Savković (2007). Two decades of ¹³⁷Cs activity measurements in cattle feed and milk in Serbia. IRPA, Regional Congress for Central and Eastern Europe – Regional and Global Aspects of radiation protection, 24-28 September, Brasov, Romania. Available from: Final programme and Full Papers CD.
- Pietrzak-Flis, Z., Rosiak, L., Suplinska, M.M, Chrzanowski, E., and S. Dembinska, (2001). Daily intakes of ²³⁸U, ²³⁴U, ²³²Th, ²³⁰Th, ²²⁸Th and ²²⁶Ra in the adult population of central Poland. *The Science of The Total Environment*, **273**,163-169.
- Snedecor, G.W. and W.G. Cochran, (1971). *Statistical Methods*, **6th ed**, pp. 593, Iowa University Press, Ames, SAD.
- Stanković, A., Stanković, S., Pantelić, G., and V. Randelović, (1997); Radiokontaminacija mahovina i lišaja sa aspekta biodiverziteta.- Zbornik radova: Rizik tehnoloških sistema i životna sredina, 149-152.

ПРИРОДНА И ВЕШТАЧКА РАДИОАКТИВНОСТ СТОЧНЕ ХРАНЕ, МАХОВИНА И ЗЕМЉИШТА У ОКОЛИНИ БЕОГРАДА

СВЕТЛАНА ГРДОВИЋ*, ГОРДАНА ВИТОРОВИЋ*, БРАНИСЛАВА МИТРОВИЋ*, В. АНДРИЋ**,
Б. ПЕТРУЈКИЋ* и М. ОБРАДОВИЋ*

* Факултет ветеринарске медицине, 11000 Београд, Србија

** Институт за нуклеарне науке «Винча», 11000 Београд, Србија

Гамаспектрометријским мерењима извршено је одређивање садржаја природних радионуклида (^{40}K , ^{238}U , ^{226}Ra , ^{232}Th) и ^{137}Cs у узорцима земљишта, луцерке, кукуруза и маховина, на шест локалитета у околини Београда. Природни радионуклиди у земљишту су били на нивоу граница специфичним за Србију, док је установљен релативно висок ниво активности ^{137}Cs (око 30 Bq kg^{-1}). Насупрот томе, у узорцима биљака, које се најчешће користе као сточна храна (луцерка, кукуруз), концентрација активности природних радионуклида и ^{137}Cs је била ниска

односно испод границе детекције. Садржај природних радионуклида у маховинама био је у стандардним границама детекције специфичним за Србију. Међутим, ниво активности ^{137}Cs у маховинама сакупљеним са широког подручја околине Београда, био је висок, а највиши са подручја Авала-Зуце ($158\text{-}221 \text{ Bq kg}^{-1}$). Добијени резултати указују да је овај радионуклид и после 20 година од несреће у Чернобилу, присутан у животној средини Београда и да су маховине добар индикатор контаминације животне средине са ^{137}Cs .