

Research article

THE INVESTIGATION OF DNA DAMAGE INDUCED BY ADRENALINE IN HUMAN LYMPHOCYTES *IN VITRO*

RADAKOVIĆ Milena^{1*}, ĐELIĆ Ninoslav¹, STEVANOVIĆ Jevrosima¹, ANĐELKOVIĆ Marko², KOLAREVIĆ Stoimir³, DAČIĆ Stefan⁴, STANIMIROVIĆ Zoran¹

¹Department of Biology, Faculty of Veterinary Medicine, University of Belgrade, Bul. Oslobođenja 18, Belgrade, Serbia; ²Serbian Academy of Sciences and Arts, Knez Mihailova 35, Belgrade, Serbia; ³Department of Microbiology, Faculty of Biology, University of Belgrade, Studentski trg 16, Belgrade, Serbia; ⁴School of Medicine, University of Niš, Bul. dr Zorana Djindžica 81, Niš, Serbia

(Received 09 May; Accepted 24 August 2014)

Adrenaline is a neurotransmitter and hormone that plays an important role in physiological regulatory mechanisms. The objective of this study was to assess primary DNA damage in isolated human lymphocytes exposed to adrenaline using the *in vitro* comet assay. Dose-response of human lymphocytes was determined at concentration range of adrenaline from 0.01 μM to 300 μM for various treatment times (1h, 2h, 4h and 24h). The obtained results showed that adrenaline induced DNA damage at concentration range from 5 μM to 300 μM after 1h, 2h and 4h of treatment. The slightest DNA damage was observed after 24 h of adrenaline treatment - only the highest concentrations of adrenaline (150 μM and 300 μM) caused increased level of DNA damage. In order to evaluate the potential contribution of reactive oxygen species (ROS) in adrenaline-induced DNA damage we used antioxidants catalase (100 IU/mL and 500 IU/mL) and quercetin (100 μM and 500 μM). Co-treatment of lymphocytes with adrenaline (300 μM) and antioxidants for 1 h, significantly reduced the quantity of DNA in the comet tails. Therefore, it can be concluded that adrenaline exhibits genotoxic effects mainly through induction of reactive oxygen species and that some of the DNA damage is repaired during the first four hours following the treatment with adrenaline.

Key words: adrenaline, antioxidants, comet assay, human lymphocytes.

INTRODUCTION

Catecholamines are a class of chemical neurotransmitters and hormones that play an important role mainly in physiological hemodynamic regulatory mechanisms. The primary effect of the catecholamine adrenaline is physiological mobilization of resources in response to emotional and physical stress. In some animals, adrenaline is involved in temperature regulation and arousal from hibernation [1]. Adrenaline is

* Corresponding author: e-mail: mradakovic@vet.bg.ac.rs

also involved in the immune response via stimulating inflammatory cytokines [2]. In human medicine, adrenaline has its application in the treatment of asthma, allergic reactions, cardiac arrest and glaucoma [3]. During normal physiological conditions, there is no constant secretion of adrenaline and the basal level of adrenaline in human plasma is in a nanomolar range [4]. Under stress, excess adrenaline is released in the circulation and, after binding to adrenergic receptors, it prepares the organism for the „fight or flight“ response [5,6].

Although catecholamines play a vital role under stressful conditions, excess circulating levels of adrenaline may induce detrimental effects in cells. Cardiotoxicity of adrenaline is well documented [7,8]. It was found that adrenaline stimulates proliferation of esophageal cell carcinoma via β -adrenoreceptor activation [9]. In breast cancer and adenocarcinoma adrenaline induces chemoresistance via α_2 -adrenergic receptors [10,11]. Besides, the mutagenic potentials of adrenaline and dopamine have been reported in tests for gene mutations in L5178Y mouse lymphoma cells [12]. More recent studies revealed that adrenaline, noradrenaline and cortisol induce DNA damage in the comet assay on 3T3 cells [13]. However, adrenaline did not induce chromosome aberrations in cultured human lymphocytes [14].

It is assumed that catecholamines exert genotoxic effects via oxidative products formed during oxidative metabolism. In a study of Djelic and Anderson [15] catalase exhibited a protective effect against DNA damaging effect of noradrenaline, *indicating the involment of ROS in genotoxicity of catecholamine*. Similarly, Miura [16] found that adrenaline and noradrenaline induce DNA strand breaks in plasmid PM2 DNA due to the creation of oxygenated derivatives of catecholamines. It is noted that oxidative DNA damage is an important mutagenic and possibly carcinogenic factor [17].

These observations prompted us to investigate the influence of adrenaline on primary DNA damage on human lymphocytes. For this purpose the effect of adrenaline was evaluated by *in vitro* comet assay. To investigate whether the mechanism underlying DNA damage of adrenaline is mediated by ROS we used antioxidants catalase and quercetin in the comet assay.

MATERIALS AND METHODS

Blood sampling and cell preparation

Peripheral blood samples (4 mL) with heparin were collected by venipuncture from two healthy male donors under 25 years of age. Whole blood was diluted 1:1 with RPMI medium, underlaid with Ficoll-Paque (Sigma) and centrifuged at 1900 g for 15 min. The lymphocyte layer (buffy coat) was washed twice in RPMI 1640 medium, each wash was followed by centrifugation for 10 min at 1800 g. The cells were suspended in a total volume of 1 ml and each reaction contained 50 μ l suspension ($\approx 10^4$ cells), various amounts (μ l) of the test agent and PBS buffer. The number of viable cells was determined by Trypan blue exclusion [18].

Chemical treatments

Isolated lymphocytes were treated with various concentrations of adrenaline (from concentrations corresponding to physiological values in humans (0.0005 μM) to a 60x higher concentration than the maximum therapeutic dose (300 μM) at different time intervals. The highest concentration of adrenaline (300 μM) was used for further analysis with antioxidants catalase and quercetin. Simultaneous treatment of human lymphocytes with adrenaline and the antioxidant catalase (100 IU/mL and 500 IU/mL) was conducted for 1 h and the same procedure was performed with quercetin (100 μM and 500 μM). Hydrogen peroxide (100 μM) was used as the positive control, while PBS was the negative control.

The Comet assay

Before the start of the experiment the cells were checked for viability using Trypan blue dye according to the method of Phillips [18]. Alkaline comet assay was performed according to Singh [19] and Tice *et al.* [20,21] technique with slight modifications. Briefly, after incubation with the tested compound for 1, 2, 4 and 24 h at 37°C, 100 μL of cell suspension was mixed with 100 μL of 1% low melting point agarose (LMPA). The 90 μL of suspension was rapidly pipetted onto the thin agarose layer of 1% normal melting point agarose (LMPA) spread with a coverslip, and kept at 4°C for 5 min. to solidify. After removal of the coverslip, the 90 μL of 0.5% LMPA was added, spread using a coverslip and kept at 4°C for 5 min. When the agarose was solidified, the slides were immersed in cold lysis solution (2.5 M NaCl, 100 mM EDTA, 10 mM Tris, 1% Triton X-100, 10% DMSO, pH 10) overnight at 4°C. After lysis, the slides were placed in a horizontal gel electrophoresis tank and kept in freshly made cold alkaline electrophoresis buffer (300 mM NaOH, 1 mM EDTA, pH > 13) for 30 min to allow unwinding of DNA. Electrophoresis was conducted at 4°C with 25 V and 300 mA for 30 min. After electrophoresis the slides were neutralized with Tris-HCl buffer (0.4 M, pH 7.5) for 5 min. The neutralization procedure was repeated three times. Then, the slides were fixed with cold methanol, dried and stored. Before analysis, the slides were rehydrated with ice cold distilled water and stained with 50 μL of 20 $\mu\text{g}/\text{mL}$ ethidium bromide.

For visualization of DNA damage, microscope slides were examined under 400 \times magnification on a fluorescence microscope Olympus, CX21 (Olympus Optical Co., GmbH Hamburg, Germany). Images of 100 randomly selected lymphocytes (50 cells from each of two replicate slides were analyzed) for each sample and the DNA damage was scored visually as described by Anderson *et al.* [22]. Namely, cells were graded by eye into five categories corresponding to the following amounts of DNA in the tail: (A) no damage, <5%; (B) low level damage, 5-20%; (C) medium level damage, 20-40%; (D) high level damage, 40-95%; (E) total damage, >95%. The number of comets in each sample was calculated (0 x No Migration (NM) + 1 x Low Migration (LM) + 2 x Medium Migration (MM) + 3 x High Migration (HM) + 4 x Extensive

Migration (EM)) which was referred by Collins [23] and expressed as the total comet score (TCS).

Statistical analysis

Statistical analysis was performed using analysis of variance (ANOVA) followed by Tukey's multiple test (GraphPad Software, USA). The value of total comet scores (TCS) is given as mean \pm SEM. A P value of at least ≤ 0.05 is considered as statistically significant.

RESULTS

The viability of cells treated with adrenaline was greater than 90% in trypan blue exclusion test at the time of the assay. DNA damage in human lymphocytes was expressed as the TCS score. Table 1 shows the effect of adrenaline on DNA damage in human lymphocytes after different incubation times. There was a tendency of dose-dependent elevation of DNA damage in lymphocytes exposed to adrenaline. A significantly increased level of DNA damage was observed at a concentration of 5 μ M - 300 μ M of adrenaline compared to untreated cells after 1 hour ($P < 0.01$, $P < 0.001$). As expected cells exposed to hydrogen peroxide (100 μ M) showed a high degree of DNA damage at all incubation times ($P < 0.001$). After 2 and 4 hours the value of TCS was significantly increased at doses 5-300 μ M ($P < 0.05$, $P < 0.001$). The lowest DNA damage in human lymphocytes was observed at one day treatment of adrenaline, when only the highest concentrations of adrenaline (150 and 300 μ M) induced significant increases level of TCS i.e. DNA damage in human lymphocytes ($P < 0.001$).

In Figure 1. The DNA damaging effect of adrenaline was presented as distribution of cells (%). It was observed that at concentration of 5-300 μ M adrenaline, 18-20% of total cells were in the category D (high damage) after 1 hour. However, a weaker effect was observed after 2 hours, since 16-10% of lymphocytes were in this category (D). A similar effect was observed after 4 hour treatment of adrenaline (Table 1, Figure 1). When cells were exposed to adrenaline for 24 hours, the highest percentage of undamaged cells was in the minimum damage category (B).

The effects of catalase and quercetin on DNA damage induced by adrenaline after 60 minutes are shown in Figure 2. After one hour, catalase (100 IU/mL) in the treatment with adrenaline (300 μ M) significantly reduced DNA damage in human lymphocytes ($P < 0.05$). Also, a significant decreasing trend of DNA damage was observed at 500 IU/mL of catalase ($P < 0.01$). Unlike catalase, quercetin at concentration of 100 μ M did not significantly reduce the level of DNA damage caused by adrenaline ($P > 0.05$), while a protective effect of quercetin was noted at a higher concentration (500 μ M) ($P < 0.01$).

Table 1. The effects of adrenaline in human lymphocytes measured by the Comet assay at different exposure times

Treatment	TCS at various incubation times			
	1h	2h	4h	24h
Negative control (PBS)	61.17±0.48	60.50±0.56	60.83±3.97	57.17±3.54
Adrenaline				
0.01 µM	62.33±0.49	61.50±0.72	61.17±1.28	56.33±0.88
1 µM	63.67±0.99	63.33±1.23	69.67±2.94	57.50±0.72
5 µM	69.33±1.02**	65.33±2.62*	75.33±3.05***	60.33±1.05
50 µM	76.33±1.41***	75.83±2.30***	92.50±2.68 ***	61.00±1.37
150 µM	95.17±1.41***	94.83±1.76***	104.50±2.78 ***	64.83±1.30***
300 µM	107.80±0.99***	108.20±1.24 ***	116.70±1.62 ***	66.67±1.12***
Positive control (H ₂ O ₂)	184.50±2.78***	195.50±7.28***	189.30±4.59***	189.30±4.34***
100 µM				

Results of the total comet score (TCS) are presented as mean values ± SE from 100 cells per experiment. Three independent experiments were performed. Statistically significant increase compared to solvent control (PBS): *P < 0.05; **P < 0.01; ***P < 0.001




Figure 1. Distribution class of comets in human lymphocytes A. treatment with adrenaline for 1 hour, B. treatment with adrenaline for 2 hours, C. treatment with adrenaline for 4 hours, D. treatment with adrenaline for 24 hours. Statistically significant increase compared to solvent control (PBS): *P < 0.05; **P < 0.01; ***P < 0.001.




Figure 2. The effect of catalase and quercetin against DNA damage of adrenaline in human lymphocytes after 60 minutes. The results are presented as mean TCS values \pm SE from 100 cells per experiment; C- negative control (PBS); A- adrenaline; A+CAT - simultaneous treatments of adrenaline and catalase (100 and 500 IU/mL); A+Q - simultaneous treatments of adrenaline and quercetin (100 and 500µM). Statistically significant increase compared to adrenaline: *P < 0.05; **P < 0.01

DISCUSSION

Some natural hormones can act as endogenous mutagens [24,25] and can also influence cell proliferation [26]. In addition to oestrogens, as the most studied endogenous mutagens, thyroid hormones and catecholamines may induce oxidative stress favoring the processes of mutagenesis.

The catecholamines have always attracted the attention of the scientific community due to their importance in physiological and pathological processes. Since adrenaline is an endogenous substance, it has not been considered as a possible mutagen. Hence, the data concerning evaluation of mutagenic effects of adrenaline are scarce. In this work we investigated the level of primary DNA damage in isolated human lymphocytes exposed to adrenaline using the *in vitro* comet assay.

We observed that adrenaline induced a significant increase of the TCS values at concentrations of 5 µM (maximum therapeutic dose) and higher applied concentrations compared to the negative control after 1 h, 2 h and 4 h of treatments. However, after 24 h treatment, only the highest concentrations of adrenaline (150 and 300 µM) induced significant DNA damage. This signifies that increased DNA migration detected at shorter incubation times in the comet assay could be due to strand breaks still unrepaired by excision repair [27]. We suppose that cells have repaired the most of DNA damage after 24 hours which resulted in reduced genotoxic effects of adrenaline.

Although a percentage of comets might have resulted as a consequence of early apoptosis, the comet assay is regarded as a specific test for genotoxicity. Namely, the results in the comet assay are not necessarily caused by concomitant processes leading to apoptosis [28]. Actually, DNA breakage detected in the comet assay is mainly caused by mutagen-induced DNA damage [29].

These results suggest that adrenaline induces DNA damage in human lymphocytes which is consistent with the findings of Flint *et al.* [13] that stress hormones, including adrenaline, induced DNA damage in murine 3T3 cell in the comet assay. Miura *et al.* [16] also indicated on the genotoxic potential of adrenaline. However, Djelic *et al.* [14] reported the absence of clastogenic effects of adrenaline in human lymphocytes using an *in vitro* cytogenetic test. The lack of clastogenic activity indicates that adrenaline induced DNA damage detected in the comet assay only at the molecular level. This assumption is in agreement with findings of Lankoff *et al.* [30] that microcystin-LR induced DNA breaks in the comet assay, but was unable to induce chromosome aberrations.

The catecholamines can be involved in the redox cycling accompanied by production of ROS [31,32,33]. Therefore, we used antioxidants catalase and quercetin to examine whether DNA damaging effects of adrenaline is mediated by ROS. Our results show that catalase (100IU/mL and 500 IU/mL) and quercetin (500 μ M) in co-treatment with adrenaline significantly reduced DNA damage in lymphocytes. However, quercetin at 100 μ M did not significantly reduce DNA damage. We assume that lower doses of quercetin in human lymphocytes are not high enough to reduce genotoxic effects, and possibly quercetin is also metabolized to the less effective form [34]. Protective effect of antioxidants used in our investigation indicated the involvement of free radicals in the genotoxic effect of adrenaline.

The contribution of ROS in genotoxicity of catecholamines is supported by investigation of Djelic and Anderson [15] showing that catalase reduced DNA damaging effect of noradrenaline in isolated human lymphocytes. Also, Miura *et al.* [16] showed that catalase exhibited a protective effect against genotoxicity of adrenaline and noradrenaline on plasmid DNA.

DNA damaging effect of adrenaline detected in the comet assay is likely to be due to oxidative products of adrenaline. In this regard, oxidation products of adrenaline have been described in the heart, skeletal muscle, liver and blood [35,36]. Because of the unstable nature of the catechol group, catecholamine can be easily oxidized to reactive quinone and semiquinone producing ROS. Superoxide anion (O_2^-) generated by oxidative metabolism of adrenaline can be converted to hydroxyl radical (OH^\bullet) and induce DNA breaks [37]. Also O_2^- may promote further oxidation of adrenaline and increase ROS production [31]. It was found that O_2^- can reduce the activity of antioxidant enzymes such as catalase, glutathione peroxidase and NADH dehydrogenase [38]. On the other hand, quinone formed during the adrenaline oxidation process [39] can form glutathionyl adducts [40]. GSH conjugates were found to covalently bind with DNA and generate apurinic sites, which can lead to mutations [41].

It is noteworthy that we detected DNA damage of adrenaline at a concentration which is several times higher than the normal physiological levels in human blood. However, markedly elevated concentrations of circulating catecholamines are found in ischemia [42, 43], pheochromocytoma [44] and drug abuse [45]. The obtained results indicate

that applied concentrations of adrenaline (5-300 μM) established a redox cycle that multiplies ROS production. Under this condition antioxidative enzymes do not have the ability to protect cells against excess level of ROS attack which leads to oxidative stress. There is ample evidence which suggest the strong association of oxidative stress, genetic instability and cancer development [46]. ROS have the ability to cause mutations in the proto-oncogenes and thus create the possibility to be involved in malignancy [47,48].

An elevated concentration of adrenaline occurs in acute stress and can be further increased during chronic stress [49,50]. Experimental findings of DNA damaging effects of adrenaline are significant, because there is evidence that stress hormones induce cell transformation via signal transduction pathway [13]. Namely, stress hormones induce DNA damage, preventing the entry of cells in apoptosis and cell cycle arrest contributing to the transformation of cells. In support of this it is established that stress increases oxidative damage and affects the processes of DNA repair in cells [51,52]. On coherence between adrenaline and oxidative stress pointed Pereira *et al.* [53]. Their findings pointed that adrenaline regulates the activities of catalase and glutathione peroxidase, but the exact mechanism remains to be determined.

Taken together, these results suggest that adrenaline has the ability to affect genomic stability and *further* research should, *therefore* be undertaken to fully *understand* the effects of adrenaline on genetic integrity of cells.

Acknowledgements

This research was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Grant no. III 46002).

REFERENCES

1. Bentley PJ: Endocrine Pharmacology: Physiological Basis and Therapeutic Applications. England: Cambridge University Press; 2011, 204.
2. Röntgen P, Sablotzki A, Simm A, Silber RE, Czeslick E: Effect of catecholamines on intracellular cytokine synthesis in human monocytes. *Eur Cytokine Netw* 2004, 15:14-23.
3. National Toxicology Program (NTP): Toxicology and carcinogenesis studies of l-Epinephrine Hydrochloride (CAS No. 55-31-2) in F344/N rats and B6C3F1 mice (inhalation studies). *Natl Toxicol Program Tech Rep Ser* 1990, 380:1-205.
4. Goldstein DS, Eisenhofer G, Kopin IJ: Sources and significance of plasma levels of catechols and their metabolites in humans. *J Pharmacol Exp Ther* 2003, 305:800–811.
5. Sherwood L: Human Physiology: From Cells to Systems. Belmont CA: Thomson Brooks/Cole, 7 th Revised; 2009, 928.
6. Gavrilovic Lj, Stojiljkovic V, Kasapovic J, Pejic S, Todorovic A, Pajović BS, Dronjak S: Chronic physical stress changes gene expression of catecholamine biosynthetic enzymes in the adrenal medulla of adult rats. *Acta Vet- Beograd* 2012, 62(2-3):151-169.

7. Behonick GS, Novak MJ, Nealley EW, Baskin SI: Toxicology update: the cardiotoxicity of the oxidative stress metabolites of catecholamines (aminochromes). *J Appl Toxicol* 2001, 21 Suppl 1:S15-22.
8. Costa, VM, Silva R, Ferreira LM, Branco PS, Carvalho F, Bastos ML, Carvalho RA, Carvalho M, Remião F: Oxidation process of adrenaline in freshly isolated rat cardiomyocytes: formation of adrenochrome, quinoproteins, and GSH adduct. *Chem Res Toxicol* 2007, 20(8):1183-1191.
9. Liu X, Wu WK, Yu L, Sung JJ, Srivastava G, Zhang ST, Cho CH: Epinephrine stimulates esophageal squamous-cell carcinoma cell proliferation via beta adrenoceptor-dependent transactivation of extracellular signal-regulated kinase/cyclooxygenase-2 pathway. *J Cell Biochem* 2008, 105(1):53-60.
10. Su F, Ouyang N, Zhu P, Ouyang N, Jia W, Gong C, Ma X, Xu H, Song E: Psychological stress induces chemoresistance in breast cancer by upregulating *mdr1*. *Biochem Biophys Res Commun* 2005, 329(3):888-897.
11. Yao H, Duan Z, Wang M, Awonuga AO, Rappolee D, Xie Y: Adrenaline induces chemoresistance in HT-29 colon adenocarcinoma cells. *Cancer Genet Cytogenet* 2009, 190(2):81-87.
12. McGregor, DB, Raich CG, Brown A, Edwards I, Reynolds D, West K, Willington S: Reactivity of catecholamines and related substances in mouse lymphoma L5178Y Cell assay for mutagens. *Environ Mol Mutagen* 1988, 11:523-544.
13. Flint MS, Baum A, Chambers WH, Jenkins FJ: Induction of DNA damage, alteration of DNA repair and transcriptional activation by stress hormones. *Psychoneuroendocrinology* 2007, 32(5):470-479.
14. Djelic N, Djelic D, Spremo-Potparevic B, Markovic B, Zivkovic L: Cytogenetic analysis of the effects of epinephrine on cultured human lymphocytes. *Acta Vet- Beograd* 2003, 53:113-120.
15. Djelic N, Anderson D: The effect of the antioxidant catalase on oestrogens, triiodothyronine and noradrenaline in the Comet assay. *Teratog Carcinog Mutagen* 2003, 23:69-81.
16. Miura T, Muraoka S, Fujimoto Y, Zhao K: DNA damage induced by catechol derivatives. *Chem Biol Interact* 2000, 126:125-136.
17. Waris G, Ahsan H: Reactive oxygen species: role in the development of cancer and various chronic conditions. *J Carcinog* 2006, 5:14.
18. Phillips HJ: Dye exclusion tests for cell viability. In Kruse, PF, Patterson MK, editors. *Tissue Culture: Methods and Applications*. New York: Academic Press; 1973, 407-408.
19. Singh NP, Mc Coy, MT, Tice RR, Schneider EL. A simple technique for quantitation of low levels of DNA damage in individual cells. *Exp Cell Res* 1988, 175:184-191.
20. Tice RR, Andrews PW, Hirai O, Singh NP: The single cell gel (SCG) assay: an electrophoretic technique for the detection of DNA damage in individual cells. *Adv Exp Med Biol* 1991, 283:157-164.
21. Tice RR, Agurell E, Anderson D, Burlinson B, Hartmann A, Kobayashi H, Miyamae Y, Rojas E, Ryu JC, Sasaki YF: Single cell gel/comet assay: Guidelines for *in vitro* and *in vivo* genetic toxicology testing. *Environ Mol Mutagen* 2000, 35:206-221.
22. Anderson D, Yu T-W, Phillips BJ, Schmezer P: The effect of various antioxidants and other modifying agents on oxygen-radical-generated DNA damage in human lymphocytes in the COMET assay. *Mutat Res* 1994, 307:261-271.

23. Collins AR: The comet assay. Principles, applications, and limitations. *Methods Mol Biol* 2002, 203:163-177.
24. Djelic N, Spremo-Potparevic B, Markovic B, Zivkovic L, Djelic D: Cell cycle kinetics and cytogenetic changes in human lymphocytes exposed to oestradiol in vitro. *Acta Vet-Beograd* 2006, 56:37-48.
25. Djelic N, Nestic I, Stanimirovic Z, Jovanovic S: Evaluation of the genotoxic effects of thyroxine using in vivo cytogenetic test on Swiss albino mice. *Acta Vet- Beograd* 2007, 57: 487-495.
26. Djelic N: Evaluation of mitogenic effects of oxytocin on cultured human lymphocytes. *Acta Vet- Beograd* 2002, 52: 19-26.
27. Tuck A, Smith S, Larcom L. Chronic lymphocytic leukemia lymphocytes lack the capacity to repair UVC-induced lesions. *Mutat Res* 2000, 459:73–80.
28. Roser S, Pool-Zobel BL, Rechkemmer G: Contribution of apoptosis to responses in the comet assay. *Mutat Res* 2001, 497(1-2):169-175.
29. Rundell MS, Wagner ED, Plewa MJ: The comet assay: genotoxic damage or nuclear fragmentation? *Environ Mol Mutagen* 2003, 42(2):61-67.
30. Lankoff A, Carmichael WW, Grasman KA, Yuan M: The uptake kinetics and immunotoxic effects of microcystin-LR in human and chicken peripheral blood lymphocytes in vitro. *Toxicology* 2004, 204(1):23-40.
31. Genova ML, Abd-Elsalam, NM, Mahdy el SM, Bernacchia A, Lucarini, M, Pedulli GF, Lenaz G: Redox cycling of adrenaline and adrenochrome catalysed by mitochondrial Complex I. *Arch Biochem Biophys* 2006, 447(2):167-173.
32. Miyazaki I, Asanuma M: Dopaminergic neuron-specific oxidative stress caused by dopamine itself. *Acta Med Okayama* 2008, 62:141–150.
33. Manini P, Panzella L, Napolitano A, d'Ischia M: Oxidation chemistry of norepinephrine: partitioning of the O-quinone between competing cyclization and chain breakdown pathways and their roles in melanin formation. *Chem Res Toxicol* 2007, 20:1549–1555.
34. Sousa RL, Marletta MA: Inhibition of cytochrome P450 activity in rat liver microsomes by the naturally occurring flavonoid, quercetin. *Arch Biochem Biophys* 1985, 240: 345–357.
35. Dhalla, SN, Sasaki H, Mochizuki S, Dhalla SK, Liu X, Elimban V: Catecholamine-induced cardiomyopathy. London: Taylor and Francis, 3rd ed; 2001, 269-318.
36. Silva R, Boldt S, Costa VM, Carmo H, Carvalho M, Carvalho F, Bastos Mde L, Lemos-Amado F, Remião F: Evaluation of GSH adducts of adrenaline in biological samples. *Biomed Chromatogr* 2007, 21:670-9.
37. Dizdaroglu M, Jaruga P, Birincioglu M, Rodriguez H: Free-radical-induced damage to DNA: mechanisms and measurement. *Free Rad Bio Med* 2002, 32:1102–15.
38. Willcox JK, Ash SL, Catignani GL: Antioxidants and prevention of chronic disease. *Crit Rev Food Sci Nutr* 2004, 44(4):275-295.
39. Remião F, Rettori D, Han D, Carvalho F, Bastos ML, Cadenas E: Leucoisoprenochrome-o-semiquinone formation in freshly isolated adult rat cardiomyocytes. *Chem Res Toxicol* 2004, 17:1584–1590.
40. Monks TJ, Jones DC, Bai F, Lau SS: The role of metabolism in 3,4-(()-methylenedioxyamphetamine and 3,4-(()-methylenedioxymethamphetamine (Ecstasy) toxicity. *Ther Drug Monit* 2004, 26:132–136.
41. Cavalieri EL, Li KM, Balu N, Saeed M, Devanesan P, Higginbotham S, Zhao J, Gross ML, Rogan EG: Catechol ortho-quinones: the electrophilic compounds that form depurinating

- DNA adducts and could initiate cancer and other diseases. *Carcinogenesis* 2002, 23:1071-1077.
42. Akiyama T, Yamazaki T: Myocardial interstitial norepinephrine and dihydroxyphenylglycol levels during ischemia and reperfusion. *Cardiovasc Res* 2001, 49:78–85.
 43. Milovanovic A, Milovanovic J, Milovanovic A, Konstatinovic Lj, Petrovic M, Kekus D, Petronijevic Vrzic S, Artiko V: Metabolism Of Biogenic Amines In Acute Cerebral Ischemia: Influence Of Systemic Hyperglycemia. *Acta Vet- Beograd* 2012, 62(4):385-401.
 44. Hoffman BB: Catecholamines, sympathomimetic drugs, and adrenergic receptor antagonists. In: Goodman & Gilman's: the Pharmacological Basis of Therapeutics, Hardman, JG, Limbird, LE and Gilman, AG (eds). New York: McGraw-Hill; 2001, 215–268.
 45. Carvalho F, Remião F, Amado F, Domingues P, Correia AJF and Bastos ML: d Amphetamine interaction with glutathione in freshly isolated rat hepatocytes. *Chem Res Toxicol* 1996, 9:1031-1036.
 46. Klaunig JE, Kamendulis LM: The role of oxidative stress in carcinogenesis. *Annu Rev Pharmacol Toxicol* 2004, 44:239-267.
 47. Behrend L, Henderson G, Zwacka, RM. Reactive oxygen species in oncogenic transformation. *Biochem Soc Trans* 2003, 31:1441-1444.
 48. Wu LL, Chiou CC, Chang PY, Wu JT: Urinary 8-OHdG: a marker of oxidative stress to DNA and a risk factor for cancer, atherosclerosis and diabetics. *Clin Chim Acta* 2004, 339: 1-9.
 49. Gold SM, Zakowski SG, Valdimarsdottir HB, Bovbjerg DH: Higher Beck depression scores predict delayed epinephrine recovery after acute psychological stress independent of baseline levels of stress and mood. *Biol Psychol* 2004, 67:261-73.
 50. Wuren M, Yougang Z, Huangrong L, Banchao S, Jiefeng L, Jin Y, Changrong W, Xiaolong X, Yuying H, Xiaoxi L, Fenghua L, Jianqin X: Effect of transport stress on peripheral blood lymphocyte subsets and Th cytokines in pigs. *Acta Vet- Beograd* 2013, 63(2-3):177-190.
 51. Cohen L, Marshall GD Jr, Cheng L, Agarwal SK, Wei Q: DNA repair capacity in healthy medical students during and after exam stress. *J Behav Med* 2000, 23(6):531-544.
 52. Forlenza, MJ, Latimer JJ, Baum A: The effects of stress on DNA repair capacity. *Psychology and Health* 2000, 15:881-889.
 53. Pereira B, Costa-Rosa LF, Bechara EJ, Newsholme P, Curi R: Changes in the TBARS content and superoxide dismutase, catalase and glutathione peroxidase activities in the lymphoid organs and skeletal muscles of adrenomedullated rats. *Braz J Med Biol Res* 1998, 31(6):827-833.

ISPITIVANJA OŠTEĆENJA DNK IZAZVANIH ADRENALINOM U LIMFOCITIMA ČOVEKA *IN VITRO*

RADAKOVIĆ Milena, ĐELIĆ Ninoslav, STEVANOVIĆ Jevrosima, ANĐELKOVIĆ Marko, KOLAREVIĆ Stoimir, DAČIĆ Stefan, STANIMIROVIĆ Zoran

Adrenalin je neurotransmiter i hormon koji ima važne uloge u fiziološkim regulatornim mehanizmima. Predmet istraživanja u ovom radu bila je evaluacija primarnih oštećenja DNK na izolovanim humanim limfocitima izloženim dejstvu adrenalina pri-

menom *in vitro* Komet testa. Odnos doza-odgovor na limfocitima čoveka određen je u rasponu koncentracija adrenalina od 0.01 μM do 300 μM pri različitim dužinama tretmana (1h, 2h, 4h i 24h). Dobijeni rezultati pokazuju da adrenalin indukuje oštećenja DNK u rasponu od 5 do 300 μM nakon 1, 2 i 4 h tretmana. Najmanje oštećenje DNK zapaženo je nakon 24 h – samo su najveće koncentracije adrenalina (150 μM i 300 μM) prouzrokovale povećan stepen oštećenja DNK. Da bi odredili moguć doprinos reaktivnih kiseoničnih vrsta (ROS) u nastanku oštećenja DNK pod dejstvom adrenalina, koristili smo antioksidanse katalazu (100 IU/mL i 500 IU/mL) i kvercetin (100 μM i 500 μM). Kotretman limfocita adrenalinom (300 μM) i antioksidansima u trajanju od 1 h, značajno je smanjio količinu DNK u repovima kometa. Prema tome, može se zaključiti da adrenalin ispoljava genotoksične efekte uglavnom preko stvaranja reaktivnih kiseoničnih vrsta, a jedan deo oštećenja DNK se popravi tokom prva četiri časa, nakon tretmana adrenalinom.