

Originalni naučni rad

**UČESTALOST UTVRĐIVANJA PROMENA NA PARENHIMATOZNYM ORGANIMA
TOKOM ULTRAZVUČNOG PREGLEDA GENITALNOG TRAKTA****

Strahinja ĆIBIĆ^{1*}, Miloš PAVLOVIĆ²

1 Strahinja Ćibić, Fakultet veterinarske medicine Univerziteta u Beogradu, Bulevar oslobođenja 18, Beograd, Srbija

2 Prof.dr Miloš Pavlović, Katedra za porodiljstvo, sterilitet i veštačko osemenjavanje, Fakultet veterinarske medicine Univerziteta u Beogradu, Bulevar oslobođenja 18, Beograd, Srbija

* Korespondentni autor: Strahinja Ćibić: strahinjacicibic@yahoo.com

Kratak sadržaj: Ultrazvučna *Real time 2D* dijagnostika, kao dostupna i neinvazivna metoda, sa uspehom se koristi u medicini, veterini i biologiji, za dijagnostiku kako fizioloških tako i patoloških stanja. Kod malih životinja, ultrazvuk se često koristi u dijagnostici graviditeta, broja, veličine i vitalnosti fetusa, kao i za postavljanje dijagnoze patoloških stanja koja se odvijaju u genitalnom traktu, a kod muških još i za pregled statusa testisa i prostate. Napredna primena ultrazvučne dijagnostike je dovela do toga da se u današnje vreme slobodno mogu povezati brojna patološka stanja reproduktivnog trakta kako ženskih, tako i muških životinja, koja direktno utiču na zdravstveno stanje ostalih unutrašnjih organa. Cilj ovog rada je dijagnostika primarnih poremećaja genitalnog trakta putem ultrazvučne dijagnostike, kao i dokazivanje njihove povezanosti sa promenama na parenhimatoznim organima. Ultrazvučni pregled izvršen je na ukupno 12 pasa, šest mužjaka i šest ženki. Jajnici, materice i abdominalni organi su pregledani sondama u opsegu od 5 do 8 MHz. Za pregled prostate i testisa je korišćena sonda od 7,5 MHz. Pacijenti su bili položeni u ležeći položaj. Od pregledanih šest mužjaka, tri su imala promene na prostati, dok je jedan pas imao tumor testisa. Kod ovih pasa su takođe primećeni cistitis i hiperplazija prostate. Od šest pregledanih ženki, jedna je imala cističnu hiperplaziju endometrijuma, tri su imale piometru, dok je kod jedne dijagnostikovana folikularna cista jajnika. Pored ovih patoloških stanja, utvrđene su promene na jetri i pankreasu. Ultrazvučnom dijagnostikom se pored primarnih promena na genitalnom traktu kako mužjaka tako i ženki, mogu vizuelizovati i posledične promene na parenhimatoznim organima.

Кljučне речи: ultrazvuk, genitalni trakt, parenhimatozni organi, psi

UVOD

Ultrazvučna *Real time 2D* dijagnostika, kao dostupna i neinvazivna metoda, sa uspehom se koristi u medicini, veterini i biologiji, za dijagnostiku kako fizioloških tako i patoloških stanja (Marković i sar, 2003). Primena ove metode je bila pionirska u ranoj dijagnostici

graviditeta kobilu osamdesetih godina prošlog veka, a zatim opšte prihvaćena u humanoj medicini koju su lekari uveli u ekspanziju u eksperimentima i svakodnevnoj kliničkoj praksi. Danas je ovaj dijagnostički postupak gotovo nezamenljiv u svakodnevnoj kliničkoj praksi.

** Rad je prezentovan na 23. Godišnjem savjetovanju doktora veterinarske medicine Republike Srpske (BiH) sa međunarodnim učešćem, Teslić 2018.

Učestalost utvrđivanja promena na parenhimatoznim organima tokom ultrazvučnog pregleda genitalnog trakta

Najbolji opis ove metode je da ona predstavlja stetoskop lekara i veterinara 21. veka.

Početak masovne primene ultrazvučne dijagnostike je doneo i suočavanje sa jako visokim cenama aparata, ali omasovljenje i tehnološki napredak je učinio da ova metoda postane dostupna u širokoj praksi (Lukač i sar, 1994). Što se tiče veterinarske kliničke prakse ultrazvuk je doneo revolucionarni napredak kako kod velikih životinja tako i kod malih. Primena ultrazvučne dijagnostike kod pasa i mačaka je doprinela poboljšanju pouzdanosti u postavljanju dijagnoza, ali i proširenju primene lekova u različitim patološkim stanjima i predupredila da ordinirajući veterinar sazna šta se dešava u organizmu pacijenta. Mnoge dijagnoze koje se postavljaju u današnje vreme su nekada bile potpuno skrivene i prepuštene više patolozima nego kliničarima.

Kod malih životinja, ultrazvuk se često koristi u dijagnostici graviditeta, broja, veličine i vitalnosti fetusa, kao i za postavljanje dijagnoze patoloških stanja koja se odvijaju u genitalnom traktu, a kod muških još i pregled statusa testisa i prostate (Vogas i sar, 1996). Ultrazvučna dijagnostika nam takođe omogućava uvid u funkcionalno stanje ostalih organa kao što su srce, jetra, pankreas, bubrezi, nadbubrežne žlezde, tireoidea, pluća.

Najveći izazov u ultrazvučnoj dijagnostici predstavlja ogroman raspon u veličini pacijenata koji su predmet istraživanja i dijagnostike u veterinarskoj praksi. Težina pacijenata u maloj praksi varira 0,5 – 75 kg. Samom ovom činjenicom se i nameće pitanje postavljanja objektivne dijagnoze. Kada se upoređuje primena ultrazvučne dijagnostike u humanoj i veterinarskoj praksi uočljivo je da su varijacije daleko veće kada se radi o pregledu malih životinja. Osim toga i građa pasa i mačaka u odnosu na građu ljudskog tela je potpuno drugačija (Marković i sar., 2003). Psi i mačke imaju uzak i dubok trup, grudni koš i abdomen dok su humani pacijenti više pljosnate građe, širokog abdomena. Iako postoji velika razlika koja se može konstatovati ne sme se zaboraviti

da je suština ultrazvučnog pregleda mogućnost apsorpcije ultrazvučnih talasa od strane tkiva različite gustine.

Ključnu ulogu ima i kolagen koji se nalazi na površini kapsula svih unutrašnjih organa, što daje definiciju primene ultrazvučne dijagnostike kod malih životinja i ostavlja mogućnost visoke rezolucije i definicije fizioloških stanja ili promena na unutrašnjim organima, a koji se nalaze na relativno malom prostoru (Marković i sar, 2003). Ova različitost u građi je i presudna kada je u pitanju izbor adekvatnih ultrazvučnih sonde koje se koriste u kliničkoj veterinarskoj praksi (Lukač i sar, 1994).

Sonde koje se koriste u ultrazvučnoj dijagnostici su istog opsega kao i u humanoj medicini. Ultrazvučne sonde koje se koriste u kliničkoj praksi su elektronske i dele se na linearne, sektorske i konveksne. Kod linearnih sonde se baterije kristala aktiviraju jedna za drugom i dobija se slika pravougaonog oblika, po dimenzijama jednaka sonde. Kod elektronskih sektorskih sonde aktiviranje baterija kristala vrši se po posebnom pravilu, na osnovu faznog kašnjenja da bi se dobila karakteristična sektorska slika. Dobre strane su iskorišćene, a nedostaci oba tipa sonde znatno su smanjeni kod konveksnih sonde. Posebnom konstrukcijom ove vrste sonde povećan je ugao pri vrhu -glavni nedostatak sektorske sonde, a otklonjen i nedostatak linearne sonde koja daje samo sliku onog što je neposredno pod njom. Tako konveksna sonda ima pri vrhu veliko vidno polje, a može da vizualizuje i zaklonjene delove. Na svu sreću u današnjoj kliničkoj praksi zahvaljujući tehnološkom napretku nije neophodno posedovati više sonde različite frekvence obzirom da današnji aparati imaju mogućnost regulacije frekvenci sonde u samom aparatu. Potreba za različitim sondama se sastoji samo u potrebi veterinara da želi da radi endorektalnim, kardiološkim ili najčešće transabdominalnim, multifrekventnim sondama kojima mogu da pokriju najveći opseg svojih pregleda (Skrobonja, 2003).

Učestalost utvrđivanja promena na parenhimatoznim organima tokom ultrazvučnog pregleda genitalnog trakta

Ultrazvučni aparati koji se danas koriste u kliničkoj veterinarskoj paraksi imaju širok opseg u zavisnosti od potreba i mogućnosti. Tako se veterinari uglavnom kreću u opsegu aparata koji su manje ili više portabilni i višenamenski. Danas postoje i aparati koji osim 2D *Reall Time*, i aparati koji poseduju opciju primenu kolor dopler ultrasonografije (CD), što predstavlja napredniju tehnološku varijantu, zahtevaju neprekidno izučavanje i usavršavanje i donose napredniji i precizniji pristup u dijagnostici. Kolor dopler predstavlja neinvazivnu evaluaciju protoka kroz krvne sudove genitalnih organa u dvodimenzionalnoj, visoko rezolutoj slici. Identifikacija ili mapiranje krvnih sudova i protoka postiže se promenama u pravcu kretanja talasa u odnosu na sondu. Protok krvi ka sondi mapira se crvenom bojom. Protok krvi od sonde, plavom bojom. Varijacijama u frekvenciji CD talasa postiže se promena u koloru osnovnih boja: Tamnija nijansa boje proporcionalna je manjoj brzini protoka i obrnuto. Laminarni protok boji se čistom crvenom ili plavom bojom. Zelena boja identifikuje postojanje turbulentnog kretanja i to: otvorenija nijansa zelene boje saglasna je većoj turbulenciji. Ukoliko se ona odigrava u pravcu sonde, vizuelizuje se žutom bojom, a nakon insoniranog talasa, nishodno, modrom bojom. (Đukić, 2003)

Reproduktivni organi kao predmet ultrazvučnog ispitivanja se dele na ženske i muške. Zahvaljujući veličini životinje koja se ispituje i rezultat je delikatniji. Pregledač je veoma često u situaciji da se prilagođava na licu mesta veličini životinje koja diktira i opseg, odnosno frekvencu sonde koja se koristi ili dubini na kojoj je organ koji se pregleda. Ultrazvučni pregled je zato najbolje raditi po standardnom protokolu i u istom položaju tela, s obzirom da već postoji delikatan broj varijacija koje pre svega zavise od samog pacijenta i ukoliko pregledač poremeti pristup tokom pregleda, akomodacija oka i rada ruku ga lako mogu odvesti u zabunu i pogrešno tumačenje. Kako se mnogi znalci u ovoj dijagnostičkoj oblasti

izjašnjavaju, aparat nikada ne greši, ali greše oni koji čitaju i tumače sliku na monitoru.

Ono što se smatra komocijom u pregledu je pregled životinje u leđnom položaju ili maksimalno stojećem. Ukoliko se neki od abdominalnih organa ne prikazuje prilikom pregleda, treba osigurati bolju poziciju i raditi van ovih standarda, ali tek nakon što su postavljene neke ključne teze dijagnoze. Ovo se pre svega odnosi na bubrege ili jetru. Obavezno se mora napomenuti da je za pregled srca koje može biti organ pogođen posledicama primarnih patoloških promena na reproduktivnom traktu, neophodna bočna tj. postrana pozicija.

Prilikom ultrazvučnog pregleda genitalnog trakta najbolje je poštovati protokol o pregledu. Ultrazvučni pregled genitalnog trakta kod malih životinja se retko izvodi u uslovima kada je pacijent pripremljen, praznog stomaka i pune bešike (Klein, 1996). To su uslovi koje bi svako poželeo kako bi imao transparentnost mokraćne u pristupu grliću materice i bifurkaciji, a zatim materičnim rogovima i jajnicima bez gasova u crevima koji ometaju normalni pregled. Zbog minimalne apsorptivne moći telesnih tečnosti i bolje transparentnosti tkiva koja se dobija prolaskom ultrazvučnih talasa kroz tkiva različite gustine, eho odjek talasa koji se dobija na monitoru ekrana ultrazvučnog aparata dobija na kvalitetu rezolucije i kod lošijih aparata.

Sam pregled se započinje pripremom, brijanjem dlačnog pokrivača kako se ne bi stvarali vazdušni džepovi koji bi ometali postojanje homogene celine ultrazvučne slike. Kod ženskih i muških pacijenata je najbolje da se nalaze u leđnoj poziciji za pregled, koji se odvija od kaudalnog dela male karlice pa kranijalno duž *linea alba*-e ka sternalnoj kosti. Za obe kategorije pacijenata, ženki i mužjaka ispunjenost mokraćne bešike čini pregled daleko objektivnijim. Puna mokraćna bešika je ključni orijentir za definisanje veličine jetre i slezine. Kod pasa i mačaka s obzirom na jako veliki broj rasa, a i mešanaca, ne postoje standardi koji su prisutni u humanoj medicini. Hepatomegalija kao dijagnoza se definiše kada kaudalni rubovi

lobusa jetre prelaze transversalnu liniju pune mokraćne bešike ka karlici. Isto važi i za splenomegaliju.

U svakodnevnoj kliničkoj praksi vršimo ultrazvučni pregled materice, materičnih rogova i jajnika (Diez i sar, 1998). Mogućnost vizuelizacije ne zavisi od veličine pacijenta nego od relaksiranosti creva od sadržaja i gasova, odnosno ispunjenosti bešike mokraćom. Ovo je najosnovnije pravilo u ultrazvučnoj dijagnostici. Najveći problem veterinarima predstavlja činjenica da je skoro 90% pacijenata nepripremljeno i to čini da su pregledi skoro uvek u zoni interventnog ultrazvuka (Marković i sar, 2003). Povećana ili promenjena prostata i promene na testisima se mogu otkriti rutinskim ultrazvučnim pregledom, čak i u onim slučajevima kada ne dolazi do ispoljavanja karakteristične kliničke slike.

Napredna primena ultrazvučne dijagnostike je dovela do toga da se u današnje vreme slobodno mogu povezati brojna patološka stanja reproduktivnog trakta kako ženskih, tako i muških životinja, koja direktno utiču na zdravstveno stanje ostalih unutrašnjih organa. Kod nekih oboljenja promene koje nastaju su posledica nekad mehaničkih ili metastatskih pojava, funkcionalnih smetnji. Posledice znaju i da se reperkutuju promenama na koži, što često ume da dovede u zabludu kod neiskusnog pristupa u lečenju.

Ženski genitalni trakt

Kod malih životinja je veoma teško i delikatno izvršiti ultrazvučni pregled jajnika, s obzirom da veličina jajnika varira od 0,5 do 2cm, a veličina pacijenata od 0,5 do 75 kilograma (Klein, 1996). Zahvaljujući strukturama kao što su: folikul, žuto telo i ciste žutog tela, možemo u fazi proestrusa, estrusa i metestrusa da identifikujemo i pregledamo jajnike. Jajnici su ovalnog oblika, lokalizovani kaudalno i često bočno u odnosu na kranijalni pol bubrega, koji se koristi kao orijentir za njihovu identifikaciju. Izgled jajnika varira tokom polnog ciklusa.

Normalna, negravidna materica je neupadljiva i teško se vizuelizuje kod malih životinja. Najbolji pristup u prepoznavanju materice je pregled koji započinjemo od vrata ispunjene mokraćne bešike koja nam otvara "prozor" svojom transparentnošću i daje mogućnost vizuelizacije cerviksa, tela materice i bifurkacije. Paralelno sa matericom se pruža vena cava caudalis. Zabuna u zameni ne postoji, s obzirom da se zidovi materice uočavaju hiperehogenim izgledom usled pojačane apsorpcije ultrazvučnih talasa (perimetrijum, miometrijum i endometrijum). Pravilnost kontinuiteta zida je očuvana u proestrusu i estrusu, a u anestrusu su zidovi manje više nepravilnog izgleda. Zidovi vene su ravnomerni i hipoehogeni. Materica se najlakše prepoznaje u kaudalnom abdomenu, gde izgleda kao cevasta struktura između mokraćne bešike (ventralno) i descendentnog kolona (dorzalno) (Lukač i sar, 1994). Veličina materice je promenljiva i zavisi od veličine životinje, broja prethodnih porođaja i faze polnog ciklusa. Usled prisustva creva i njihovog sadržaja matericu i materične rogove možemo da vizualizujemo samo na segmente, nikad u kontinuitetu, ali je to za validan pregled sasvim dovoljno. Za pregledača je bitan promer kao i eventualno prisustvo gnojnog sadržaja koji se karakteriše prisustvom korpuskula ehogenog ili hiperehogenog karaktera. Materica se sastoji iz tri sloja: mukoze, mišićnog sloja i seroze. Endometrijum i miometrijum se obično ne mogu razdvojiti. Periferno je naznačen tanki ehogeni rub. Kod negravidne materice vidljivo je telo, dok se rogovi materice gube u delovima tankog creva i mezenteričnom masnom tkivu. Materica se razlikuje od tankog creva po izostanku peristaltike, nedostatku intraluminalnog gasa i odsustvu slojeva karakterističnih za tanka creva. Patološka stanja koja se dešavaju na ženskom genitalnom traktu su endometritisi različitog tipa, piometra, hemometra, cistična hiperplazija endometrijuma karakterističnog izgleda sličnog brojanicama, tumori jajnika ili ciste jajnika (Klein, 1996).

Muški genitalni trakt

Normalni testisi su ehogeni do hipoehogeni i imaju finu, homogenu ehostrukturu. Testis ograničava tanka, glatka i hiperehogena tunica albuginea. Pri sagitalnim skenovima vidljiva je centralna hiperehogena linija koja predstavlja medijastinum testisa, dok je pri poprečnim skenovima medijastinum vidljiv kao hiperehogeni fokus (Johnston i sar, 1991a). Veličina testisa je direktno povezana sa telesnom težinom. Glava i rep epididimisa nalaze se na kranijalnom i kaudalnom polu testisa, dok je telo epididimisa smešteno dorzalno od testisa. Pri poređenju sa parenhimom testisa, delovi epididimisa su hipoehogeni i grublje strukture.

Lokacija, veličina i izgled prostate zavisi od starosti i od toga da li je pas kastriran ili ne. Zdrava prostata kod mladih i nekastriranih pasa srednje dobi, gradi homogenu sliku, srednje do fine strukture, okružena tankom hiperehogenom kapsulom (Atalan i sar, 1999). Prostata kranijalno graniči sa mokraćnom bešikom, ventralno sa karličnim dnom i ventralnim trbušnim zidom, dok dorzalno graniči sa rektumom. U podužnom skenu prostata je okruglog do ovalnog oblika. Kod polno nezrelih i kastriranih pasa, prostata je znatno manja, relativno hipoehogene i homogene strukture.

Patološke promene na koje najčešće nailazimo kod muških životinja su na testisima hidrocele, varikocеле, tumori i orhitis različite etiologije. Promene koje se mogu identifikovati na prostati su intraprostatične ciste, hiperplazija, tumori, ekstraprostatične ciste.

Abdominalni organi

Položaj jetre zavisi od konstitucije, širine rebarnog luka i veličine jetre. Kaudalno kontura jetre graniči sa konturom desnog bubrega, dok se kranijalno naslanja na dijafragmu. Na položaj mogu uticati promene koje menjaju intratorakalni i intraabdominalni pritisak. Tako je jetra kranijalno pomerena kod tumora abdomena, ascitesa, relaksacije ili rupture dijafragme i atelektaze (Cuccovillo i sar, 2002).

Kaudalno je pomerena kod većeg pleuralnog izliva i hepatomegalije. Određivanje dimenzije i veličine jetre je nepouzdan podatak, jer su oblik i položaj jetre individualno varijabilni. Parenhim jetre je homogenog izgleda. Sastoji se od fino zrnastih ehoa, srednjeg intenziteta, iste jačine u svim delovima jetre. Ipak slabljenje ehoa u dubokim slojevima ne mora biti patološko. Normalna ehogenost jetre je nešto jača u odnosu na ehogenost bubrega, a slabija u odnosu na pankreas. U jetrinom parenhimu se vide okruglaste i tubularne strukture koje predstavljaju krvne sudove. Grane vene porte konvergiraju prema hilusu jetre i njihove konture su hiperehogene. Grane hepatičnih vena konvergiraju prema kranijalno, njihove konture su glatke i nežne bez ehogenosti zidova. Ova diferencijacija je moguća samo kod većih krvnih sudova. Intrahepatični žučni vodovi se ne vide ukoliko njihov lumen nije dilatiran.

Patološke promene su ultrazvučno jasno vidljive, a nastaju kao posledica primarnih oboljenja (*cirosis hepatitis* i fokalne promene) za koje nismo sigurni bez dodatnih pretraga da li su maligne ili benigne prirode. One predstavljaju ograničene promene nepravilnih ovalnih oblika koje svojom hiperehogenošću odudaraju od normalnog parenhima jetre. Njihova prisutnost u jetri može biti solitarna ili multipla. Svojom ukupnom površinom smanjuju funkcionalnu aktivnost organa i utiču na prognozu i dalji tok lečenja. Metastatske promene kao tumorske promene su jasno ograničene, nepravilne strukture, ispresecane hiperehogenim strukturama nalik gredicama između kojih se nalaze kolekcije tečnog anehogenog sadržaja (Mattoon i sar, 1995). Tumori jetre su najčešće lokalizovani u hilusu jetre i retko dostupni tokom pregleda s obzirom da se nalaze u retroperineumu. Tumorske formacije u hilusu jetre posledično dovode do slabosti desnog srca i upućuju na ultrazvučni kardiološki pregled. Prisustvo ovih novonastalih promena dovode do uvećanja tkiva jetre i pomeranja levog lobusa ka *trunkusu celiakusu*, kojeg prepoznamo po anehogenoj senci u oliku "brkova". Patološke

formacije u jetri takođe vrše pritisak i pomeranje levog lobusa ka pankreasu, što se vidi po raširenosti *ductus choledochusa*, a koji ima zajednički izvod sa *papilom major* pankreasa.

Pankreas je organ koji se kod pasa izuzetno teško razaznaje osim u slučajevima navedenih patoloških stanja na jetri ili samom pankreasu.

MATERIJAL I METODE

Ultrazvučni pregled je izvršen na ukupno dvanaest pasa. Šest pasa je bilo muškog, a ostalih šest ženskog pola. Psi su bili starosti između 4 i 8 godina.

Jajnici, materice i abdominalni organi su pregledani sondama u opsegu od 5 do 8 MHz. Ultrazvučni aparat koji smo koristili je Toshiba color dopler, eccocoo, stacionarni aparat sa multifrekventnim sondama, konveksnog i linearnog tipa. Pacijenti su bili položeni u dorzalni ležeći položaj. Za pregled prostate i testisa je korišćena sonda od 7,5 MHz, sa pacijentom položenim u dorzalni ležeći položaj. Pri pregledu jetre ultrazvukom analiziran je položaj, oblik, veličina, konture, parenhim, vaskularizacija i ehostruktura. Za pregled jetre korišćen je longitudinalni, transversalni, kosi i interkostalni presek.

Uzdužni ili longitudinalni presek korišćen je za posmatranje levog i desnog režnja, kupole dijafragme, merenje uzdužnog dijametra, praćenje velikih krvnih sudova, žučnih vodova i holeciste. Ovaj presek je izveden postavljanjem

sonde vertikalno na rebarni luk i pomicanjem u medijalni i lateralni pravac uz blagu rotaciju.

U poprečnom preseku je pregledan levi režanj, merena je njegova veličina, praćen oblik i posmatrana ehostrukturu oba režnja. Sonda je prvo postavljena u transversalni pravac ispod procesusa ksifoideusa, zatim se angulirala u pravac koji je paralelan sa rebarnim lukom, pomerana je kranijalno i kaudalno sa lepezastim rotiranjem pri svakom pomeranju.

Desni kosi presek jetre je vršen vertikalano na rebarni luk, a levi paralelno. Ovim skenovima je pregledan parenhim oba režnja jetre, hilusa i krvnih sudova.

Interkostalni skenovi vršeni su u levom dekubitusu pacijenta pri, mirnoj respiraciji. Sonda je postavljena u bilo koji međurebarni prostor, paralelno sa rebrima i prikazivan je centralni deo jetre sa krvnim suvodima.


Patološke promene jetrinog parenhima podeljene su na difuzne promene, hemangiomi i cirkumskriptne (fokalne) lezije jetre.

REZULTATI

Od pregledanih šest mužjaka, tri su imala promene na prostati. Kod dva psa je dijagnostifikovana hiperplazija prostate, dok je kod jednog psa dijagnostifikovana intraprostatska cista. Kod ovih pasa su takođe primećeni cistitis, hidronefroza i hipertrofija nadbubrega.

Kod jednog mužjaka je utvrđen tumor testisa. Pored tumora su dijagnostifikovani hiperplazija prostate i cistitis. Preostala dva mužjaka nisu imala patološke promene na genitalnim organima.


Учесталост утврђивања промена на parenhimatozним organима током ултразвучног прегледа гениталног тракта



Графикон 1. Резултати ултразвучног прегледа мушјака

Од шест прегледаних женки једна је имала цистићну хиперплазију ендометријума, три су имале пиометру, док је код једне дијагностикована фолкуларна циста јajника. Поред ових патолошких

стања, утврђене су промене на јетри и панкреасу. Једна женка није имала патолошке промене на гениталним organима.



Графикон 2. Резултати ултразвучног прегледа женки

DISKUSIJA

Цисте и неоплазме представљају два најчешћа патолошка стања јajника (Marković и sar, 2003).

У цистама долази до стварања естрогена, који се депоњује у организму и делује као канцерогени

hormon (Klein, 1996). Target organ za nastanak tumora predstavljaju prvenstveno mlečna žlezda, jetra, pluća, pankreas, slezina. Tumorozne tvorevine mogu biti benigne ili maligne prirode, ali se najčešće javljaju maligni karcinomi, sarkomi i adenokarcinomi. Primarni tumori jajnika mogu poticati od primitivnog kličinog epitela, stromalnih ćelija i od površinskog celomskog epitela (Diez i sar, 1998). Svi stromalni tumori polnih žlezda dele sklonost da budu hormonski aktivni i tako produkuju znakove hiperestrogenizma, maskulinizacije ili upornog anestrusa.

Fiziološki u toku ovulacije na jajovodima i jajnicima često dolazi do pojave hemoragija, čija je dijagnostika veoma komplikovana. Na mestima gde su se pojavile hemoragije, može da dođe do formiranja priraslica usled prisutne vaskularne tečnosti. Priraslice nastaju između jajnika (burze ovarike) i omentuma (Cruz-Arambulo i sar, 2004). Kada dođe do pojave priraslica, često dolazi do unilateralne pojave tumora jajnika. Novonastali tumor jajnika kompromituje omentum i vezujući se za njega nastavlja svoj razvoj. Širenje tumora najčešće nastaje trascelomično, nakon rupture cista unutar neoplazme ili invazije tumora kroz kapsulu jajnika (Meuten, 2017). Posledično može da nastane ascites i abdominalna distenzija izazvana limfnom blokadom. Ovakvi procesi najčešće metastaziraju jer su naslonjeni na velike krvne sudove omentuma, koji je bogato vaskularizovan i povezan sa digestivnim traktom. Jetra je najčešća meta metastaza, uglavnom kroz portalni sistem koji drenira većinu abdominalnih struktura (Nyland i sar, 1983). Kod pasa se metastatske neoplazme jetre približno tri puta češće sreću u odnosu na primarne neoplazme. Varijabilne karakteristike tkiva kod primarnih i metastatskih neoplastičnih procesa uključujući: gustinu tkiva, vaskularni crtež, nekrozu, tečnost i kalcifikaciju, uzrokuju varijabilan ultrasonografski prikaz (Klein, 1996). Fokalne hipoehogene lezije sa hiperehogenim centrom, nazvane target lezije, obično predstavljaju metastaze. Ranije studije

pokazuju da kod nalaza najmanje jedne target lezije u jetri ili slezini, postoji pozitivna korelacija od 74% za malignitet kod malih životinja (Cuccovillo i sar, 2002). Ostale osobine koje ukazuju na malignitet su veličina lezije (više od 3 cm) i prisustvo peritonelanog izliva. Dodatne karakteristike za razlikovanje malignih od benignih jetrinih neoplazmi se mogu ustanoviti putem CT-a, kao i tumormarkerima.

Najčešća patološka stanja materice čine: cistična hiperplazija endometrijuma, mukometra, hemometra, hidrometra, piometra, segmentna piometra, granulomi i tumori materice (Voges i sar, 1996). Cistična hiperplazija endometrijuma je hormonske etiologije i obično prethodi piometri. Intraluminalno se može pojaviti sterilna sluzava tečnost, koja uzrokuje hidrometru ili mukometru, zavisno od stadijuma hidratacije mukusa. Kod kuja cistična hiperplazija endometrijuma nastaje pod uticajem progesterona, ali njena inicijacija nastaje pod dejstvom estrogena iz jajnika (Klein, 1996). Zbog dejstva progesterona, endometrijum je osetljiv na infekcije, što dovodi do toga da cistična hiperplazija endometrijuma prelazi u piometru. Promene van genitalnog trakta koje prate piometru ispoljavaju se u vidu izražene kaheksije, poremećaja metabolizma i glomerulonefritisa prouzrokovanog imunskim kompleksima (Klein, 1996). Ove sekundarne promene su naročito zapažene kod kuja i mogu dovesti do intoksikacije sa fatalnim posledicama.

Lejomijomi predstavljaju najčešće tumore materice kuja. Javljaju se kao multiple neoplazme, ne samo u uterusu već i u cerviksu i vagini. Često su udruženi sa drugim poremećajima, kao što su cistična hiperplazija endometrijuma, folikularne ciste i tumori mlečne žlezde. Estrogen najverovatnije ima značajnu ulogu u nastajanju i održavanju ove neoplazme kod kuja. Tumori materice kuja, naročito starijih od 8 godina, su često malignog karaktera i daju metastaze na mlečnoj žlezdi (Mattoon i sar, 1995). Tumori mlečne žlezde

Učestalost utvrđivanja promena na parenhimatoznim organima tokom ultrazvučnog pregleda genitalnog trakta

predstavljaju jedne od najčešćih tumora kod kuja, a najviše pažnje je posvećeno uticaju stereoidnih hormona na tumorogenezu. Receptori za estrogen, progesteron i prolaktin otkriveni su na normalnim i tumorskim ćelijama mlečne žlezde. Oko 40 do 60% benignih i malignih tumora mlečne žlezde su pozitivni na ove receptore. Rana kastracija, do starosti od 2 godine, ima zaštitni efekat i smanjuje verovatnoću pojave karcinoma mlečne žlezde kod kuja.

Od patoloških stanja na testisima najzastupljenije su neoplazme. Pored njih često se dijagnostikuju: orhitis, epididimitis, granulom, apsces, hematoma, hidrocele, varikokele, kriptorhizam i skrotalna hernija (Johnston i sar, 1991a). Tumori testisa mogu da vode poreklo iz germinativnog epitela (seminom i teratom) ili intersticijalnih ćelija (tumor Sertolijevih ćelija i tumor Lajdigovih ćelija). Seminomi se često javljaju kod kriptorhida i pokazuju lokalnu invazivnost, dok su teratomi benigni i javljaju se pretežno kod mladih životinja. Tumor Sertolijevih ćelija se najčešće sreće kod pasa, daje metastaze u unutrašnjem ilijačnom limfnom čvoru. Tumorske ćelije mogu da produkuju estrogen i da kod muških životinja dovedu do hiperestrogenizma koji se manifestuje ginekomastijom i alopecijom, dok se u pojedinim slučajevima razvija i aplazija kostne srži (Hart i sar, 2013).

Slika orhitisa je veoma slična neoplazmama. Često je prisutna ekstraparastatična tečnost i povećanje epididimisa.

Benigna hiperplazija prostate, prostatitis, apsces, prostatična i paraprostatična cista, kao i neoplazme čine najčešća patološka stanja ove akcesorne polne žlezde (Johnston i sar, 2000). Hiperplazija prostate je često oboljenje starijih pasa koje nastaje delovanjem androgenih i estrogenih hormona. Vremenom dolazi do fibroziranja tkiva, što posledično dovodi do kompresije uretre i otežanog mokrenja. Unutar parenhima mogu biti prisutne ciste različitog broja i veličina. Solitarne ciste mogu biti dovoljno velike da dovedu do opstrukcije

uretera, a posledično i do hidronefroze (Lukač i sar, 1994).

Neoplazme prostate uglavnom su maligne prirode i u velikom postotku metastaziraju u regionalne limfne čvorove, kosti i pluća (Atalan i sar, 1999). Za razliku od benigne hiperplazije, dolazi do razaranja kapsule i širenja u okolno tkivo. Najčešće se javljaju adenokarcinomi i nediferencirani karcinomi.

Patološke promene jetrinog parenhima dele se na difuzne promene jetre, hemangiome i cirkumskriptne (fokalne) lezije jetre (Penninck i sar, 2013). Difuzne lezije mogu biti inflamatorne, metaboličke i vaskularne. Fokalne lezije jetre mogu biti primarni i benigni tumori. Vaskularni tumori su najčešći primarni tumori jetre uopšte. Kapilarni hemangiomi predstavljaju hiperehogenu fokalnu leziju, odlično ograničenu od okoline bez hipoehogenog haloa, koji se obično viđa u malignim tumorima, mada mogu biti i prolaznog karaktera, kao i da se pojave na novom mestu u parenhimu (Cruz-Arambulo i sar, 2004). Najčešće su okruglasti. Ponekad mogu biti hipoehogeni sa sitnim, brojnim internim ehoima. Adenomi su okrugle fokalne promene koje mogu biti i hiper i hipoehogene. Često su multipli i onda mogu da pokažu različitost ehostrukture. Nodularna hiperplazija je obično hiperehogena. Ovi benigni tumori su inkapsulirani i imaju dobre ograničene ivice. Često su prostrani i mogu biti multipli. Poneki su izoehogeni. Maligne tumore jetre čine: hepatocelularni karcinomi, fibrolamelarni hepatokarcinomi i holangiokarcinomi (Marković i sar, 2003). Ehogenost hepatokarcinoma zavisi od njegove tkivne strukture, odnosno od toga da li je bogat krvnim sudovima, mastima ili vezivnim septama. Bogatstvo u mastima daje hiperehogeni izgled. Ako se radi o solitarnom tumoru obično je veoma krupan, režnjevito. Tumorske mase su dosta dobro ograničene od okoline, sa hipoehogenim haloom. Hepatokarcinom ima tendenciju invazije vena. Hepatokarcinomi su

dobro vaskularizovani i mogu da krvare spontano.

Metastaze u jetri su najčešće multinodularne, brojne, ali postoje i solitarne (Cuccovillo i sar, 2002). Kod metastaza u jetri nalazimo promene u konturama jetre i abnormalnosti tkiva. U metastazi se ultrazvukom ponekad može videti kalcifikacija u trenutku kada se ona još ne vidi na rentgenu. Ovakve kalcifikacije se najčešće vide u mucinoznim adenokarcinomima kolona, ili u karcinomu ovarijuma. U metastazi može da se razvije nekroza. U tom slučaju govorimo o nekrotičnim metastazama. Nekroza može da se dogodi spontano u koliko veličina metastaze pređe 6-7 cm. Dilatacija intrahepatičnih žučnih puteva može pratiti metastaze (Lukač i sar, 1994). Do ovoga dolazi zbog kompresije izazvane metastazama. Hepatične vene takođe, mogu da promene izgled zbog metastaza. Vene uz metastaze su izmenjene, deformisane ili razorene. Može doći i do komprese ili invazije portne vene i njenih grana. Eksperimentalne studije pokazuju da stepen vaskularizacije metastaza utiče na njen izgled (Cuccovillo i sar, 2002). Uopšte hipervaskularne lezije su ehogene, a hipovaskularne su anehogene. Najčešće se uz ove promene vidi i

splenomegalija i retroperitonealna limfadenopatija.

Kao posledica intoksikacija, poremećaja metabolizma, hronične kongestije i nekih infektivnih agenasa nastaje ciroza jetre. U diferencijaciji ciroze od metastatskih nodusa mora se paziti na koeficijent atenuacije i konture jetre. U cirozi je atenuacija pojačana, a konture postaju bikonveksne (Stowater i sar, 1990). Veličina jetre može biti od pomoći. U cirozi jetra se postepeno smanjuje, a u metastatskim promenama jetra je rapidno povećana. Povećana atenuacija vidi se samo u cirozi. Splenomegalija i dilatirana v. lienalis javlja se samo u cirozi.

Portna hipertenzija nastaje kao rezultat intra i ekstrahepatičneotežane portne cirkulacije, zbog ciroze, fibroze, steatoze, primarnih i metastatskih tumora (Lukač i sar, 1994). Zastoj u portnoj cirkulaciji dovodi do primarnog uvećanja jetre, promene strukture parenhima i proširenja vene porte i njenih grana. Sekundarno se razvija splenomegalija, kolateralna cirkulacija i ascites.

Takođe, metaboličke bolesti jetre su praćene promenom veličine jetre, ponekad i ehostrukture parenhima.

ZAKLJUČCI

Ultrazvučnom dijagnostikom se pored primarnih promena na genitalnom traktu, kako mužjaka tako i ženki, mogu vizuelizovati i posledične promene na parenhimatoznim organima.

Promene na jetri kao posledica primarnog oboljenja reproduktivnog trakta ženki mogu biti *cirrhosis hepatis*, kao i fokalne promene koje izazivaju hepatomegaliju i mehanički ugrožavaju *pancreas*.

Kod mužjaka tumori prostate izazivaju inkontinenciju urina, stvaranje fibrinskih

naslaga u početku kao polipa koji kalcifikuju, do stvaranja komora sa zaostalim mokraćom, sačastog izgleda, dilatacijom bešike koja ne može da se prazni.

Zahvaljujući ultrazvučnoj dijagnostici su se proširila znanja u veterini kao i opseg primene lekova.

Glavni izazov ordinirajućeg veterinara je da zahvaljujući ultrazvučnoj dijagnostici pretpostavi primarno oboljenje, a ne da leči posledična.

LITERATURA

1. Atalan G, Holt PE, Barr FJ (1999) Ultrasonographic estimation of prostate size in normal dogs and relationship to bodyweight and age, *J Small Anim Pract*, 40:119-122.

Učestalost utvrđivanja promena na parenhimatoznim organima tokom ultrazvučnog pregleda genitalnog trakta

2. Barella G, Lodi M, Sabbadin LA, Faverzani S (2012) A new method for ultrasonographic measurement of kidney size in healthy dogs *J Ultrasound*, 15:186–191.
3. Barthez PY, Nyland TG, Feldman EC (1995) Ultrasonographic evaluation of the adrenal glands in dogs, *J Am Vet Med Assoc*, 207:1180-1183.
4. Cruz-Arambulo R, Wrigley R, Powers B (2004) Sonographic features of histiocytic neoplasms in the canine abdomen, *Vet Radiol Ultrasound*, 45:554–558.
5. Cuccovillo A, Lamb CR (2002) Cellular features of sonographic target lesions of the liver and spleen in 21 dogs, *Vet Radiol Ultrasound*, 43:275–278.
6. Diez-Bru N, Garcia-Real I, Martinez EM, Rollan E, Mayenco A, Llorens P (1998) Ultrasonographic appearance of ovarian tumors in 10 dogs, *Vet Radiol Ultrasound*, 39:226–233.
7. Feeney DA, Johnston GR, Klausner JS, Bell FJ (1989) Canine prostatic ultrasonography, *Semin Vet Med Surg (Small Anim)*, 4:44–57.
8. Hecht S, Henry G (2007) Sonographic evaluation of the normal and abnormal pancreas, *Clin Tech Small Anim Pract*, 22:115–121.
9. Johnston GR, Feeney DA, Johnston SD, O'Brien TD (1991a) Ultrasonographic features of testicular neoplasia in dogs: 16 cases (1980–1988), *J Am Vet Med Assoc*, 198:1779–1784.
10. Johnston SD, Kamolpatana K, Root-Kustritz MV, Johnston GR (2000) Prostatic disorders in the dog, *Anim Reprod Sci*, 60–61:405–415.
11. Kantrowitz BM, Nyland TG, Feldman EC (1986) Adrenal ultrasonography in the dog, *Vet Radiol*, 27:91-96.
12. Klein MK (1996) Tumors of the female reproductive system. In: Withrow SJ, MacEwen EG, eds, *Small Animal Clinical Oncology*, Philadelphia: WB Saunders, pp 347–355
13. Lukač I, Kovačević N (1994) Dijagnostički ultrazvuk u gastroenterologiji i nefrologiji, Beograd, Dunaj.
14. Marković A (2003) Ultrazvuk u medicini, Udruženje za primenu ultrazvuka u medicini, Biologiji i veterini Srbije, 360-375,
15. Mattoon JS, Nyland TG (1995) Ovaries and uterus, In: Nyland TG, Mattoon JS, eds, *Small Animal Diagnostic Ultrasound*, Philadelphia: WB Saunders, pp 231–249.
16. Meuten DJ (2017) Tumors in domestic animals, Collage of Veterinary Medicine North Carolina State University Raleigh, NC USA, Fift edition, 602-615, 689-702.
17. Nyland TG, Park RD (1983) Hepatic ultrasonography in the dog, *Vet Radiol*, 24:74-84.
18. Older RA, Van Moore A Jr, Glenn JF, Hidalgo HJ (1984) Diagnosis of adrenal disorders, *RCNA*, 22:433-454.
19. Penninck DG, Zeyen U, Taeymans O, Webster CR (2013) Ultrasonographic measurement of the pancreas and pancreatic duct in clinically normal dogs, *Am J Vet Res* 74:433–437.
20. Penninck DG, d' Anjou, MA (2015) *Atlas of Small animal ultrasonography*, Wiley Blackwell, Second Edition, 183-195, 311-314, 403-411.
21. Stowater JK, Lamb CR, Schelling SH (1990) Ultrasonographic feature of canine hepatic nodular hyperplasia, *Vet Radiol*, 31:268-272
22. Trigo, FJ, Thompson, H (1982) The pathology of liver tumours in the dog, *J Comp Pathol*, 92:21–39.
23. Voges AK, Neuwirth L (1996) Ultrasound diagnosis: Cystic uterine hyperplasia, *Vet Radiol Ultrasound*, 37:131–132.

Rad primljen: 28.12.2018.
Rad prihvaćen: 07.02.2019.

DOI 10.7251/VETJEN1901184P

UDK 636.7/.8.09:618.3-073.432.19

*Original scientific paper***FREQUENCY OF DETERMINATION OF PARENCHYMAL ORGANS DISTURBANCES DURING ULTRASOUND EXAMINATION OF REPRODUCTIVE TRACT******Strahinja ĆIBIĆ^{1*}, Miloš PAVLOVIĆ²**

1 Strahinja Ćibić, Faculty of Veterinary Medicine, University of Belgrade, Serbia

2 Prof.dr Miloš Pavlović, Department of Obstetrics, Reproduction and Artificial Insemination, Faculty of Veterinary Medicine, University of Belgrade, Serbia

* contact person e-mail: Strahinja Ćibić: strahinjacicibic@yahoo.com

Abstract: Ultrasound *Real time 2D* diagnostics, being an available and non-invasive method, is successfully used in human medicine, veterinary medicine and biology to diagnose both physiological and pathological conditions. In the case of small animals, the ultrasound is often used to diagnose gravidity, number, size and vitality of fetuses as well as to diagnose pathological conditions present in the genital tract, and is also used to examine testicles and prostate of male animals. Today's advanced use of the ultrasound diagnostics provides us with an opportunity to detect numerous pathological conditions of reproductive tract of both female and male animals, and those conditions directly affect health status of other internal organs. The aim of this study is to diagnose primary genital tract disorders using the ultrasound diagnostics, as well as proving its connection to the changes in parenchymal organs. The ultrasound examination was performed on the total of 12 dogs, 6 male and 6 female dogs. Ovaries, uterus and the organs in the abdominal area were examined using a transducer ranging from 5 to 8 MHz. A transducer of 7.5 MHz was used to examine prostates and testicles. The patients were in dorsal position. Three out of six examined male dogs exhibited changes on their prostates, and the remaining one had a tumor on the testicles. Cystitis and hyperplasia of prostate were also found upon the examination of these dogs. One out of six examined female dogs, had cystic endometrial hyperplasia, three of them had pyometra, and one of them was diagnosed with an follicular cyst. In addition to these pathological conditions, changes were noticed in the liver and pancreas. Along with the detection of primary changes in the genital tract of both male and female animals, the ultrasound diagnostics can be used to discover consequent changes in parenchymal organs.

Key words: ultrasound, genital tract, parenchymal organs, dogs

** Work is presented on the 23rd Annual Counselling of Doctors of Veterinary Medicine of Republic of Srpska (B&H) with International participation, Teslić 2018.

INTRODUCTION

Ultrasound Real Time 2D Diagnostics, as an available and non-invasive method, is successfully used in medicine, veterinary and biology, for the diagnosis of both physiological and pathological conditions (Marković et al., 2003). The use of this method was pioneering in the early diagnosis of pregnancy in the eighties and then generally accepted in human medicine, which the physicians introduced in experiments and everyday clinical practice. Today, this diagnostic procedure is almost irreplaceable in everyday clinical practice. The best description of this method is that it represents a stethoscope of doctors and veterinarians of the 21st century. Massive application of ultrasound diagnostics brought about facing very high prices of the device, but massification and technological advancement made this method available in wide practice (Lukač et al, 1994). As for veterinary clinical practice, ultrasound has made revolutionary progress both in large animals and small ones. The use of ultrasound diagnostics in dogs and cats has improved diagnostic reliability, broaden the use of drugs in various pathological conditions and allowed a veterinarian to find out what is happening in the patient's body. Many of the diagnoses that are being made today were once completely concealed and left more to pathologists than clinicians. In small animals, ultrasound is often used in the diagnosis of gravity, number, size and vitality of the fetus, as well as for the diagnosis of pathological conditions occurring in the genital tract, and prostate and testicular examination in males (Vogas et al, 1996). Ultrasound diagnostics also allows us to inspect the functional state of other organs such as the heart, liver, pancreas, kidneys, adrenal glands, thyroid, lungs.

The greatest challenge in ultrasound diagnostics is the huge range in the size of patients who are the subject of research and diagnostics in

veterinary practice. The weight of patients in small animal practice varies from 0.5 to 75 kg. This fact itself poses the question of setting up an objective diagnosis. When comparing the use of ultrasound diagnostics in human and veterinary practice, it is noticeable that variations are far greater when examining small animals. In addition, the structure of dog's and cat's bodies in relation to the human body construction is completely different (Marković et al., 2003). Dogs and cats have a narrow, deep trunk, chest and abdomen while humans are more plump with broad abdomen.

Although there is a large difference, one must not forget that the essence of the ultrasound examination is the ability to absorb ultrasound waves from different tissue densities. The collagen on the surface of the capsules of all internal organs plays a key role, which defines the application of ultrasound diagnostics in small animals, and leaves the possibility of high resolution and definition of physiological conditions or changes in internal organs, which are located in a relatively small area (Marković et al, 2003). This diversity in body construction is also crucial when it comes to the selection of adequate ultrasonic probes used in clinical veterinary practice (Lukač et al, 1994).

The probes used in ultrasonic diagnostics have the same circumference as in human medicine. Ultrasonic probes used in clinical practice are electronic and are divided into linear, sectoral and convex probes. In linear probes, the crystal batteries are activated one after the other, the image produced is rectangular in shape, with the dimensions equal to the probe. In electronic sector probes, the activation of crystal batteries is carried out according to a special rule, based on phase delay in order to obtain a characteristic sector image. In convex probes good sides are used and the defects of both types of probes are significantly reduced. A special construction of

this type of probes has increased the angle at the top - the main lack of a sector probe, and the lack of a linear probe which gives only the image of what is directly below it has also been eliminated. The convex probe has a large field of view at the top, and it can also visualize hidden parts.

Fortunately, in today's clinical practice, due to technological advancement, it is not necessary to have multiple probes of different frequencies since today's devices have the ability to regulate the probe frequency in the device itself. The need for different probes consists only in the need of a veterinarian to work with endorectal, cardiological or most often transabdominal, multi-frequency probes that can cover the largest extent of his examination (Skrobonja, 2003).

Ultrasound devices that are used today in clinical veterinary practices have a wide range depending on the needs and possibilities. Thus, veterinarians generally use devices that are more or less portable and multifunctional. Today, there are devices that, besides 2D Real Time, have the option of using color doppler ultrasonography (CD), which is a more advanced technological variant. They require continuous learning and training and bring a more advanced and more accurate approach in the diagnostic process. Color Doppler presents a non-invasive evaluation of the flow through blood vessels of the genital organs in a two-dimensional, high-resolution image. Identification or mapping of blood vessels and blood flows is achieved by changes in the direction of wave motion in relation to the probe. The blood flow toward the probe is mapped in red. Blood flow from the probe in blue. Variations in the frequency of the CD waves make a change in the color of the basic colors: Darker color shade is proportional to the lower flow rate and vice versa. The laminar flow is marked with red or blue color. The green color identifies the existence of turbulent movement: lighter shades of green color are concordant with greater turbulence. If it takes place toward the probe, it is visualized in

yellow, and in indigo blue after insonated wave. (Đukić, 2003)

Reproductive organs as subjects of ultrasound examination are divided into female and male reproductive organs. The result is more delicate due to the size of the animal being examined. The examiner very often has to adapt to the size of the animal which dictates the range and the frequency of the probe being used or the depth at which the body is examined. Ultrasound examination is therefore best done by the standard protocol and in the same position of the body, since there is already a delicate number of variations that primarily depend on the patient itself, and if the examiner disrupts the approach during the examination, the adjustment of the eye and the work of the hand can easily lead to confusion and misinterpretation. As many experts in this diagnostic area claim, the device never fails, but those who read and interpret the image on the monitor may be wrong. Examining the animal in the back or standing position provides the greatest commotion during the examination. If some of the abdominal organs are not displayed during the examination, a better position should be provided and work out of these standards, but only after some key elements of diagnosis are set. This primarily refers to kidneys or liver. It is mandatory to note that for the examination of the heart that can be an organ affected by the consequences of primary pathological changes on the reproductive tract, lateral decubitus position is necessary.

In the ultrasound examination of the genital tract, it is best to comply with the examination protocol. Ultrasound examination of the genital tract in small animals is rarely performed under conditions where the patient is prepared, with empty stomach and full bladder (Klein, 1996). These are the conditions anyone would want in order to have urine transparency in the access to the cervical os and bifurcation, and then the uterine horns and ovaries without interfering intestinal gasses. Due to the minimal absorptive power of body fluids and better tissue

transparency obtained when ultrasonic waves pass through tissues of different densities, the echo of the wave received on the monitor screen of the ultrasonic device gets on the quality of resolution even with poorer devices.

The examination itself begins with the preparation, shaving of the hairy covering to prevent creation of air pockets that would disturb the existence of a homogeneous whole of the ultrasound image. For female and male patients it is best to be lying on back during the examination which occurs from the caudal portion of the pelvic cavity and cranially along the linea alba to the sternal bone. For both categories of patients, females and males, bladder fullness makes the examination far more objective. Full bladder is a key landmark for defining the size of the liver and spleen. In dogs and cats, given the very large number of breeds and mixed breeds, there are no standards present in human medicine. Hepatomegaly as a diagnosis is defined when the caudal edges of the liver lobes cross the transversal line of the full bladder to the pelvis. The same applies to splenomegaly.

In everyday clinical practice we perform ultrasound examination of the uterus, uterus horns and ovaries (Diez et al, 1998). The possibility of visualization does not depend on the size of the patient, but on the relaxation of the intestine from the contents and gases or bladder fullness. This is the most basic rule in ultrasound diagnostics. The biggest problem for veterinarians is the fact that almost 90% of patients are unprepared, and that makes examinations almost always in the zone of interventional ultrasound (Marković et al, 2003). Increased or altered prostate and changes in the testicles can be detected by routine ultrasound examination, even in those cases where the characteristic clinical picture does not occur. The advanced application of ultrasound diagnostics has led to the possibility that today many pathological conditions of the reproductive tract

of both female and male animals, which directly influence the health condition of other internal organs, can be linked. In some diseases, changes that occur are the result of mechanical or metastatic occurrences, functional disturbances. The consequences are also known to be reflected by changes in the skin, which can often be misled by inexperienced treatment.

Female genital tract

In small animals it is very difficult and delicate to perform an ultrasound examination of the ovaries, since the size of the ovaries varies from 0.5 to 2cm, and the size of the patients is from 0.5 to 75 kilograms (Klein, 1996). Thanks to structures such as follicle, yellow body and corpus luteum cyst, we can identify and examine the ovaries at the stage of proestrus, estrus and metestrus. The ovaries are oval in shape, located caudally and often laterally to the cranial pole of the kidney, which is anatomic landmark for localizing the ovary. The appearance of the ovary varies during the cycle.

A normal, non-gravid uterus is unobtrusive and difficult to visualize in small animals. The best way to recognize the uterus is an examination that begins from the neck of the filled bladder that opens the "window" with its transparency and provides the possibility of visualizing the cervix, the uterine body and bifurcation. There is also the vena cava caudalis. They cannot be mistaken one for another since the uterine walls are recognized by hyperechoic look due to increased absorption of ultrasonic waves. (perimetry, myometrium and endometrium). The regularity of wall continuity is preserved in proestrus and estrus, while during anestrus the walls are more or less irregular. The walls of the vein are uniform and hypoechogenic. The uterus is most easily recognized in the caudal abdomen, where it looks like a tubular structure between the urinary bladder (ventral) and the descending colon (dorsal) (Lukač et al, 1994).

The size of the uterus is variable and depends on

the size of the animal, the number of previous deliveries, and uterine cycle phase. Due to the presence of the gut and their contents, the uterus and uterine horns can be visualized only in segments, never in continuity, but this is enough for a valid examination. The diameter is important for the examiner, as well as the possible presence of purulent contents, which is characterized by the presence of corpuscles of the echogenic or hyperechogenic character. The uterus consists of the following 3 tissue layers: the internal mucosa, the muscularis, and the serosa. Endometrium and myometrium usually can not be separated. The thin echogenic rim is visible peripherally. In non-gravid uterus the body is visible, while the horns of the uterus are lost in the parts of the small intestine and the mesenteric fat tissue. The uterus differs from the small intestine by the absence of peristalsis, the lack of intraluminal gas and the absence of layers characteristic for the small intestine. Pathological conditions that occur on the female genital tract are different kinds of endometriosis: pyometra, hemometra, cystic endometrial hyperplasia with characteristic rosary-like appearance, ovarian tumors and ovarian cysts. (Klein, 1996).

Male genital tract

Normal testicles are echogenic to hypoechogenic and have a fine, homogeneous echo structure. The testis is covered by a thin, hyperechoic tissue layer known as the tunica albuginea. In sagittal scans a central hyperechoic line that represents the mediastinum testis is visible, whereas in transverse scanning the mediastinum appeared as a hyperechoic focus. Testicle size is directly related to body weight. The head and tail of the epididymis are located on the cranial and caudal pole of the testis, while the epididymal body is located along the dorsal aspect of the testicle. Compared to the parenchyma of the testis, epididymis are hypoechoic and have coarser structure. The location, size and appearance of the prostate

depends on the dog's age and whether the dog is castrated or not. The healthy prostate in young and uncastrated dogs of the middle age builds a homogeneous, medium to fine structure, surrounded by a thin, hyperechogenic capsule (Atalan et al., 1999). The prostate is bordered cranially by the bladder, ventrally by the pelvic floor and ventral abdominal wall, and dorsoally by the rectum. In the longitudinal scan the prostate is round to oval. In sexually immature and neutered dogs, the prostate is significantly smaller, with relatively hypoechoic and homogeneous structure. The pathological changes in testis most commonly encountered in male animals are hydrocele, varicocele, tumors and orchitis of different etiologies. The changes that can be identified in the prostate are intraprostatic cysts, hyperplasia, tumors, extraprostatic cysts.

Abdominal organs

The liver position depends on body built, the breadth of the ribs and the liver size. The contour of the liver is caudally bordered by the right kidney contour, and cranially by the diaphragm. Changes in intrathoracic and intraabdominal pressures can affect the position. Thus, the liver is cranially shifted in case of abdominal cancer, ascites, relaxation or rupture of the diaphragm and atelectasis (Cuccovillo et al., 2002). It is caudally shifted in higher pleural effusion and hepatomegaly. Determining the size of the liver is unreliable because the shape and position of the liver are individually variable. The liver has homogeneous parenchyma. It consists of fine grainy echoes, of medium intensity and the same strength in all parts of the liver. However, attenuation of echoes in deep layers does not have to be pathological. Normal liver echogenicity is somewhat stronger than echogenicity of the kidney, and less than that of the pancreas. In the liver parenchyma there are circular and tubular structures that represent blood vessels. Portal vein branches converge toward liver hilus and their walls are

hyperechoic. Hepatic vein branches converge cranially, their contours are smooth and gentle without the echogenicity of the vein walls. This differentiation is possible only in larger vessels. Intrahepatic bile ducts are not seen if their lumen is not dilated.

Pathological changes are clearly visible in the ultrasound, and are the result of primary diseases (cirrhosis hepatis and focal changes) for which we are unsure of whether they are malignant or benign. They represent limited changes in irregular oval shapes which, by their hyperechogenicity, deviate from normal hepatic parenchyma. Their presence in the liver may be solitary or multiple. With their total surface they reduce the functional activity of the organ and affect the prognosis and further course of treatment. Metastatic changes as tumor changes are clearly limited, irregular structures, containing hyperechogenic structures like the

beds between which there are collections of liquid anechoic content (Mattoon et al., 1995). Liver tumors are most commonly localized in the liver hilus and are rarely available during the examination given that they are in the retroperinum. Tumor formation in the liver hilus eventually leads to weakness of the right heart and point to an ultrasound cardiologic examination. The presence of these newly formed changes leads to the enlargement of the liver tissue and the displacement of the left lobe to the celiac trunk which is recognized by the anechoic shadow in the shape of "mustache". Pathological formation in the liver also exerts pressure and shift the left lobe to the pancreas, which can be seen by dilatation of common bile-duct which has the same opening, papilla major, as pancreatic duct. Pancreas is an organ that is extremely difficult to detect in dogs except in the cases of these pathological conditions of the liver or the pancreas.

MATERIAL AND METHODS

Ultrasound examination was performed on a total of twelve dogs. The six dogs were male and the other six females. Dogs were between 4 and 8 years old. The ovaries, uterus and abdominal organs were examined with the probes in the range of 5 to 8 Mhz. We used Toshiba color doppler ultrasound, a stationary device with multi-frequency probes, convex and linear type. The patients were placed in dorsal lying position. A probe of 7.5 MHz was used to examine the prostate and the testicles, with the patient placed in dorsal lying position. Position, shape, size, contours, parenchyma, vascularisation and echostructures of the liver were examined with ultrasound. Longitudinal, transverse, oblique and intercostal cross-sections were used for the examination of the liver. Longitudinal cross-section was used for observation of the left and right lobes, diaphragm cupola, for longitudinal diameter measurement and for observation of large blood vessels, bile ducts and gallbladder. This cross-


section is made by placing the probe vertically on the costal margin and moving in the medial and lateral direction with slight rotation. The left lobe was examined using cross section, its size was measured and the shape and echostructure of both lobes was observed. The probe was first positioned in the transverse direction below the xiphoid process then it was angulated in a direction parallel to the costal margin, and shifted cranially and caudally with fan-like rotation at each shift. Right oblique scan of the liver was performed vertically on the costal margin and the left one parallel to it. Liver parenchyma of both lobes was examined as well as gallbladder and blood vessels. Intercostal scans were performed in the left lateral decubitus of the patient in a quiet respiration. The probe was placed in any intervertebral space, parallel to the ribs, and the central part of the liver with blood vessels was shown. Pathological changes in the liver parenchyma are divided into diffuse changes, hemangiomas,

and focal liver lesions.

RESULTS

Of the six males examined, three had changes of the prostate. Prostatic hyperplasia is diagnosed in two dogs, while intraprostatic cyst is diagnosed in one dog. Cystitis, hydronephrosis and adrenal gland hypertrophy were also


observed in these dogs. Testicular cancer was found in one male. In addition to tumors, prostatic hyperplasia and cystitis are diagnosed. The remaining two males did not have pathological changes in the genital organs.



Graph 1. Results of ultrasound examination of male dogs

Of the six examined females, one had cystic hyperplasia of the endometrium, three were diagnosed with pyometra, while one was diagnosed with follicular cyst of the ovary. In

addition to these pathological conditions, changes in the liver and pancreas have been identified. One female had no pathological changes in the genital organs.



Graph 2. Results of ultrasound examination of female dogs

DISCUSSION

Cysts and neoplasms are the two most common pathological conditions of the ovaries (Marković et al., 2003). Cysts produce estrogen, which is deposited in the body and acts as a carcinogenic hormone (Klein, 1996). The target organ for the development of tumors primarily are mammary gland, liver, lung, pancreas and spleen. Tumors can be benign or malignant, but malignant carcinomas, sarcomas, and adenocarcinomas are most common. Primary tumors of the ovaries can be induced by primitive epithelium, stromal cells and surface (coelomic) epithelium (Diez et al., 1998). All stromal tumors of sex glands share a tendency to be hormonal and thus produce signs of hyperestrogenism, masculinisation or persistent anestrus. Physiologically, hemorrhage may occur during ovulation and diagnosing a hemorrhage can be very complicated. At places where haemorrhage has occurred, there may be a formation of adhesions due to the presence of vascular fluid. Adhesions are formed between the ovary (bursa ovarica) and omentum (Cruz-Arambulo et al, 2004).

When the adhesion occurs, there is often unilateral occurrence of ovarian tumors. Newborn ovarian tumors compromise omentum and binding to it continues its development. There is usually a spread of micrometastases after cyst rupture within the neoplasm or invasion of the tumor through the ovary capsule (Meuten, 2017). Consequently, ascites and abdominal distension can be caused by lymph blockade. Such processes often develop metastases because they are leaning against large vessels of omentum, which is highly vascularized and connected with the digestive tract. The liver is the most common site of metastatic spread, mainly through a portal system that drains most of the abdominal structures (Nyland et al, 1983).

In dogs, metastatic neoplasms of the liver are approximately three times more likely to occur compared to primary neoplasms. Variable tissue characteristics in primary and metastatic neoplastic processes, including tissue density, vascular pattern, necrosis, fluid, and

calcification, cause a variable ultrasonographic display (Klein, 1996). Focal hypoechoic lesions with hyperechoic center called target lesions, are usually metastases. Early studies show that in the finding of at least one target lesion in the liver or spleen, there is a positive correlation of 74% for malignancy in small animals (Cuccovillo et al., 2002). Other signs indicating malignancy are the size of the lesion (more than 3 cm) and the presence of peritoneal effusion. Additional characteristics for distinguishing malignant from benign liver neoplasms can be established using CT as well as tumor-markers.

The most common pathological conditions of the uterus are: cystic hyperplasia of the endometrium, mucometra, hematometra, hydrometra, pyometra, segmental pyometra, granulomas and tumors of the uterus (Voges et al., 1996). The cystic hyperplasia of endometrium has hormonal etiology and usually precedes pyometra. Intraluminally there can be sterile mucus which causes hydrometra or a mucometra, depending on the degree of hydration of the mucin. In bitches, cystic hyperplasia of endometrium occurs under the influence of progesterone, but its initiation occurs under the influence of estrogen from the ovary (Klein, 1996). Due to the effect of progesterone, endometrium is susceptible to infection which causes cystic hyperplasia of the endometrium develop into pyometra. Changes outside the genital tract that accompany pyometra are expressed in the form of expressed caexia, metabolic disorders and glomerulonephritis caused by immune complexes (Klein, 1996). These secondary changes are especially noticed in bitches and can lead to intoxication with fatal consequences.

Leiomyomas are the most common tumors of the bovine uterus. They appear as multiple neoplasms, not only in the uterus, but also in the cervix and vagina. They are often associated with other disorders, such as cystic hyperplasia of the endometrium, follicular cysts and tumors of the mammary glands. Estrogen probably plays a significant role in the development and

maintenance of this neoplasm in bitches. Bladder cancer, especially in the bitches older than 8 years, are often malignant and results in metastases to the mammary gland (Mattoon et al., 1995). Mammary gland cancers represent one of the most common tumors in female dogs, and most attention is devoted to the effect of steroid hormones on tumorigenesis. Receptors for estrogen, progesterone and prolactin have been detected on normal and tumor cells of the mammary gland. About 40 to 60% of benign and malignant mammary tumors are positive for these receptors. Early castration, up to 2 years of age, has a protective effect and reduces the likelihood of mammary gland tumor in female dogs.

Of the pathological conditions on the testicles, neoplasms are the most common ones. In addition, they are often diagnosed with: orchitis, epididymitis, granuloma, abscesses, hematoma, hydrocele, varicocele, cryptorchidism and scrotal hernia (Johnston et al., 1991a). Testicular tumors can originate from the germinal epithelium (seminoma and teratoma) or interstitial cells (Sertoli-Leydig cell tumors) Seminomas often occur in cryptorchide and show local invasiveness, while teratomas are benign and occur predominantly in young animals. Sertoli cells tumor is most commonly found in dogs, giving metastases in the internal iliac lymph node. Tumor cells can produce estrogen and lead to hyperestrogenism that is manifested by gynaecomastia and alopecia in male animals, and bone marrow aplasia develops in some cases (Hart et al, 2013). The picture of orchitis is very similar to neoplasms. Often there is extratesticular fluid and an increase in epididymis. Benign prostatic hyperplasia, prostatitis, abscesses, prostatic and paraprostatic cysts, as well as neoplasms, are the most common pathological conditions of this accessory sex glands (Johnston et al., 2000). Prostatic hyperplasia is often a disease of older dogs resulting from the action of androgenic and estrogenic hormones. Over time, tissue fibrosis develops, which results in compression of the

urethra and difficulty urinating. In the parenchyma, there may be cysts of varying numbers and sizes. Over time, tissue fibrosis develops, which results in compression of the urethra and difficulty in urinating. Cysts of different numbers and sizes may be present within the parenchyma. Solitary cysts can be large enough to lead to ureteral obstruction, and consequently to hydronephrosis (Lukač et al, 1994). Prostatic neoplasms are mostly malignant in nature and in large percentage metastasized to regional lymph nodes, bones, and lungs (Atalan et al., 1999). Unlike with benign hyperplasia, there is capsule destruction and spread into the surrounding tissue. The most common are adenocarcinomas and undifferentiated carcinomas.

Pathological changes in the liver parenchyma are divided into diffuse changes in the liver, hemangiomas and focal liver lesions (Penninck et al, 2013). Diffusive lesions can be inflammatory, metabolic and vascular. Focal liver lesions can be primary and benign tumors. Vascular tumors are the most common primary liver tumors in general. Capillary hemangiomas represent a hyperechoic liver lesion, well-limited from the environment without a hypoechoic halo usually seen in malignant tumors, although it may also be transient in nature, as well as appear in a new location of parenchyma (Cruz-Arambulo et al 2004). They are usually rounded. Sometimes they can be hypoechogenic with small, numerous internal echoes. Adenomas are round focal changes that can be hyper and hypoechoic. They are often multiples and then they can show the difference in the echostructure. Nodular hyperplasia is usually hyperechoic. These benign tumors are encapsulated and have good limited edges. They are often large and can be multiple. Some of the are isoechoic. Malignant liver tumors include: hepatocellular carcinomas, fibrolamellar hepatocellular carcinoma and cholangiocarcinoma (Marković et al., 2003).

Echogenicity of hepatocarcinoma depends on its

tissue structure, whether it is rich in blood vessels, fats or connective tissue. Significant amount of fat gives a hyperechoic look. When it comes to solitary tumor it is usually very large and lobatus. Tumor masses are fairly well-restricted from the environment, with a hypoechoic halo. Hepatocarcinoma has a tendency to invade veins. Hepatocarcinomas are well vascularized and can bleed spontaneously. Metastases in the liver are usually multinodular, numerous, but there are also solitary ones (Cuccovillo et al., 2002). In liver metastases, there are changes in liver contour and tissue abnormalities. In liver metastases, there are changes in liver contour and tissue abnormalities. In metastases, ultrasound can sometimes detect calcification at a time when it is not yet seen on X-rays. Such calcification is most commonly seen in mucinous adenocarcinoma of colon or in ovarian cancer. Necrosis may occur in the metastases. In this case we are talking about necrotic metastases. Necrosis can occur spontaneously when the size of the metastasis exceeds 6-7 cm. dilatation of the intrahepatic bile duct can follow metastases (Lukač et al, 1994). This is due to compression caused by metastases. Hepatic veins can also change the appearance due to metastases. Veins with metastases are altered, deformed or destroyed. There may also be an invasion of the portal vein and its branches. Experimental studies show that the degree of metastatic vascularization affects its appearance (Cuccovillo et al., 2002). In general, hypervascular lesions are echogenic, and hypovascular are anechoic. Most commonly, along with these changes splenomegaly and retroperitoneal lymphadenopathy are also seen. Cirrhosis develops as a result of intoxication, metabolic disorders, chronic congestion and some infectious agents. In the differentiation of cirrhosis from the metastasis nodus, one must pay attention to attenuation coefficient and contour of the liver. In cirrhosis, attenuation is increased, and contours become biconvex (Stowater et al, 1990). Liver size can be helpful.

In cirrhosis the liver gradually decreases, and in the metastatic changes the liver is rapidly increased. Increased attenuation is seen only in cirrhosis. Splenomegaly and dilated v.lienalis occur only in cirrhosis. Portal hypertension arises as a result of intra and extrahepatic portal vein obstruction due to cirrhosis, fibrosis, steatosis, primary and metastatic tumors (Lukač et al, 1994). The halt in portal circulation leads

to a primary enlargement of the liver, a change in the structure of the parenchyma, and the expansion of the portal vein and its branches. Splenomegaly, collateral circulation and ascites develop secondarily. Also, metabolic diseases of the liver are accompanied by changes in the liver size, and sometimes also in parenchymal echostructure.

CONCLUSIONS

Apart from primary changes in the genital tract in both males and females, ultrasound diagnosis can also visualize subsequent changes in parenchymatous organs. Changes in the liver as a consequence of the primary disease of the reproductive tract in females may be cirrhosis hepatitis, as well as focal changes which cause hepatomegaly and mechanically endanger pancreas. In males, prostate tumors cause urinary incontinence, the formation of fibrin deposits starting with calcification process in

polyps and then with formation of a honeycomb-like unit with residual urine due to dilation of the urinary bladder.

Thanks to ultrasound diagnostics, veterinary knowledge as well as the extent of drug use have expanded. The main challenge for an attending veterinarian is not to cure consequential diseases but to suspect a primary disease using ultrasound.

LITERATURE

1. Atalan G, Holt PE, Barr FJ (1999) Ultrasonographic estimation of prostate size in normal dogs and relationship to bodyweight and age, *J Small Anim Pract*, 40:119–122.
2. Barella G, Lodi M, Sabbadin LA, Faverezani S (2012) A new method for ultrasonographic measurement of kidney size in healthy dogs *J Ultrasound*, 15:186–191.
3. Barthez PY, Nyland TG, Feldman EC (1995) Ultrasonographic evaluation of the adrenal glands in dogs, *J Am Vet Med Assoc*, 207:1180–1183.
4. Cruz-Arambulo R, Wrigley R, Powers B (2004) Sonographic features of histiocytic neoplasms in the canine abdomen, *Vet Radiol Ultrasound*, 45:554–558.
5. Cuccovillo A, Lamb CR (2002) Cellular features of sonographic target lesions of the liver and spleen in 21 dogs, *Vet Radiol Ultrasound*, 43:275–278.
6. Diez-Bru N, Garcia-Real I, Martinez EM, Rollan E, Mayenco A, Llorens P (1998) Ultrasonographic appearance of ovarian tumors in 10 dogs, *Vet Radiol Ultrasound*, 39:226–233.
7. Feeney DA, Johnston GR, Klausner JS, Bell FJ (1989) Canine prostatic ultrasonography, *Semin Vet Med Surg (Small Anim)*, 4:44–57.
8. Hecht S, Henry G (2007) Sonographic evaluation of the normal and abnormal pancreas, *Clin Tech Small Anim Pract*, 22:115–121.
9. Johnston GR, Feeney DA, Johnston SD, O'Brien TD (1991a) Ultrasonographic features of testicular neoplasia in dogs: 16 cases (1980–1988), *J Am Vet Med Assoc*, 198:1779–1784.
10. Johnston SD, Kamolpatana K, Root-Kustritz MV, Johnston GR (2000) Prostatic disorders in the dog, *Anim Reprod Sci*, 60–61:405–415.

11. Kantrowitz BM, Nyland TG, Feldman EC (1986) Adrenal ultrasonography in the dog, *Vet Radiol*, 27:91-96.
12. Klein MK (1996) Tumors of the female reproductive system. In: Withrow SJ, MacEwen EG, eds, *Small Animal Clinical Oncology*, Philadelphia: WB Saunders, pp 347–355
13. Lukač I, Kovačević N (1994) *Dijagnostički ultrazvuk u gastroenterologiji i nefrologiji*, Beograd, Dunaj.
14. Marković A (2003) Ultrazvuk u medicini, Udruženje za primenu ultrazvuka u medicini, Biologiji i veterini Srbije, 360-375,
15. Mattoon JS, Nyland TG (1995) Ovaries and uterus, In: Nyland4 TG,Mattoon JS, eds, *Small Animal Diagnostic Ultrasound*, Philadelphia: WB Saunders, pp 231–249.
16. Meuten DJ (2017) Tumors in domestic animals, *Collage of Veterinary Medicine North Carolina State University Raleigh, NC USA*, Fift edition,602-615, 689-702.
17. Nyland TG, Park RD (1983) Hepatic ultrasonography in the dog, *Vet Radiol*, 24:74-84.
18. Older RA, Van Moore A Jr, Glenn JF, Hidalgo HJ (1984) Diagnosis of adrenal disorders, *RCNA*, 22:433-454.
19. Penninck DG, Zeyen U. Taeymans O, Webster CR (2013) Ultrasonographic measurement of the pancreas and pancreatic duct in clinically normal dogs, *Am J Vet Res* 74:433–437.
20. Penninck DG, d' Anjou, MA (2015) *Atlas of Small animal ultrasonography*, Wiley Blackwell, Second Edition, 183-195, 311-314, 403-411.
21. Stowater JK, Lamb CR, Schelling SH (1990) Ultrasonographic feature of canine hepatic nodular hyperplasia, *Vet Radiol*, 31:268-272
22. Trigo, FJ, Thompson, H (1982) The pathology of liver tumours in the dog, *J Comp Pathol*, 92:21–39.
23. Voges AK, Neuwirth L (1996) Ultrasound diagnosis: Cystic uterine hyperplasia, *Vet Radiol Ultrasound*, 37:131–132.

Paper received: 28.12.2018.

Paper accepted: 07.02.2019.
