

The objectives  
and the rationale  
of Manuscript is  
clearly stated.

# WBC

WORLD BUIATRICS CONGRESS

SEPTEMBER 4<sup>TH</sup> TO 8<sup>TH</sup>

MADRID 2022

[www.wbc-madrid2022.com](http://www.wbc-madrid2022.com)

## ABSTRACT BOOK



# 31<sup>st</sup> World Buiatrics Congress

September 4<sup>th</sup> to 8<sup>th</sup>, 2022

Madrid, Spain

## Organizers:

**The National Association of Spanish Specialists in Bovine Medicine  
(ANEMBE)**



## The World Association for Buiatrics (WAB)



**WORLD  
ASSOCIATION  
FOR BUIATRICS**

**ASSOCIATION MONDIALE DE BUIATRIE**

**ASOCIACION MUNDIAL DE BUIATRIA**

**WELT-GESELLSCHAFT FÜR BUIATRIK**

**WBC 2022 Congress Technical Secretariat:**

**GRUPO PACIFICO**  
The power of meeting

Marià Cubí, 4, 08006,  
Barcelona, Spain.

Email: [wbcmadrid2022@pacifico-meetings.com](mailto:wbcmadrid2022@pacifico-meetings.com)



# INDEX

## VOLUME 2

POSTERS .....	7
Animal Health <b>(AH)</b> .....	9
Buffaloes and Camelids <b>(BC)</b> .....	43
Biotechnology <b>(BT)</b> .....	52
Cattle Welfare <b>(CW)</b> .....	55
Diagnostic Imaging <b>(DI)</b> .....	67
Epidemiology <b>(EP)</b> .....	76
Feedlot <b>(FE)</b> .....	83
Genetics and Breeding <b>(GB)</b> .....	89
Herd Health Management, Smart Farming and Economics <b>(HH)</b> .....	91
Infectious Diseases <b>(ID)</b> .....	110
Internal Medicine and Toxicology <b>(IM)</b> .....	142
Immunology and Vaccinology <b>(IV)</b> .....	152
Lameness <b>(LA)</b> .....	180
Nutrition and Metabolic Diseases <b>(NU)</b> .....	191
Parasitism <b>(PA)</b> .....	216
Public Health, Food Safety and Antimicrobial Resistance <b>(PH)</b> .....	228
Pharmacology and Therapeutics <b>(PT)</b> .....	245
Reproduction in Beef Cattle <b>(RB)</b> .....	248
Reproduction in Dairy Cattle <b>(RD)</b> .....	262
Sustainable Cattle Productive Systems <b>(SP)</b> .....	286
Small Ruminants <b>(SR)</b> .....	293
Surgery <b>(SU)</b> .....	318
Teaching and Continuing Education <b>(TE)</b> .....	324
Udder Health and Mastitis <b>(UH)</b> .....	327
INDEX OF AUTHORS .....	365

**AH-P45****How climate change affects dairy cow farming in Serbia**

Ivan Vujanac<sup>1</sup>, Radiša Prodanović<sup>1</sup>, Sveta Arsić<sup>1</sup>, Sreten Nedić<sup>1</sup>, Ljubomir Jovanović<sup>1</sup>, Milica Stojić<sup>1</sup>, Julijana Trifković<sup>2</sup>, Danijela Kirovski<sup>1</sup>.

<sup>1</sup>Faculty of Veterinary Medicine University of Belgrade, Belgrade, Serbia; <sup>2</sup>Faculty of Agriculture University of East Sarajevo, East Sarajevo, Bosnia And Herzegovina.

**Objective:** Due to climate change, for the period 2001 to 2030 temperature in Serbia is projected to rise by about 1°C, compared with 1961-1990. The aim of this study was to evaluate the effects of increased environmental temperature during summer season in Serbia, which is in the zone of moderate-continental climate, on milk production and milk composition in dairy cows, as well as postnatal adaptive capability of neonatal calves.

**Material and methods:** The trial was carried out on 40 Holstein Friesian cows and their neonatal calves (20 cows in summer and 20 cows in the spring season). Average daily feed consumption was measured for each cow. The cows were housed in a stable which during the summer was not equipped with additional cooling. The temperature-humidity index was recorded hourly. Based on the results calculated for average all-day THI, during the spring season the cows were not exposed to heat stress. During the summer season, the cows were exposed to the effects of moderate to severe heat stress. The cows' colostrum was sampled at 2, 14, and 26 h after calving, before feeding of their calves. Calves' blood samples were taken before the first colostrum intake and on days 1, 2, 3, and 7 of life. Calves' physiological parameters were measured on days 0 and 7. Milk production was measured daily from calving to day 90 of lactation.

**Results:** The quality of cows' colostrum was significantly reduced during summer season. The ingestion of the low-quality colostrum, combined with the thermal discomfort during summer season, provoked impaired physiological, biochemical, hormonal, and oxidative stress parameters in samples taken from the post-colostrum calves. Average daily milk yield in the spring season from day 60 to day 90 of lactation was significantly higher compared to the same lactation period during the summer season, while there was no significant difference in average milk yield from day 30 to day 60 of lactation between spring and summer season. Average daily feed consumption was significantly lower during the summer compared to spring during whole trial period.

**Conclusion:** Our results indicate that, during summer season, the negative impact of climate changes on milk production, milk composition and post neonatal adaptive capability of cows are strongly expressed. However, negative impact on milk production is not effective until day 60 of lactation, probably due to homeorhetic mechanisms which are important for the rearrangement of metabolism. This study highlights the importance of adequate supporting strategies for the care of the dairy cows and postnatal calves during the summer season.

This research was supported by the Science Fund of the Republic of Serbia.

Grant No 7750295, "Mitigation of methane production from

dairy cattle farm by nutritive modulation of cow's metabolism -MitiMetCattle."

**Keywords:** dairy cows, neonatal calves, climate changes.

**AH-P46****Effect of seasonality (spring × winter) on metabolism, health and immunity of dairy cows and their calves**

Fernanda Carolina Ramos Dos Santos<sup>1</sup>, Bianca Paola Santarosa<sup>2</sup>, Felipe Eduardo Dal Más<sup>3</sup>, Érica Cristina Bueno Do Prado Guirro<sup>3</sup>, Viviani Gomes<sup>1</sup>.

<sup>1</sup>School of Veterinary Medicine and Animal Science (FMVZ), University of São Paulo (USP), São Paulo, Brazil; <sup>2</sup>Institute of Agricultural Sciences (ICA), Federal University of Jequitinhonha and Mucuri Valleys (UFVJM), Unaí, Brazil; <sup>3</sup>Federal University of Paraná (UFPR), Palotina, Brazil.

**Objectives:** The objective of this study was to evaluate the effect of the season (spring and winter) on the metabolism, health, and immunity of neonate Holstein calves.

**Materials & Methods:** This research was in a dairy farm, in Descalvado, São Paulo, Brazil. It was evaluated 28 healthy multiparous Holstein cows, that calved female neonates, in eutocic delivery, evaluated during winter (July and August, n=15) and spring (September and October, n=13), which the average maximum, minimum, and thermal amplitude temperature were 28.2°C, 29.3°C, 9.2°C; 15.9°C, 21.2°C and 13.4°C, respectively. Dry cows were housed at indoor compost-barn, with cross-ventilation, 30 days before expected calving date. Immediately after birth, calves were separated in a "cuddle box" to maternal initial contact, and receive colostrum (≥21% BRIX) until 1h. From the 1<sup>st</sup> to the 14<sup>th</sup> day, calves stayed in closed-sided individual housing, hay bedding, covered barn, and the 15<sup>th</sup> to 28<sup>th</sup> in individual housing system outside the covered barn. At calving, the cow body condition score (BCS) was classified (1-5) and blood collected to analyse. Calves were blood sampled immediately after birth (D1) and on D2-3, D7, D14 and D28 days of life, 2h before the morning milk feeding, when has done a physical examination. Blood samples were collected in vacuum tubes containing fluoride sodium and with no anticoagulant, to obtain plasma and serum, respectively. Fecal and bovine respiratory disease (BRD) scores were assessed based in Calf Health Scoring Criteria (University of Wisconsin). With serum samples were analyzed cholesterol, triglycerides, iron, total protein and albumin, and with plasma, glucose, non-esterified fatty acids (NEFA) and β-hydroxybutyrate (BHB), through commercial kits in automated biochemical analyzer. Haptoglobin (Hp) was measured by spectrophotometric technique. The IgG levels in calf serum samples were quantified using an in-house sandwich ELISA. Measurement of endogenous and induced reactive oxygen species (ROS) production were conducted in whole blood cultures by using the fluorescent dye dihydrorhodamine 123. The cells were stimulated with inactivated antigens like *Staphylococcus aureus*; *Escherichia coli*; *S. hyicus* and Phorbol Myristate Acetate at 10<sup>-6</sup>M (PMA), and the capacity