

**Congress Proceedings**  
**XIII Middle European Buiatric's Congress**  
Best Western Hotel M, Belgrade, Serbia, June 5 - 8, 2013

Publisher: Serbian Buiatric's Association, Faculty of Veterinary  
Medicine, University of Belgrade  
Chief Editors: Prof. dr Dragan Gvozdić  
dr Branko Petrujkić  
Prepres and technical editor: BBN Congress Management d.o.o., 9 Deligradska, Belgrade  
Cover Design: Marija Marković  
Printing Office: Naučna KMD, Belgrade  
Circulation: 230

CIP - Каталогизација у публикацији  
Народна библиотека Србије, Београд

636.2/.3(082)  
614.9(082)  
637.1(082)

MIDDLE European Buiatric's Congress (13 ;  
2013 ; Beograd)

Congress Proceedings / XIII Middle  
European Buiatric's Congress, Belgrade,  
Serbia, June 5-8, 2013 ; organizers Serbian  
Buiatric's Association [and] Faculty of  
Veterinary Medicine [and] University of  
Belgrade ; chief editors Dragan Gvozdić,  
Branko Petrujkić. - Belgrade : Serbian  
Buiatric's Association : Faculty of  
Veterinary Medicine : University, 2013  
(Belgrade : Naučna KMD). - 596 str. : ilustr.  
; 24 cm

Tiraž 230. - Bibliografija uz svaki rad.

ISBN 978-86-916767-0-4 (SBA)  
1. Serbian Buiatric's Association (Beograd)  
a) Преживари - Зборници b) Ветерина -  
Зборници c) Млеко - Зборници  
COBISS.SR-ID 198690572

**COMMITTEES**

**National Organizing Committee**

President: Dragan Gvozdić  
Organizing secretary: Natalija Fratrić  
Technical secretary: Branko Petrujkić

**Members**

Vlado Teodorović  
Vera Katić  
Sanja Kovačević-Aleksić  
Milica Kovačević-Filipović  
Sonja Radojičić  
Slobodanka Vakanjac  
Horea Šamanc  
Miodrag Lazarević  
Ivan Jovanović  
Branislav Prokić  
Dragan Damjanović  
Dragan Šefer

**International Scientific Committee**

Prof. W. Baumgartner, PhD  
Prof. O. Szenci, PhD  
Ozbal Podpecan, PhD  
Borut Zemljic, PhD  
Prof. J. Kos, PhD  
Doc. Nikica Prvanovic, PhD  
Prof. K. Doll, PhD  
Prof. G. Kovác, PhD  
Doc. J. Illek, PhD  
J. Twardoń  
Prof. Natalija Fratric, PhD

**Conference Technical Organizer**

BBN Congress Management doo

## THE INFLUENCE OF DYLAIED SUBCLINICAL HYPOCALCEMIA IN COWS ON THE MACROELEMENTS LEVELS AND HEALTH STATUS OF NEWBORN CALVES

Lutnicki K, Kurek L, Kleczkowski M<sup>1</sup>, Banach A.

*Chair and Clinic of Internal Medicine, Faculty of Veterinary Medicine, University of Life Science in Lublin, Department of Internal Diseases of Farm Animals and Horses, Lublin, Głęboka 30, Poland. <sup>1</sup>Department of Pathology and Veterinary Diagnostics, Faculty of Veterinary Medicine, Warsaw University of Life Science, 159C Nowoursynowska, Warsaw, Poland.*

Corresponding author: krzysztof.lutnicki@up.lublin.pl

The most often noticed health disturbances in dairy cows herd in Poland are calcium (Ca), phosphorus (P), magnesium (Mg), copper (Cu) and zinc (Zn) deficiency. The mineral turnover impair exist as subclinical and clinical forms. Prolonged subclinical disturbances in mineral metabolism evoke the clinical symptoms in perinatal period, decreased milk production, and have the influence on fetus development. There are many papers and points of view concerning the maternal mineral deficiency and its influence on mineral levels in calves, especially in newborns. It is possible, that prolonged subclinical and clinical disturbances may affect on the health and mineral status in calves. The aim of this work was to asses the influence of prolonged subclinical hypocalcaemia in adult cows on calves macroelements levels, health status and body weight gain in first week after birth.

The investigation were carried out in 300 HF cows, 3-5 years old, mean milk yield 35 L per day. The experimental group consisted of 30 cows with abnormal Ca levels (hypocalcemic) in 2 and 5 months after calving and one month before next delivery. The control group cows exhibited no disturbances in the same experimental periods. The materials for analyses were obtained from newborn calves first 24 – 48 h and next 6-7 days after birth. In the same time the clinical status and welfare was assessed. The mean calcium content in experimental group was significantly lower in comparison to control, whereas the hypocalcemic cows exhibited no clinical signs.

The calves originated from this cows exhibited mean level of Ca -10.7 mg/dl; Mg- 2.9 mg/dl and Pi- 7.3 mg/dl after birth. The second proof after one week revealed 10.5; 2.8; 7.1 mg/dl respectively. Calves from control group demonstrated after birth: Ca concentration 10.6; Mg-2.6; P-6.4 mg/dl and after one week 10.4; 2.4; 6.3 mg/dl respectively. The Pi levels were significant higher in experimental group in both proofs. Calves in experimental group not exhibited worse weight gain as well as the susceptibility to infectious diseases. Obtained results do suggest, that prolonged subclinical hypocalcemia have no influence on clinical status, but evoke significant alterations in P levels. Moreover, the mean levels of above mentioned molecules are higher in calves obtained from hypocalcemic cows and its metabolism and hormonal regulation possibility is more efficient.

KEY WORDS: subclinical hypocalcaemia, macroelements, dairy cows, calves.

## PREPARTAL PLATELET COUNT AND FIBRINOGEN CONCENTRATION IN DAIRY COWS WITH AND WITHOUT RETAINED PLACENTA

Lužajić T\*, Katić M\*\*\*, Danilović Luković J\*\*, Blond B\*\*\*, Bogdanović D\*\*, Kovačević Filipović M.\*

\*Department of Pathophysiology, Faculty of Veterinary Medicine, University of Belgrade, \*\*Department of Biomedical Sciences, State University of Novi Pazar, \*\*\*PKB Corporation

Corresponding author: Milica Kovačević Filipović, Department of Pathophysiology, Faculty of Veterinary Medicine, University of Belgrade, Serbia, Bulevar Oslobođenja 18, e-mail: milkovac@yahoo.com

### ABSTRACT

Although extensive data are available on leukocyte changes in prepartal period in cows with and without retained placenta, there is no data describing changes in platelet count. The aim of this study was to define changes in platelet count in cows, with and without retained placenta, three weeks before and one day after calving. Using fibrinogen level as a general reference to cow's health, we also tested the hypothesis that cows, that develop placenta retention in prepartal period, have a subclinical systemic inflammatory reaction. Blood samples taken from 14 Holstein cows were analyzed, three times in a week interval before and 24 hours after calving. After calving, cows were retrospectively divided, in the group without (n=9) and the group with retained placenta (n=5). Clauss method was used for determination of fibrinogen concentration and blood smear examination for assessment of platelet count. The results have shown that retained placenta is associated with increase in platelet count seven days before calving and decrease in platelet count 24 hours after calving comparing to precalving values. The week before calving group of cows, that developed retained placenta had platelet count significantly higher, comparing to control group of cows. Fibrinogen level was not significantly different between groups. The recorded increase in platelet count, one week before calving can be related to unidentified stress factors, and decrease of platelet count 24 hours after calving, can be explained by their increased consumption, due to activation of haemostatic mechanisms after placenta abruption.

KEY WORDS: cattle, fibrinogen, platelets, prepartal period, stress

### INTRODUCTION

Retained placenta is a common problem in dairy cow farms and leads to major economic losses. Numerous studies have shown an association between retained placenta and twin pregnancy, cow's age, cow's body mass index and size of the calves (Laven and Peters, 1996; Gaafar et al., 2011). The prepartal risk factors that lead to disorders associated with retained placenta, also include some infective diseases and disorders of the immune system caused by deficiency of different nutritional factors (rev: Bealey et al., 2010). Unfortunately, in clinical conditions, individuals that will develop retained placenta cannot be recognized in advance. Our previous results showed that, total and differential leukocyte count, three weeks before calving and 24 hours after calving, were the same in cows with and without retained placenta (Lužajić et al., 2013, unpublished results). The only difference was a slight prepartal increase in segmented

rophil count in the group of cows, that developed retained placenta, comparing with cows without this disorder (Lužajić et al., 2013, unpublished results). However, in cattle, total leukocyte and differential counts are not good indicators of systemic reaction to tissue damage or stress (Nemi, 1993).

Deviation of platelet count is the most common hematological abnormality in veterinary medicine, yet it has never been assessed in dairy cows during peripartal period. In this study, we described changes in platelet count in cows, with and without retained placenta, three weeks before and 24 hours after calving. Using fibrinogen level, as a general reference to cow's health, we also tested the hypothesis that cows, that develop placenta retention in prepartal period, have a subclinical systemic inflammatory reaction.

MATERIALS AND METHODS

The study was performed on 14 Holstein dairy cows during the summer period. The animals were kept outdoors until the calving. They were 3 to 9 years old, with a uniform average annual milk yield. All animals were clinically examined and had no signs of disease a month before parturition. After calving, retained placenta was diagnosed in five cows (experimental group) and nine cows had puerperium without visible problems (control group). Blood was sampled three times before parturition, in a week interval, before the expected date of calving, and once again 24 hours after calving. To determine the concentration of fibrinogen, blood samples were obtained by puncture of *v.coccigea* and collected in tubes containing 3.2% Na-citrate (Vacuette, Greiner Bio-One). Blood smears were made immediately after sampling and stained with Hemacolor® (Merck). Fibrinogen concentration was determined by Clauss method (Clauss, 1957) and the platelet count was estimated from blood smear (Harvey, 2001). Statistical analysis was conducted using Student's t-test and analysis of variance for repeated measures (Repeated Measures ANOVA) with application of the Greenhouse-Geisser test.

RESULTS

Control group of cows had no variation in platelet count before and after calving (Table 1, Figure 1A). In the experimental group, the platelet count increased during prepartal period reaching a nadir one week before calving and significantly decreased after calving (Table 1, Figure 1A). Change in the platelet count in time, within the cows in the experimental group, was statistically significant (Table 2). Also, change between groups occurred in significantly different manner (Table 2). Platelet count was significantly lower in experimental group of cows three weeks and significantly higher one week before calving (Table 1).

Table 1. The platelet count and fibrinogen levels in cows from the control (n = 14) and experimental group (n = 5)

Variable and group	21 days before calving	14 days before calving	7 days before calving	24 hours after calving
<b>Platelets (x10<sup>9</sup>/L)</b>				
Control group	452.89±79.39	440.89±135.60	448.00±69.66	431.56±97.91
Experimental group	288.00±82.66*	391.20±126.31	572.80±131.74*†	365.60±145.54
<b>Fibrinogen (g/L)</b>				
Control group	3.43±1.07	3.93±0.88	3.28±0.57‡	4.87±1.63
Experimental group	3.24±0.56	4.60±1.17	3.82±1.89	4.54±2.54

NOTES: All values are given as mean ± standard deviation;  
 \* - p<0.01 in comparison with control group.  
 † p<0.01 in comparison with mesurment after calving




Figure 1. A and B. Changes in the platelet count and fibrinogen level before and after calving

Fibrinogen levels, during prepartal period, were similar in both groups of cows, yet the values were slightly higher in the experimental group (Table 1, Figure 1B). Significant increase in fibrinogen level has been observed in the control group of cows one day after calving, in comparison with levels one week before calving (Table 1). Analysis of variance for repeated measures, confirmed that in the control group, these changes were statistically significant (Table 2).

Table 2. Assessment of the impact of time and group on the investigated characteristics

Variable and type of effect	F	P	Partial Eta Squared
<b>Platelets</b>			
Within subjects effects of time in control group	0.075	0.922	0.009
Within subjects effects of time in experimental group	4.336	0.047	0.520
Within subjects effects of time in all cows	3.808	0.036	0.241
Within subjects effects of interaction time*group	3.751	0.038	0.238
Between subjects effects of group	2.109	0.172	0.150
<b>Fibrinogen</b>			
Within subjects effects of time in control group	4.473	0.023	0.359
Within subjects effects of time in experimental group	0.758	0.433	0.174
Within subjects effects of time in all cows	3.411	0.046	0.221
Within subjects effects of interaction time*group	0.540	0.601	0.043
Between subjects effects of group	0.144	0.711	0.012

NOTE: \* - p<0.05

DISCUSSION

Our results evidenced that retained placenta cows had an increase in platelet count a week before calving. It was previously shown that pro-inflammatory cytokines stimulate increase in platelet count in humans (rev: Klinger and Jelkmann, 2002) and some animal species (Sellon et al., 1997, Hammer, 1991; Neel et al., 2012). Corticosteroids also enhance platelet count (Mandell, 2000). In our experimental group, prepartal increase in platelet count, as well as slight increase in segmented neutrophils, (Lužajić et al., 2013, unpublished results) can be related to unidentified stress factors. Response to stress includes hypothalamic - pituitary -

cortex axis activation and leads to rise in glucocorticosteroids and possible enhancement of thrombopoiesis. Simultaneously, glucocorticosteroids prevent the transendothelial migration of neutrophils (Vlahos et al., 2012; Mandell, 2000). As the prepartal fibrinogen concentration was in reference range and was not significantly different between two groups before calving, we can conclude that subclinical inflammation was not present in that period. However, inclusion of more sensitive biomarkers (Eckersall and Bell, 2010) could be more reliable method for subclinical inflammation diagnostics.

Our results also showed that 24 hours after calving, platelet count decreased in a significant manner, when animals develop retained placenta. These data suggest that, platelets are activated and rapidly consumed in cows with retained placenta. In addition, 24 hours after birth, in the control group, fibrinogen concentration increased significantly, and was equal to the concentration of fibrinogen in the group with retained placenta. These data, as well as, data showing significant increase in the number of band neutrophils 24 hours after calving, (Lužajčić et al., 2013, unpublished results) as well as, increase in globulin concentrations, and a decrease in the total number of leukocytes, albumin and iron, (Katić, 2011) suggests that, tissue damage during calving, induce a systemic inflammatory response in all cows.

Based on all previous results, we consider, that the recorded increase in platelet count, one week before calving, in cows that develop retained placenta, can be related to unidentified stress factors, and decrease of platelet count 24 hours after calving, can be explained by their increased consumption, due to activation of haemostatic mechanisms after placenta abruption.

#### ACKNOWLEDGEMENTS

This work was supported by the Ministry of Science, Republic of Serbia (grant No. 175061)

#### REFERENCES

- Bealey JC, Whitman KJ, Baptiste KE, Scherzer J, 2010, Physiology and Treatment of Retained Fetal Membranes in Cattle, *J Vet Intern Med*, 24, 261-8.
- Clauss A, 1957, Fast coagulation method for estimation of fibrinogen, *Acta Haematologica*, 17, 237-46.
- Eckersall PD, Bell R, 2010, Acute phase proteins: Biomarkers of infection and inflammation in veterinary medicine, *Vet J*, 185(1), 23-7.
- Gaafar HM, Shamiyah ShM, El-Hamd MA, Shitta AA, El-Din MA, 2011, Dystocia in Friesian cows and its effects on postpartum reproductive performance and milk production, *Trop Anim Health Prod*, 43(1), 229-34.
- Hammer AS, 1991, Thrombocytosis in dogs and cats: a retrospective study. *Comp Haematol Int*, 1, 181-6.
- Harvey JW, 2012, Veterinary Hematology, a Diagnostic Guide and Color Atlas, *Saunders, Elsevier Inc*, 10-1.
- Katić M, 2011, Parameters of iron status in newborn calves and cows of different ages in the periparturient period, *Specialist thesis*, 60-5.
- Klinger MH, Jelknam W, 2002, Role of blood platelets in infection and inflammation, *J Interferon Cytokine Res*, 22(9), 913-22.
- Mandell PC, 2000, Essentials thrombocythemia and reactive thrombocytosis, In: Feldman FB, Zinkl GJ, Nemi CJ, editors, *Veterinary hematology*, 503-4.
- Neel JA, Snyder L, Grindem CB, 2012, Thrombocytosis: a retrospective study of 165 dogs, *Vet Clin Pathol*, 41(2), 216-22.
- Nemi C Jain, 1993, Essentials of veterinary hematology, 298-9.
- Peters AR, Laven RA, 1996, Treatment of bovine retained placenta and its effects, *Vet Rec*, 139(22), 535-9.

- Sellon DC, Levin JF, Palmer K, Millikin E, Grindem C, Covington P, 1997, Thrombocytosis in 24 horses (1989-1994), *J Vet Intern Med*, 11, 24-9.
- Vlahos R, Wark PA, Anderson GP, Bozonovski S, 2012, Glucocorticosteroids differentially regulate MMP-9 and neutrophil elastase in COPD, *PLoS One*, 7(3), 11-3.