

UDK 619 (05)

ISSN 1840-2887 Штампа-Print

ISSN 2303-4475 Online

ВЕТЕРИНАРСКИ ЖУРНАЛ РЕПУБЛИКЕ СРПСКЕ



Veterinary Journal of Republic of Srpska

Волумен/Volumen XXII, бр./No 1-2, стр./page 01-209, Бања Лука/Ванја Лука, 2022



**„Једно здравље“ за бољу будућност!
„One Health“ for a better future!**

UDK 619 (05)

ISSN 1840-2887 – Штампа-Print

ISSN 2303-4475 - Online

ВЕТЕРИНАРСКИ ЖУРНАЛ РЕПУБЛИКЕ СРПСКЕ

VETERINARY JOURNAL OF REPUBLIC OF SRPSKA

Научни часопис - Scientific journal

Ветеринарски журнал Републике Српске, Вол. 22, број 1-2, стр. 1-209, Бања Лука, 2022
Veterinary Journal of Republic of Srpska, Vol. XXII, No 1-2, page 1-209, Banja Luka, 2022

ИЗДАВАЧ – PUBLISHER:

ЈУ ВЕТЕРИНАРСКИ ИНСТИТУТ РЕПУБЛИКЕ СРПСКЕ „Др Васо Бутозан“ БАЊА ЛУКА
PI VETERINARY INSTITUTE REPUBLIC OF SRPSKA „Dr. Vaso Butozan“ BANJA LUKA

СУИЗДАВАЧ – CO-PUBLISHER:

ФАКУЛТЕТ ВЕТЕРИНАРСКЕ МЕДИЦИНЕ, УНИВЕРЗИТЕТ У БЕОГРАДУ, СРБИЈА
FACULTY OF VETERINARY MEDICINE, UNIVERSITY OF BELGRADE, SERBIA

ГЛАВНИ УРЕДНИК – EDITOR IN CHIEF:

Проф.др Драго Н. Недић, Prof.dr. Drago N. Nedic (БиХ-РС, В&Н-RS)

МЕЂУНАРОДНИ УРЕЂИВАЧКИ ОДБОР – INTERNATIONAL EDITORIAL BOARD:

Др Милорад Мириловић, Dr Milorad Mirilović, PhD, (Србија-Serbia)
Др Жељко Цветнић, Dr Željko Cvetnić, PhD, (Хрватска-Croatia)
Др Андреј Кирбиш, Dr Andrej Kirbiš, PhD (Словенија-Slovenia)
Др Нихад Фејзић, Dr Nihad Fejzić, PhD, (БиХ, В&Н)
Др Лазо Пендовски, Dr Lazo Pendovski, PhD, (Северна Македонија, North Macedonia)
Др Жељко Сладојевић, Dr Željko Sladojević, PhD, (БиХ-РС, В&Н-RS)
Др Јанко Иванов, Dr Yanko Ivanov, PhD (Бугарска, Bulgaria)
Др Предраг Слијепчевић, Dr Predrag Slijepčević, PhD (Велика Британија-United Kingdom)
Др Нектариос Д. Гиадинис, Dr Nektarios D. Giadinis (Грчка, Greece)
Др Мајкл Гилсдорф, Dr Michael Gilsdorf, PhD (САД-United State of America)
Др Семир Лончаревић, Dr Semir Lončarević, PhD (Шведска-Sverige)
Др Миломир Ковач, Dr Milomir Kovač, PhD, (Руска Федерација, Россијска Федерација)
Др Неђељко Карабасил, Dr Nedjeljko Karabasil, PhD, (Србија-Serbia)
Др Милан Балтић, Dr Milan Baltić, PhD, (Србија-Serbia)
Др Смиљана Параш, Dr Smiljana Paraš, PhD, (БиХ-РС, В&Н-RS)
Др Весна Калаба, Dr Vesna Kalaba, PhD, (БиХ-РС, В&Н-RS)
Др Драган Касагић, Dr Dragan Kasagić, PhD, (БиХ-РС, В&Н-RS)
Др Виолета Сантрач, Dr Violeta Santrač, PhD, (БиХ-РС, В&Н-RS)

ТЕХНИЧКИ УРЕДНИК: -TECHNICAL EDITOR: др Бојан Голић, dr Bojan Golić (БиХ-РС, В&Н-RS)

ЛЕКТОР - ЛЕКТОР: Маја Martinović

Преводилац на енглески/Translator in English: Проф. Нивес Јаковљевић, Prof. Nives Jakovljević

РЕЦЕНЗИЈА: Часопис се упућује на двоструку рецензију еминентним стручњацима. The journal is referred to a dual review by eminent experts.

ГОДИШЊЕ СЕ ОБЈАВЉУЈЕ 2 БРОЈА ЧАСОПИСА (јун/децембар или двоброј крајем године)

2 ISSUES OF MAGAZINES ARE PUBLISHED ANNUALLY (June / December or double issue at the end of the year)

Часопис је бесплатан и објављује се у електронској верзији и штампа у 300 примјерака. The magazine is free and is published in an electronic version and printed in 300 copies.

На основу Мишљења Министарства науке и технологије Републике Српске часопис је ослобођен пореза на промет. Based on the Opinion of the Ministry of Science and Technology of the Republic of Srpska, the magazine is exempt from sales tax.

Штампа: Штампариија Alfa studio Gradiška; **Press:** Alfa studio, Gradiška

78000 Бања Лука, Бранка Радичевића 18; 78000 Banja Luka, Branka Radičevića 18,

Тел: 051/229-210, Е-mail: drago.nedic@virs-vb.com,

<http://virs-vb.com/veterinarski-zurnal-rs/> <http://doisrpska.nub.rs/index.php/VJRS>

Обим и политика часописа

Ветеринарски журнал Републике Српске је покренут 2001. године како би истраживачи код нас, у региону и широм свијета могли да објављују своја истраживања и радове. Часопис је посвећен унапређењу и ширењу научних сазнања о ветеринарским наукама и сродним академским дисциплинама. Ветеринарски журнал Републике Српске је отвореног приступа, рецензирани научни часопис ЈУ Ветеринарски институт Републике Српске ”Др Васо Бутозан” Бања Лука, Република Српска/БиХ, посвећен објављивању оригиналних научних радова, прегледних научних радова, кратких саопштења и приказа случаја. Ветеринарски журнал Републике Српске објављује се на једном од језика у службеној употреби у Босни и Херцеговини и енглеском језику, у годишњим волуменима који се састоје од два броја, на моделу *Open Access* који омогућава високу видљивост радова. Ветеринарски журнал Републике Српске је индексиран као национални часопис. Сваки рукопис биће обрађен од стране два рецензента који пружају квалитетну уредничку обраду. Упутства за пријављивање рукописа, припрему рада, процедуру рецензије и све остале детаље доступна су на страницама часописа. Ова страница ће ауторе упутити корак по корак кроз процес пријављивање (Упутство за ауторе). Ветеринарски журнал Републике Српске је часопис отвореног приступа. Сви радови могу се бесплатно преузети и користити у складу са *Creative Commons Attribution 4.0 International License (CC BY 4.0)*.

Scope and Policy of the Journal

Veterinary Journal of Republic of Srpska was launched in 2001 to offer researchers and professionals in the region and worldwide a place to publish their research findings and work. The Journal is devoted to the advancement and dissemination of scientific knowledge concerning veterinary sciences and related academic disciplines. *Veterinary Journal of Republic of Srpska* is an open access, peer-reviewed scientific journal published by the PI Veterinary Institute Republic of Srpska „Dr. Vaso Butozan“ Banja Luka, Republic of Srpska/B&H, and dedicated to the publication of original scientific paper, review scientific paper, short communications, case report. *Veterinary Journal of Republic of Srpska* is published in one of the languages in official use in Bosnia and Herzegovina and English, in yearly volumes comprising two issues, on an Open Access model which allows high visibility of articles. *Veterinary Journal of Republic of Srpska* is indexed as a national journal. Each manuscript will be handled by two reviewers providing quality editorial processing. The instructions for manuscript submission, paper preparing, peer-review procedure and all other details are available on the Journal homepage. This site will guide Authors stepwise through the submission process (Author Guidelines).

Veterinary Journal of Republic of Srpska is an Open Access Journal. All articles can be downloaded free of charge and used in accordance with the Creative Commons Attribution 4.0 International License (CC BY 4.0).

САДРЖАЈ / CONTENTS

| | Стр./pp. |
|---|----------|
| QUORUM SENSING У МИКРОБИОЛОГИЈИ ХРАНЕ <i>Снежана Булајић, Тијана Ледина, Јасна Ђорђевић</i> | 5-17 |
| QUORUM SENSING IN FOOD MICROBIOLOGY <i>Snežana Bulajić, Tijana Ledina, Jasna Đorđević</i> | 18-30 |
| ДЕКЛАРИСАЊЕ МЕСА– МИШЉЕЊЕ И ПОВЕРЕЊЕ ПОТРОШАЧА <i>Јелена Јањић, Милан Ж. Балтић, Мирјана Ловреновић, Славен Грбић, Драго Недић, Споменка Ђурић, Бранислав Вејновић, Лазо Пендовски, Милорад Мириловић</i> | 31-43 |
| MEAT DECLARATION – CONSUMER OPINION AND CONFIDENCE <i>Jelena Janjić, Milan Ž. Baltić, Mirjana Lovrenović, Slaven Grbić, Drago Nedić, Spomenka Đurić, Branislav Vejnović, Lazo Pendovski, Milorad Mirilović</i> | 44-56 |
| МИКРОБИОЛОШКА ИСПРАВНОСТ I ФИЗИКАЛНО-КЕМИЈСКИ ПАРАМЕТРИ КВАЛИТЕТА КОЗЈИХ I ОВЉИХ СИРЕВА ИЗ ХЕРЦЕГОВИНЕ <i>Ана Сесар, Маја Дрмаћ, Виктор Ландека, Бојан Голић</i> | 57-67 |
| MICROBIOLOGICAL SAFETY AND PHYSICOCHEMICAL QUALITY PARAMETERS OF GOAT AND SHEEP CHEESE FROM HERZEGOVINA <i>Ana Sesar, Maja Drmać, Viktor Landeka, Bojan Golić</i> | 58-79 |
| QUORUM ANALIZA KVALITETA SEKSIRANOG SEMENA BIKOVA КОМПЈУТЕРСКИ АСИСТИРАНОМ АНАЛИЗОМ SEMENA-CASA <i>Слободанка Ваканјак, Горан Пантић, Љубомир Јовановић, Владимир Магаš, Данијела Кировски</i> | 80-89 |
| ANALYSIS OF THE QUALITY OF SEXED BULL SEMEN BY COMPUTER-ASSISTED SEMEN ANALYSIS-CASA <i>Slobodanka Vakanjac, Goran Pantić, Ljubomir Jovanović, Vladimir Magaš, Danijela Kirovski</i> | 90-100 |
| АНТИХЕЛМИНТИЧКИ ПОТЕНЦИЈАЛ ЕТАРСКОГ УЛЈА КОРИЈАНДЕРА (CORIANDRUM SATIVUM L.) КОД ОВАЦА <i>Филип Штрбац, Антонио Боско, Коста Петровић, Драгича Стојановић, Радомир Ратајак, Наташа Симић, Дејан Орчић, Ђиuseppe Cringoli, Лаура Риналди</i> | 101-113 |
| АНТИХЕЛМИНТИЧКИ ПОТЕНЦИЈАЛ КОРИЈАНДЕРА (CORIANDRUM SATIVUM L.) У ОВЦАМА <i>Filip Štrbac, Antonio Bosco, Kosta Petrović, Dragica Stojanović, Radimir Ratajac, Nataša Simin, Dejan Orčić, Giuseppe Cringoli, Laura Rinaldi</i> | 114-126 |
| КРИПТОРХИЗАМ КОД ПАСА <i>Јован Спасојевић, Иван Галић, Бојан Тохолј, Иван Станчић, Сандра Никوليћ, Ивана Давидов, Тијана Кукурић, Ненад Поповић</i> | 127-141 |
| CRYPTORCHIDISM IN DOGS <i>Jovan Spasojević, Ivan Galić, Bojan Toholj, Ivan Stančić, Sandra Nikolić, Ivana Davidov, Tijana Kukurić, Nenad Popović</i> | 142-155 |

| | Стр./pp. |
|---|--------------------|
| ЗНАЧАЈ САРАДЊЕ ЛЕКАРА И ВЕТЕРИНАРА У КОНЦЕПТУ ЈЕДНО ЗДРАВЉЕ <i>Бранислава Белић, Марко Цинцковић, Николина Новаков, Марија Јефтић, Драгана Димитријевић, Бобан Ђурић</i> | 156-166 |
| IMPORTANCE OF COLLABORATION BETWEEN PHYSICIANS AND VETERINARIANS IN ONE HEALTH CONCEPT <i>Branislava Belić, Marko Cincović, Nikolina Novakov, Marija Jeftić, Dragana Dimitrijević, Boban Đurić</i> | 167-177 |
| АКТИВНОСТ СЕЛЕНОЕНЗИМА GP _x -1 И GP _x -3 У КРВИ РАДНИХ КОЊА НА ТЕРИТОРИЈИ ЦЕНТРАЛНЕ СРБИЈЕ <i>Оливера Валчић, Петар Милосављевић, Иван Јовановић, Светлана Милановић</i> | 178-185 |
| ACTIVITY OF SELENENZYMES GP _x -1 AND GP _x -3 IN THE BLOOD OF WORKING HORSES IN THE TERRITORY OF CENTRAL SERBIA <i>Olivera Valčić, Petar Milosavljević, Ivan Jovanović, Svetlana Milanović</i> | 186-193 |
| УПУТСТВО АУТОРИМА И РЕЦЕНЗЕНТИМА INSTRUCTIONS FOR AUTHORS AND REVIEWERS | 194-201 202-209 |



DOI 10.7251/VETJEN2201018B

UDK 006.4:[663/664:658.562

Review scientific paper

QUORUM SENSING IN FOOD MICROBIOLOGY

Snežana BULAJIĆ, Tijana LEDINA, Jasna ĐORĐEVIĆ*

Faculty of Veterinary Medicine University of Belgrade, Belgrade, Serbia

*Corresponding author: Jasna Đorđević, jasna.djordjevic@vet.bg.ac.rs

Summary

In the last few decades, our perception of bacteria and their communities has changed significantly. In the past, the prevailing opinion was that bacterial communities represent a population of cells that act individually. Today, there is a well-documented fact that the mechanism of intercellular communication, known as "*quorum sensing*", plays an important role in controlling various cellular processes of bacteria such as bioluminescence, virulence, tolerance to disinfectants, sporulation, motility, biofilm formation and antimicrobial resistance.

Understanding the intercellular communication system gives the possibility to control the growth of undesirable bacteria in the food matrix and to develop a new strategy in ensuring the food quality and safety.

Keywords: quorum sensing, food quality and safety.

INTRODUCTION

„*Quorum sensing*“ (QS), term established by Fuqua and Winans (1994), represents a universal mechanism by which bacteria recognize changes in the environment. Recognizing newly established changes, the microbial world has the ability to react in a timely manner, by developing a specific defense strategy in order to adapt to unfavorable conditions in environment and during a time. In this way, by the act of socialization (association) at the moment when the population density reaches a critical level, bacteria synchronize their behavior and thereby optimize the chances of survival in competitive niches.

Numerous gram-negative and gram-positive species of bacteria are capable of adapting their activities to the social community and coordinating collective behavior within a multimicrobial community. The mechanism of combined expression involves the synthesis of low-molecular signal molecules. Bacteria ("emitter" cells) continuously generate a signal, at the beginning of growth in a

low concentration, and then, as the population density increases over time, the signal accumulates. Once, when the concentration of the signal reaches a critical level, which the bacteria perceive as a "quorum", the signal reacts with the receptor protein of the "responder" cell, and through the coordinated expression of certain genes, a unique, common response of the community is triggered in order to survive.

Processes controlled by the "*quorum sensing*" system include bioluminescence, sporulation, competence, production of antimicrobial peptides, biofilm formation, antimicrobial resistance, secretion of virulence factors, toxin production, transfer of conjugative plasmids, as well as production of enzymes whose activity is linked to food spoilage (Smith et al., 2004; Zhao et al., 2020; Wang and Xie, 2020).

Types of signaling molecules

Basically, the "*quorum sensing*" system involves the production and reading of extracellular signals. Signals are chemicals, signal molecules, so-called autoinducers. Often, although not always, the genes responsible for the synthesis of signaling molecules and the response to them activate their own expression, which explains the name autoinducer. According to Ammor et al. (2008), the identified signaling molecules are divided into 4 categories:

1. N-acyl homoserine lactones (AHL), derivatives of fatty acids, generically called autoinducers 1 (AI-1); they are produced and used by gram-negative bacteria mainly for intraspecies communication. The intracellular accumulation of a sufficient concentration of AHL triggers the transcriptional activation of various promoters within the bacterial genome. Several factors such as temperature, pH, NaCl content, growth medium composition and growth phase can affect the amount of AHLs produced. The production of C₄-homoserine lactone by *Aeromonas hydrophila* 519 depends on the growth temperature, glucose and salt concentration, as well as the pH value of the cultivation medium (Medina-Martinez et al., 2006). The same authors determine that at a basic pH value, AHLs become unstable and undergo hydrolysis. As bacteria enter the stationary phase, the amount of AHL decreases (Ravn et al., 2001). Examples of AHL-regulated phenotypes include antimicrobial peptide production, antimicrobial resistance, biofilm formation, DNA uptake competence status, cell differentiation, bioluminescence, growth, plasmid transfer, expression of virulence factors, as well as production of various extracellular enzymes.
-

2. furanosyl borate diester, known as autoinducer 2 (AI-2); they are created by gram-negative and gram-positive bacteria and thus serve as a universal signal for communication within and between species; responsible for the modulation of several phenotypic changes, such as expression of virulence factors, levels of ABC transporters in *Salmonella Typhimurium*, production of proteases, transcription of numerous genes in *Escherichia coli*, bioluminescence in *Vibrio* spp. Production of AI-2, as in the case of AI-1, is conditioned by the temperature and formulation of the growth medium, as well as changes in intracellular metabolism and the action of stressors.
 3. autoinducer 3 (AI-3) is a signal for virulence genes of enterohemorrhagic *Escherichia coli* (EHEC); the production of this autoinducer has also been found in the intestinal microbiota of humans, which implies a possible role in communication between species.
 4. autoinduced peptides (AIP) produced and used by gram-positive bacteria. AIPs are synthesized on ribosomes as precursor peptides, transformed into an active peptide and exported outside the cell by the action of ABC transporters in an ATP-dependent process. These peptides are characterized by low molecular weight (5-26 amino acid residues), high stability, specificity and diversity. The signal of these peptides is usually read on the cell surface via the histidine kinase sensor, when the autophosphorylation of the kinase triggers the phosphorylation of the regulatory protein, which ultimately leads to the transcription of the corresponding genes. Examples of phenotypes modulated by the activity of autoinduced peptides include genetic competence and sporulation in *Bacillus subtilis* and *Streptococcus pneumoniae*, virulence in *Enterococcus faecalis*, as well as production of antimicrobial compounds such as nisin (*Lactococcus lactis*) and subtilisin (*Bacillus subtilis*). Nevertheless, a typical example of an AIP-mediated QS mechanism is the *agr* system in *Staphylococcus aureus* (Abisado et al., 2018) and *Listeria monocytogenes*. *Staphylococcus aureus* represents the commensal microbiota of humans, but it also exhibits pathogenic properties after penetrating the host's tissue. The pathogenicity mechanism of staphylococci implies a biphasic strategy: at low population density, staphylococci express proteins that promote adherence to mucosal surfaces and colonization, while at high population concentration, these properties are extinguished, and the secretion of toxins and proteases is initiated, which promotes the dissemination of staphylococci. This "switch" in gene expression is regulated by the *agr* QS mechanism.
-

More recently, other types of signaling molecules have been described, such as cholera autoinducer 1, diketopiperazines, 4-hydroxy-2-alkylquinoline diffusible signaling factors (Wang and Xie, 2020), various metabolic products such as indole, short-chain fatty acids, secondary bile acids (Buffie et al., 2015).

Quorum sensing in the context of microbial ecology

In most cases, previous studies of the QS mechanism were related to the study of the molecular aspect of intercellular communication, that is, how QS affects virulence, sporulation or conjugation. Less attention was given to the ecological concept at the center of which lies the main question - why do bacteria produce signal molecules in the first place, guided by the basic ecological paradigm - everything is related to everything else, but also how the determinants of the ecological niche affect communication, especially in the case when the niche is represented by complex food ecosystems. Although the production of signaling molecules (AI-1, AI-2) is associated with certain members of the food microbiota (*Pseudomonas* species, members of the *Enterobacteriaceae* family, lactic acid bacteria), very little is known about the influence of process parameters and storage conditions on quality and quantity of these signals in food. It is known that the dominance of one group of bacteria in the continuum of food production and processing is the result of a complex process of microbial succession, when a specific population, thanks to its implicit characteristics and the adoption of a specific strategy, acquires numerical superiority in a niche that is established as an interplay of the physicochemical characteristics of the food matrix and storage conditions. In most cases, the food matrix implies the association of microbial cells with a solid substrate, and these cells are immobilized and localized in certain microenvironments in high concentration, and growth is present in the form of microcolonies or biofilms. At different places within the food matrix, there are variations in the level of oxygen, water activity, nutrients and pH values. Based on the interaction of all these factors, the food matrix is presented through a series of interconnected microenvironments, some of which can favor the growth of bacteria. The growth dynamics and activity of bacteria present in food, whether they are spoilage agents, pathogenic bacteria or beneficial microbiota, is determined, to a large extent, by *in situ* intercellular ecological interactions. For this reason, an ecological approach is necessary in understanding intercellular communication in different food ecosystems. Very likely, looking at the ecological dimension of the QS mechanism would provide answers to the following important questions:

- What is the critical concentration of the QS signal required for the bacteria to recognize the quorum and initiate gene expression?
-

- Is there a concentration gradient or chemical signal degradation due to dynamic (abiotic) conditions?
- Is the spatial distribution of cells more important than population densities in the generation and reading of QS signals?
- Is it possible that other species or strains of bacteria, which coexist in the same niche, deactivate and/or produce the same autoinducer?

Quorum sensing in food microbiology

Intercellular communication attracts the attention of food microbiologists due to the evidence of the connection between QS and food spoilage (Ammor et al., 2008; Skandamis et al., 2009; Galić et al., 2018) and the fact that different signaling molecules have been isolated from the matrix of spoiled food (Blana and Nychas, 2014), but also due to the importance of QS in bacterial pathogenesis (Li et al., 2018). In addition, the role of QS in the formation of biofilm in the process environment is undeniable (Galić et al., 2018).

Quorum sensing and food spoilage

Spoilage of food of animal origin is mainly associated with the activity of gram-negative proteolytic psychrotrophic bacteria, mostly *Pseudomonas* spp., and genera of the *Enterobacteriaceae* family, i.e. lactic acid bacteria if the food is stored in modified atmosphere conditions. In addition, the activity of hydrolases, mainly phospholipases originating from gram-positive aerobic *Bacillus* spp., is responsible for the spoilage of milk and milk products.

Pseudomonas spp. represent the dominant microbiota of aerobically packaged refrigerated meat, reaching a level of 10^9 CFU/g in case of meat spoilage (Liu et al., 2006). Jay et al. (2003) determined the connection between QS and meat spoilage caused by the activity of *Pseudomonas* spp., *Hafnia alvei* and *Serratia* spp., as dominant psychrotrophic species of the *Enterobacteriaceae* family isolated from spoiled meat packed in vacuum and identified as significant AHL producers. A large number of AHLs is determined in refrigerated beef and poultry meat stored in aerobic conditions, in a situation where a high concentration of *Pseudomonadaceae* (108-109 CFU/g) and *Enterobacteriaceae* (103-104 CFU/g) is recorded, with significant proteolytic activity (Liu and et al., 2006).

Yuan et al. (2018) assume the role of the QS mechanism during the growth and metabolic activity of *Pseudomonas fluorescens*, a frequent cause of spoilage of milk and milk products, due to the ability of the species to synthesize AHL and extracellular proteases. Most (84.5%) strains of psychrotrophic proteolytic microbiota isolated from raw milk are characterized by AHL production (Pinto et al., 2007). Although *Pseudomonadaceae* and *Enterobacteriaceae* cannot be

isolated from pasteurized and sterilized milk, the determined prevalence of AHLs indicates that AHLs produced by the initial microbiota of raw milk retain their activity completely, or at least partially, after thermal treatment. The production of alkaline metalloprotease in *Pseudomonas fluorescens* strain 395, which has a pronounced proteolytic activity, is regulated by the QS system (Liu et al., 2007). The production of extracellular lipolytic and proteolytic enzymes by *Serratia proteamaculans* strain B5a is controlled by the QS system (Christensen et al., 2003).

Quorum sensing and biofilm formation

Biofilm represents a complex ecosystem of one or more types of bacteria immersed in an extracellular polymer matrix. It develops according to patterns of multicellular behavior and in its final form functions as a cooperative consortium of bacteria in a complex but coordinated manner. This sessile way of life offers numerous advantages to associated members - greater resistance to stressful environmental conditions, insensitivity to antimicrobial agents as well as sanitation and disinfection agents, which makes it impossible to eradicate biofilm from the process environment. In addition, the microbial community of the biofilm shows a primitive homeostasis, a primitive circulatory system, the exchange of genetic material and the principle of metabolic cooperation.

The first step in biofilm formation involves conditioning the surface of the material and reversibly attaching cells to the same material. Then, binding becomes irreversible and microcolonies are formed. Finally, the film acquires a three-dimensional structure and the complex ecosystem is ready for dispersion. The milk industry, due to the specificity of the raw substrate (fluid) and the variety of technological processes (different temperature profiles), provides numerous opportunities for biofilm colonization. Tanks for raw milk, pipelines, centrifuges, duplicators for cheese, flow plate heat exchangers and packaging machines are examples of the so-called surface substrates on which biofilms are ideally formed. They are of particular importance since they represent a continuous, recurring source of contamination, which in the extreme case damages the quality and microbiological integrity of the final product.

The QS system is involved in all phases of biofilm formation: regulation of population density, synchronization of metabolic activity, so that the activity corresponds to the nutritional requirements, but also to the availability of the nutrient substrate. In addition, biofilm-resident bacteria show a significantly different genome/transcriptional program compared to free-living planktonic cells. The role of the QS system in the formation of biofilms was investigated in most cases in gram-negative bacteria, where acylated homoserine lactones (AHL) act as

signaling molecules. In a situation when the AHL concentration reaches a critical level, AHL binds to the receptor molecule and the formed complex, in response to stressful environmental conditions, triggers the expression and secretion of virulence factors, extracellular proteases, biofilm formation, as well as other physiological functions (Li et al., 2018).

In vitro studies show that *Salmonella* spp. readily adhere to work surfaces and form a biofilm (Coughlan et al., 2016). *Salmonella* spp. and *E. coli* do not have the *luxI* gene that encodes AHL synthetase and therefore do not have the ability to synthesize AHL. However, these microorganisms possess a LuxR homologue, SdiA, which detects a signal produced by other bacterial species. The presence of small concentrations of AHL produced by other bacteria induces biofilm formation in *E. coli*, *Salmonella enterica* serovar Typhimurium and *Vibrio* spp. (Jamuna Bai and Ravishankar Rai, 2016). Silagyi et al. (2009) determined that AI2 mediates the formation of *E. coli* biofilm, where the activated QS system also participates in the regulation of chemotaxis, flagella synthesis and expression of genes responsible for motility.

The presence of C12 HSL, one of the AHL signals, affects the formation of *Salmonella* Enteritidis biofilm and promotes the expression of virulence factors in anaerobic conditions (Almeida et al., 2017). Lamas et al. (2016) establish a link between biofilm formation by *S. enterica* and the expression of QS-related genes. In *Campylobacter jejuni*, AI-2 synthesis plays a major role in biofilm formation (Bezek et al., 2016). Biofilm formation in *Vibrio* spp. depends on several key enzymes, involved in the biosynthesis of flagella, pila and polysaccharides, as well as protein regulators that control the expression of the same genes (Giaouris et al., 2015). *Vibrio cholerae*, *V. vulnificus* and *V. parahaemolyticus* are present in contaminated food and water, especially seafood, where they form densely packed biofilms, which is a key factor in their survival in the environment (Galić et al., 2018). The QS system also controls the formation of biofilms in gram-positive foodborne pathogenic bacteria: *L. monocytogenes*, *Clostridium perfringens* and *B. cereus* (Coughlan et al., 2016). *L. monocytogenes* forms biofilms on abiotic surfaces in community with other bacteria, such as *Pseudomonas* spp. and *E. coli* (Giaouris et al., 2015). It was found that lactose induces biofilm formation in *B. cereus*, in a way that increases the production of AI-2 molecules (Duanis-Assaf et al., 2015). The Agr QS system in *S. aureus* differs from other QS systems since it has been proven that the expression of Agr leads to the down-regulation of genes responsible for bacterial adhesion. Thus, the expression of Agr reduces the adhesion potential of bacteria and indirectly affects the reduced capacity of biofilm formation.

Inhibition of QS system

As the QS system relies on the creation and transfer of small signaling molecules between bacteria, it is realistic to expect that the same molecules (AI) can be used to control the growth and survival of bacteria, as well as to modulate their virulence profile. Inhibition of the QS system, a mechanism known as "Quorum Quenching" (QQ), represents a new strategy in the control of specific bacterial phenotypes (bioluminescence, biofilm formation, virulence, swarming effect).

There are several possible ways of inhibiting the QS system:

1. Inhibition of AHL synthesis,
2. Degradation of AHL signal by enzyme activity (AHL-acylase and AHL-lactonase)
3. Interference with receptors or blocking of the AHL/receptor complex (Lade et al., 2014).
4. Post-transcriptional control of QS enzymes via sRNA.

Natural (isolated from various sources) and synthetic QS inhibitors are intensively studied. Extracts of different herbs show great potential. Methanolic extract of mango leaves reduces the production of protease, elastase, pyocyanin, as well as exopolysaccharides, and the swarming effect and biofilm formation in *Pseudomonas aeruginosa* (Husain et al., 2017). In the study of the same authors, more than 14 ingredients, including trifluoromethylketones and phenothiazines, were identified in the extract of mango leaves, with possible anti-QS action, namely by inhibiting "efflux" pumps, although it is very likely that they act on more than one target site.

Teas, especially polyphenols extracted from tea leaves, represent a traditional source of natural components, which have recently been intensively tested as possible anti-QS agents. Epigallocatechin gallate, the most abundant polyphenolic antioxidant metabolite from green tea, shows a strong inhibitory effect in relation to the expression of genes (biofilm formation, and motility) regulated by the QS system of *E. coli* (Lee et al., 2009). Spices such as black pepper, garlic, cumin and cinnamon show promising results as anti-biofilm agents.

Fruits, such as cranberries, strawberries and blackberries, contain, among other ingredients, phenols, quinones, flavonoids, alkaloids, terpenoids and polyacetylenes, which have proven to be successful as anti-QS agents, based on interference with AHL synthesis, inhibition of the production of AI- 2 (Bezek et al., 2016) or down-regulation of QS-related genes.

Quercetin (present in fruits, vegetables, nuts and cereals) shows a significant reduction of QS-dependent phenotypes of certain pathogenic food-borne bacteria (*P. aeruginosa*, *Y. enterocolitica* and *K. pneumoniae*), especially biofilm formation, exopolysaccharide production and motility (Gopu et al., 2015).

Vegetables and grains such as rice, tomatoes, soybeans, onions and broccoli also produce substances that mimic the activity of AHL signals (Jakobsen et al., 2012). Essential oils have proven to be good inhibitors of biofilm formation, primarily through the mechanism of AHL signal degradation or inhibition of AHL synthesis.

As alternative sources of QS system inhibitors, bioactive components originating from bacteria are mentioned, since it is quite certain that bacteria have adopted this mechanism in order to provide a competitive advantage in the niches in which they exist. In this sense, the members of soil microbiota are particularly interesting, which have been confirmed to produce enzymes (AHL-lactonases and AHL-acylases) capable of breaking down QS molecules, but also many so-called -ocins, such as bacteriocins, lactocins, enterocins and nisin (Choyam et al., 2019). Bee products are also recognized as a significant source of molecules with an anti-QS mechanism of action, especially propolis (Bulman et al., 2011) and honey (Maddocks and Jenkins, 2013).

One of the possible solutions in "turning off" the QS system is the application of paraoxonases, enzymes isolated from the serum of mammals, fungi and various plants. These enzymes block the QS system based on the hydrolysis of the lactone ring of N-acyl homoserine lactone (AHL) (Galić et al., 2018).

CONCLUSION

The QS system regulates the expression of genes responsible for coding many vital functions, which enable bacteria to survive in unfavorable environmental conditions. In an effort to bring the QS system under control, numerous antagonists of signaling molecules have been identified. It is important that food microbiologists are aware of the importance, but also understand the QS system, since strategies that "turn off" the quorum sensing mechanism in the bacterial population represent an effective tool for controlling the growth of undesirable bacteria in food.

Acknowledgment

The study was supported by the Ministry of Education, Science and Technological Development of the Republic of Serbia (Contract number 451-03-68/2022-14/200143).

Conflict of interest statement: The authors declare that there is no conflict of interest.

REFERENCES

- Abisado R. G., Benomar S., Klaus J. R., Dandekar A. A., Chandler J. R. (2018): Bacterial quorum sensing and microbial community interactions. *mBio*, 9(3):e02331-17.
- Almeida F. A., Pimentel-Filho N. J., Pinto U. M., Mantovani H. C., Oliveira L. L., Vanetti M. C. (2017): Acyl homoserine lactone-based quorum sensing stimulates biofilm formation by *Salmonella* Enteritidis in anaerobic conditions. *Archives of Microbiology*, 199(3):475-486.
- Ammor M. S., Michaelidis C., Nychas G. J. E. (2008): Insights into the Role of Quorum Sensing in Food Spoilage. *Journal of Food Protection*, 71(7):1510-1525.
- Bezek K., Kurincic M., Knauder E., Klancnik A., Raspor P., Bucar F., Smole Mozina S. (2016): Attenuation of adhesion, biofilm formation and quorum sensing of *Campylobacter jejuni* by *Euodia ruticarpa*. *Phytotherapy Research*, 30(9):1527-1532.
- Blana V. A., Nychas G. J. E. (2014): Presence of quorum sensing signal molecules in minced beef stored under various temperature and packaging conditions. *International Journal of Food Microbiology*, 173(3):1-8.
- Buffie C. G., Bucci V., Stein R. R., McKenney P. T., Ling L., Gobourne A., No D., Liu H., Kinnebrew M., Viale A., Littmann E., van den Brink M. R. M., Jenq R. R., Taur Y., Sander C., Cross J. R., Toussant N. C., Xavier J. B., Pamer E. G. (2015): Precision microbiome reconstitution restores bile acid mediated resistance to *Clostridium difficile*. *Nature*, 517(7533):205-8.
- Bulman Z., Le P., Hudson A. O., Savka M. A. (2011): A novel property of propolis (bee glue): Anti-pathogenic activity by inhibition of N-acyl-homoserine lactone mediated signaling in bacteria. *Journal of Ethnopharmacology*, 138(3):788-797.
- Choyam S., Srivastava A. K., Shin J. H., Kammara R. (2019): Ocins for food safety. *Frontiers in Microbiology*, 10:1736.
- Christensen A. B., Riedel K., Eberl L., Flodgaard L. R., Molin S., Gram L., Givskov M. (2003): Quorum-sensing-directed protein expression in *Serratia proteamaculans* B5a. *Microbiology*, 149(Pt 2):471-483.
- Coughlan L. M., Cotter P. D., Hill C., Alvarez-Ordóñez A. (2016): New weapons to fight old enemies: Novel strategies for the (bio)control of bacterial biofilms in the food industry. *Frontiers in Microbiology*, 7:1641.
- Duanis-Assaf D., Steinberg D., Chai Y., Shemesh M. (2015): The LuxS based quorum sensing governs lactose induced biofilm formation by *Bacillus subtilis*. *Frontiers in Microbiology*, 6:1517.
-

- Fuqua W. C., Winans S. C. (1994): A LuxR-LuxI type regulatory system activates *Agrobacterium* Ti plasmid conjugal transfer in the presence of a plant tumor metabolite. *Journal of Bacteriology*, 176:2796-2806.
- Galić S., García-Gutiérrez C., Miguélez E. M., Villar C. J., Lombó F. (2018): Biofilms in the food industry: Health aspects and control methods. *Frontiers in Microbiology*, 9:898.
- Giaouris E., Heir E., Desvaux M., Hebraud M., Moretro T., Langsrud S., Doulgeraki A., Nychas G. J., Kačaniova M., Czaczyk K., Olmez H., Simoes M. (2015): Intra- and inter-species interactions within biofilms of important foodborne bacterial pathogens. *Frontiers in Microbiology*, 6:841.
- Gopu V., Meena C. K., Shetty P. H. (2015): Quercetin influences quorum sensing in food borne bacteria: In-vitro and in-silico evidence. *PLoS One*, 10(8):e0134684.
- Husain F. M., Ahmad I., Al-Thubiani A. S., Abulreesh H. H., AlHazza I. M., Aqil F. (2017): Leaf extracts of *Mangifera indica* L. inhibit quorum sensing - regulated production of virulence factors and biofilm in test bacteria. *Frontiers in Microbiology*, 8:727.
- Jakobsen T. H., Bragason S. K., Phipps R. K., Christensen L. D., van Gennip M., Alhede M., Skindersoe M., Ostefeld Larse T., Hoiby N., Bjarnsholt T., Givskov M. (2012): Food as a source for quorum sensing inhibitors: Iberin from horseradish revealed as a quorum sensing inhibitor of *Pseudomonas aeruginosa*. *Applied and Environmental Microbiology*, 78(7):2410-2421.
- Jamuna Bai A., Ravishankar Rai V. (2016): Effect of small chain N acyl homoserine lactone quorum sensing signals on biofilms of food-borne pathogens. *Journal of Food Science and Technology*, 53(9):3609-3614.
- Jay J. M., Vilai J. P., Hughes M. E. (2003): Profile and activity of the bacterial biota of ground beef held from freshness to spoilage at 5–7 degrees C. *International Journal of Food Microbiology*, 81(2):105-11.
- Lade H., Paul D., Kweon J. H. (2014): Quorum quenching mediated approaches for control of membrane biofouling. *International Journal of Biological Sciences*, 10(5):550-565.
- Lamas A., Miranda J. M., Vazquez B., Cepeda A., Franco C. M. (2016): Biofilm formation, phenotypic production of cellulose and gene expression in *Salmonella enterica* decrease under anaerobic conditions. *International Journal of Food Microbiology*, 238:63-67.
- Lee K. M., Kim W. S., Lim J., Nam S., Youn M., Nam S. W., Kim Y., Kim S. H., Park W., Park S. (2009): Antipathogenic properties of green tea polyphenol epigallocatechin gallate at concentrations below the MIC against
-

- enterohemorrhagic *Escherichia coli* O157:H7. *Journal of Food Protection*, 72(2):325-331.
- Li T., Mei Y., He B., Sun X., Li J. (2018): Reducing quorum sensing mediated virulence factors expression and biofilm formation in *Hafnia alvei* by using the potential quorum sensing inhibitor l-carvone. *Frontiers in Microbiology*, 9:3324.
- Liu M., Gray J. M., Griffiths M. W. (2006): Occurrence of proteolytic activity and N-acyl-homoserine lactone signals in the spoilage of aerobically chill-stored proteinaceous raw foods. *Journal of Food Protection*, 69(11):2729-2737.
- Liu M., Wang H., Griffiths M. W. (2007): Regulation of alkaline metalloprotease promoter by N-acyl homoserine lactone quorum sensing in *Pseudomonas fluorescens*. *Journal of Applied Microbiology*, 103(6):2174-2184.
- Maddocks S. E., Jenkins R. E. (2013): Honey: A sweet solution to the growing problem of antimicrobial resistance? *Future Microbiology*, 8(11):1419-1429.
- Medina-Martinez M.S., Uyttendaele M., Demolder V., Debevere J. (2006): Influence of food system conditions on N-acyl-Lhomoserine lactones production by *Aeromonas* spp. *International Journal of Food Microbiology*, 112(3):244-252.
- Pinto U. M., de Souza Viana E., Martins M. L., Vanetti M. C. D. J. F. C. (2007): Detection of acylated homoserine lactones in gram-negative proteolytic psychro-trophic bacteria isolated from cooled raw milk. *Food Control*, 18(10):1322-1327.
- Ravn L., Christensen A. B., Molin S., Givskov M., Gram L. (2001): Methods for detecting acylated homoserine lactones produced by gram-negative bacteria and their application in studies of AHL-production kinetics. *Journal of Microbiology Methods*, 44(3):239-251.
- Silagyi K., Kim S. H., Lo Y. M., Wei C. I. (2009): Production of biofilm and quorum sensing by *Escherichia coli* O157:H7 and its transfer from contact surfaces to meat, poultry, ready-to-eat deli, and produce products. *Food Microbiology*, 26(5):514-519.
- Skandamis P. N., Stopforth J. D., Ashton L. V., Geornaras I., Kendall P. A., Sofos J. N. (2009): *Escherichia coli* O157:H7 survival, biofilm formation and acid tolerance under simulated slaughter plant moist and dry conditions. *Food Microbiology*, 26(1):112-19.
- Smith J. L., Fratamico P. M., Novak J. S. (2004): Quorum sensing: a primer for food microbiologists. *Journal of Food Protection*, 67(5):1053-70.
- Wang X. Y., Xie J. (2020): Quorum Sensing System-Regulated Proteins Affect the Spoilage Potential of Co-cultured *Acinetobacter johnsonii* and
-

Pseudomonas fluorescens From Spoiled Bigeye Tuna (*Thunnus obesus*) as Determined by Proteomic Analysis. *Frontiers in Microbiology*, 11:940.

Yuan L., Sadiq F. A., Burmolle M., Liu T., He G. (2018): Insights into bacterial milk spoilage with particular emphasis on the roles of heat-stable enzymes, biofilms, and quorum sensing. *Journal of Food Protection*, 81(10):1651-1660.

Zhao X., Yu Z., Ding T. (2020): Quorum-Sensing Regulation of Antimicrobial Resistance in Bacteria. *Microorganisms*, 8(3):425.

Paper received: 13.05.2022.

Paper accepted: 25.07.2022.
