

IN VIVO CYTOGENETIC ANALYSIS OF THE EFFECTS OF CLOPROSTENOL ON MAMMALIAN CELLS

MILOŠEVIĆ*, M. ANDELKOVIĆ**, B. SOLDATOVIĆ* and DIJANA CVETKOVIĆ***

*Department of Biology, Faculty of Veterinary Medicine, Bul JNA 18, Belgrade **Institute of Biology, Faculty of Biology, Studentski Trg 16, Belgrade ***Department of Anatomy, Faculty of Veterinary Medicine, Bul JNA 18, Belgrade

(Received, 3. March 1997.)

The aim of this study was to evaluate possible genotoxic effects of cloprostenol (Sinhrogal[®]), a synthetic analogue of prostaglandin F-2 alpha, which is used for synchronization of estrus in domestic animals. The mutagenic potential of the test substance was investigated on human peripheral blood lymphocytes (in vitro) and bone marrow cells of intramuscularly treated BALB/c AnNCR male mice (in vivo), by monitoring numerical and structural chromosome aberrations, as well as determination of the mitotic index. There were no indications for a mutagenic potential of cloprostenol within the experimental concentrations and doses applied in this investigation.

Key words: cloprostenol, prostaglandin F-2 alpha, genotoxicity, chromosome aberrations, mitotic index

INTRODUCTION

Endocrinological evidence and data from animal experiments strongly indicate the importance of hormones in development and/or maintenance of target cells (Henderson et al., 1982). Having in mind that most cancers originate as a consequence of genetic changes, possible mutagenic effects of hormones—especially steroids, have been studied in various test-systems. However, the influence of nonsteroid hormones and mediators requires deeper endocrinological characterization.

According to the available data, mediator PGF-2 alpha has genotoxic action (Bennet, 1983), may stimulate proliferation of cultured cells (Hakeda et al., 1981), cause chromatin decondensation of small and large luteal cells (Chegini et al., 1981) whereas metabolites of PGF-2 alpha can bind to DNA (Karmali et al., 1981). It is interesting that many mutagens and carcinogens stimulate PGF-2 synthetase in cells (Levine, 1977). Moreover, some tumor cells contain an elevated level of PGF-2 alpha (Bennet et al., 1981) and it is thought that changes in its levels can be the mechanism by which mutagens and carcinogens cause DNA damage.

The objective of our study was to investigate possible genotoxic properties of cloprostenol (Sinhrogal[®]) which is a synthetic analogue of PGF-2 alpha with a slower biotransformation time and a more profound effect. Cloprostenol is used in veterinary medicine since its luteinic effect synchronizes estrus and ovulation in domestic animals therefore allowing successful artificial insemination.

MATERIALS AND METHODS

Test substance. Cloprostenol (CAS No 40665-92-7) is a functional analogue of prostaglandin F-2 alpha (PGF-2 alpha), chemically (16-m-chlorophenyl)-18, 19, 20-tetranol PGF-2 alpha; molecular weight 424.92. Cloprostenol is used in veterinary medicine mainly for synchronization of estrus. The investigations were performed with Sinhrogal[®] (ICN Galenika) which contains cloprostenol as an active component.

Controls. The negative control was prepared in ICN Galenika as a placebo preparation containing all compounds present in Sinhrogal[®] except the active one. The positive control N-methyl-N'-nitro-N-nitrosoguanidine (MNNG) was dissolved in acetone and added to cultures to obtain a final concentration of 50 µg/ml whereas animals were treated intramuscularly with a dose of 50 mg/kg.

In vitro test. Human peripheral blood lymphocyte cultures were set up according to the slightly modified protocol described by Evans and O'Riordan (1975). Heparinised whole blood samples (0.8 ml) obtained from healthy men under 30 years of age were added to vials with 9.2 ml of Parker 199 medium containing 30% of inactivated calf serum (SERVA) and 0.04 mg/ml of phytohaemagglutinin (Wellcome). Cultures were incubated for 72 hours at 37°C. Exactly 47 h and 30 min after the beginning of incubation, cloprostenol (Sinhrogal[®], ICN Galenika) was added to the cultivation vials in such amounts to obtain final experimental concentrations of 1, 2, 5 and 10 µg/ml. Two hours before harvesting, colcemide (CIBA) was added to the cultures to achieve a final concentration of 0.5 µg/ml. After hypotonic treatment (0.075 M KCl) followed by three repetitive cycles of fixation in methanol-acetic acid solution (3:1), centrifugation, and resuspension, the cell suspension was dropped on clean, grease free microscopic slides, air dried and stained in 20% Giemsa (Kanto) for 15 minutes. For each of the experimental concentrations, as well as for controls, 150 mitotic plates were examined on coded slides.

In vivo test. The BALB/c AnNCR inbred strain of mouse was used for the assay. Animals were kept under standard conditions: $(21 \pm 1)^{\circ}\text{C}$, humidity 60%, food and water *ad libitum*. During the course of the experiment their body weight was 20-25 g. For each of the experimental concentrations (1, 3 and 10 mg/kg), as well as for controls, 10 male animals were treated intramuscularly with Sinhrogal[®] and control preparation respectively. At 24 hours after treatment, animals were injected intraperitoneally with colchicine (4 mg/kg). Chromosome preparations were made according to the method of Hsu and Patton (1966). Cytogenetic analysis implied observation of heteroploid and polyploid cells, structural chromosomal aberrations and mitotic activity.

was to investigate possible genotoxic properties of cloprosteno... which is a synthetic analogue of PGF-2 alpha... and a more profound effect. Cloprosteno... and a more profound effect. Cloprosteno... and a more profound effect. Cloprosteno...

MATERIALS AND METHODS

(CAS No 40665-92-7) is a functional analog... (16-m-chlorophen... F2 alpha), chemically... Cloprosteno... molecular weight 424,92. Cloprosteno... synchronization of estrus. The investig... (ICN Galenika) which contains cloprosteno...

was prepared in ICN Galenika as a... present in Sinhrogal[®] except the... nitro-N-nitrosoguanidine (MNNG) was... to obtain a final concentration of... muscularly with a dose of 50 mg/kg

l blood lymphocyte cultures were... protocol described by Evans and O'Brien... samples (0.8 ml) obtained from healthy... to vials with 9.2 ml of Parker 199... calf serum (SERVA) and 0.04 mg/ml... cultures were incubated for 72 hours... the beginning of incubation, cloprosteno... to the cultivation vials in such amounts... of 1, 2, 5 and 10 µg/ml. Two... was added to the cultures to achieve... tonic treatment (0.075 M KCl) followed... methanol-acetic acid solution (2:1)... the cell suspension was dropped on... and stained in 20% Giemsa... experimental concentrations, as well as... on coded slides.

inbred strain of mouse was used for... standard conditions: (21 ± 1)°C, humidity... the course of the experiment the... experimental concentrations (1, 2, 5, 10 µg/ml) and... animals were treated intramuscularly... respectively. At 24 hours after treatment... with colchicine (4 mg/kg). Chromosomes... to the method of Hsu and Patton (1960)... of heteroploid and polyploid... and mitotic activity.

Statistical analysis of the obtained results was performed with Student's

RESULTS AND DISCUSSION

Genotoxicological analysis of the effects of cloprosteno... (aneuploid and polyploid cells) and structural aberrations... and chromatid gaps and breaks) in PHA-activated human... lymphocytes and bone marrow of BALB/c AnNCR male mice. In... possible cytostatic or cytotoxic effects of cloprosteno... (the percentage of cells during mitosis) was established for each... concentration, negative and positive controls.

The results of the *in vitro* cytogenetic test are shown in Table 1. The level... aberrations for experimental concentrations of cloprosteno... 2.00% to 3.47% and those changes were not statistically significant in... to the value of the negative control (2.00%). Metaphase... isolated from cultures treated with 10 µg/ml of cloprosteno... Figure 1. Only treatment with the positive control (10⁻⁶ M MNNG) caused



Figure 1. Giemsa-stained human metaphase chromosomes isolated from a culture treated with 10 µg/ml of cloprosteno

... (p < 0.01) increase in the percentage of mitoses with gaps and... numerical aberrations, the number of polyploid and aneuploid cells... significantly changed either after administration of cloprosteno... or after... with positive control.

... to ascertain whether cloprosteno... can exhibit genotoxic effects in... preparations from the bone marrow of BALB/c inbred strain

were analyzed after single administration of cloprostamol (Table 2). The level of structural as well as numerical aberrations was similar to the values observed in cultured human peripheral blood lymphocytes obtained in this experiment. The positive control (MNNG) gave rise to a significant increase in aneuploid cells ($p < 0.01$) and cells with structural aberrations ($p < 0.001$) (an example is shown in Figure 2), whereas the level of polyploid cells did not depart markedly from the negative control. On the other hand, there were no significant changes in the levels of numerical and structural aberrations for all cloprostamol concentrations applied. A metaphase spread from an animal exposed to the highest dose of cloprostamol is presented in Figure 3.

Table 1. The effects of cloprostamol in lymphocyte cultures of human peripheral blood

Conc. of cloprostamol	Mitotic index (%)	Number of cells observed	Aneuploid cells		Polyploid cells		Cells with structural and numerical aberrations
			No	%	No	%	
negative control	5.60	150	7	4.67	2	1.33	3
positive control (MNNG)	3.83	150	11	7.33	5	3.33	22
1 $\mu\text{g/ml}$	5.98	144	6	4.17	1	0.69	5
2 $\mu\text{g/ml}$	5.34	153	4	2.61	2	1.31	3
5 $\mu\text{g/ml}$	6.35	152	9	5.92	2	1.97	4
10 $\mu\text{g/ml}$	6.15	151	6	3.97	1	0.66	3

** $p < 0.01$




Figure 2. Acentric fragments (indicated by arrows) as a consequence of the mutagenic action of MNNG in mouse cell

In this experiment mainly structural chromosomal aberrations like gaps and break damage of chromosomes, which indicate clastogenic properties could

...ation of cloprostenol (Table 2). The ...
 ... was similar to the values observed ...
 ... lymphocytes obtained in this experiment ...
 ... to a significant increase in aneuploid ...
 ... aberrations ($p < 0.001$) (an example is ...
 ... ploid cell did not depart markedly ...
 ... there were no significant changes ...
 ... aberrations for all cloprostenol concentrations ...
 ... an animal exposed to the highest dose ...


...cyte cultures of human peripheral blood

	Aneuploid cells		Polyploid cells		Cells with gaps and breaks
	No	%	No	%	
7	4.67	2	1.33	3	
11	7.33	5	3.33	22	
6	4.17	1	0.69	5	
4	2.61	2	1.31	3	
9	5.92	2	1.97	4	
6	3.97	1	0.66	3	



... as a consequence of the mutagenic ...
 ... chromosomal aberrations like gaps ...
 ... indicate clastogenic properties ...

... For the scoring of gaps and breaks we used criteria described by ... (1982).



... spread from mouse bone marrow after treatment with 5 mg/kg of cloprostenol

... The values of mitotic index (MI) are shown in Tables 1 and 2. Obviously, ...
 ... in the values of MI are not significant either for all experimental ...
 ... administered *in vitro*, or for all doses of cloprostenol injected into ...

... effects of cloprostenol on bone marrow cells of BALB/cAnNCR male mice

	Number of treated animals	Mitotic index (%)	Number of cells observed	Aneuploid cells		Polyploid cells		Cells with gaps and breaks	
				No	%	No	%	No	%
	10	2.05	500	21	4.20	5	1.00	13	2.60
	10	1.65	500	44	8.80**	7	1.40	77	15.40***
	10	1.95	500	20	4.00	6	1.20	10	2.00
	10	2.60	500	31	6.20	8	1.60	11	2.20
	10	2.55	500	29	5.80	6	1.20	9	1.80

... Therefore, on the basis of the results obtained by the cytogenetic analyses ...
 ... in this paper it can be concluded that cloprostenol has no expressed ...
 ... properties. It should be mentioned, however, that the same agent was ...
 ... as potentially genotoxic in SCE (sister chromatid exchange) test ...
 ... (1994). This discrepancy can be accounted for by the higher

sensitivity of the SCE test in comparison to the analysis of structural and numerical chromosomal aberrations (Perry, 1980). These findings indicate that cloprostenol might be designated as a relatively weak mutagen. In addition, it should be pointed out that cloprostenol is rapidly metabolized in treated animals. Perry et al. (1983) have shown that 13 hours after an intramuscular injection of cloprostenol its level in heifer plasma did not differ significantly from control values. Therefore, proper administration of cloprostenol in the breeding and reproduction of domestic animals implies that human exposure is almost negligible.

CONCLUSION

Cloprostenol (Sinhrogal^R), a synthetic functional analogue of PGF₂ was evaluated for possible mutagenic potential in PHA-activated human peripheral blood lymphocytes and bone marrow cells from BALB/c AnNCr mice. There were no indications for a mutagenic effect of cloprostenol at the concentrations and doses used in this experiment.

REFERENCES

1. Bennet A., Carroll M. A., Stamford I. F. et al. 1981. Prostaglandins and human lung cancer. *Pharmacol* 74, 207-212.
2. Brogger A. 1982. The chromatid gap - a useful parameter in genotoxicology? *Cytogenetics and Genetics* 33, 14-19.
3. Chegini N., Lei Z. M., Rao C. V. 1991. Nuclear volume and chromatin conformation of small and large bovine luteal cells: effect of gonadotropins and prostaglandins and dependence on estradiol phase. *Cell Tissue Res* 264, 453-460.
4. Das U. N., Ramadevi G., Rao K. P., Rao M. S. 1989. Prostaglandins can modify gamma-radiation and chemical induced cytotoxicity and genetic damage in vitro and in vivo *Prostaglandins* Vol. 38, No. 6, 689-716.
5. Djelić N., Soldatović B., Anđelković M. 1994. Effect of cloprostenol on the mitotic index and SCE frequency in cultures of human lymphocytes. *Acta Veterinaria*, Vol. 44, No. 5-6, 337-344.
6. Evans H. J., O'Riordan M. L. 1975. Human peripheral blood lymphocytes for the analysis of chromosome aberrations in mutagen tests. *Mutation Research* 31, 135-148.
7. Hakeda Y., Harada S., Matsumoto T., Tezuka K., Higashino K., Kodama H., Hashimoto-Sato Ogata E., Kumegawa M. 1991. Prostaglandin F-2 alpha stimulates proliferation of osteoblastic MC 3T3-E1 cells by up-regulation of insulin-like growth factor I receptors. *J. Biol. Chem.* 266 (31), 2144-2150.
8. Henderson B. E., Ross R. K., Pike M. C., Casagrande J. T. 1982. Endogenous hormones as a growth factor in human cancer. *Cancer Research* 42, 3232-3239.
9. Hsu T. C., Patton J. L. 1969. Human peripheral blood lymphocytes for the analysis of chromosome aberrations in mutagen tests. In "Handbook of mutagenicity test procedures" (Ed. Kilbey, L. S.) Elsevier/North Holland Biomedical Press, pp. 261-274.
10. Karmali R. A., Schiller P., Horrobin D. F. 1976. Prostaglandins can prevent binding of chromosomes to calf thymus DNA. *Prostaglandins* 12, 463-464.
11. Levine L. 1977. Chemical carcinogens stimulate canine kidney (MDCK) cells to produce prostaglandins. *Nature* 268, 447-448.
12. Perry P. E. 1980. Chemical mutagens and sister-chromatide exchange. From "Chemical Mutagens" Vol. 6, Frederic J. de Serres and Alexander Hollaender (Eds.) Plenum Publishing Corporation, pp. 1-39.

the analysis of structural and numerical changes in chromosomes. These findings indicate that cloprostenol is a weak mutagen. In addition, it should be noted that cloprostenol is metabolized in treated animals. Pictograms showed that after an intramuscular injection of cloprostenol, the results did not differ significantly from control. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

functional analogue of PGF₂ alpha. It is a potent mitogenic potential in PHA-activated human lymphocytes. Cloprostenol is a weak mutagen. In addition, it should be noted that cloprostenol is metabolized in treated animals. Pictograms showed that after an intramuscular injection of cloprostenol, the results did not differ significantly from control. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

prostaglandins and human lung cancer. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

in genotoxicology? Cytogenetic analysis of the effects of cloprostenol on the mitotic index and chromatin conformation of mammalian lymphocytes and dependence on the dose of cloprostenol.

prostaglandins can modify gamma-radiation induced DNA damage in vitro and in vivo. Prostaglandin synthase inhibition by cloprostenol on the mitotic index and chromatin conformation of mammalian lymphocytes.

cloprostenol on the mitotic index and chromatin conformation of mammalian lymphocytes. *Veterinaria*, Vol. 44, No. 5-6, 337-344. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

alpha stimulates proliferation of mammalian lymphocytes. *Research* 31, 135-148. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

Endogenous hormones and their effects on the mitotic index and chromatin conformation of mammalian lymphocytes for the analysis of chromosome aberrations. *Research* 31, 135-148. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

can prevent binding of chromosomes to the cell cycle. *Research* 31, 135-148. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

kidney (MDCK) cells to produce prostaglandins. *Research* 31, 135-148. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

exchange. From "Chemical Mutagenesis and Carcinogenesis" (Eds. K. Hayashi and T. Friedberg) Plenum Publishing Corporation, New York, NY, 1982, pp. 135-148. The results of the study of cloprostenol in the breeding of animals show that human exposure is almost negligible.

cloprostenol. *Veterinaria* (Beograd), Vol. 47. No. 2-3, 151-158, 1997. Cytogenetic: in vitro and in vivo cytogenetic analysis of the effects of cloprostenol on mammalian cells

IN VITRO I IN VIVO CITOGENETIČKA ANALIZA EFEKATA KLOPROSTENOLA NA ČELIJAMA SISARA

N. DELIĆ, M. ANDELKOVIĆ, B. SOLDATOVIĆ I DIJANA CVETKOVIĆ

SADRŽAJ

Ovim istraživanjima bio je da se izvrši evaluacija mogućih genotoksičnih efekata kloprostenola (Sinhrogal%) – sintetičkog analoga prostaglandina R2 alfa. Kloprostenol se koristi za sinhronizaciju estrusa domaćih životinja. Mutageni potencijal ove supstance ispitan je na humanim limfocitima periferne krvi (in vitro) i na limfocitima intramuskularno tretiranih mužjaka miševa soja BALB/cAnNCR (in vivo). Utvrđene su numeričke i strukturne hromazomske aberacije i određivan je mutageni indeks (procenat ćelija u mitozima). Kloprostenol nije ispoljio mutagene aktivnosti u okviru eksperimentalnih koncentracija i doza upotrebljenih u ovom istraživanju.

U ovom istraživanju bio je da se izvrši evaluacija mogućih genotoksičnih efekata kloprostenola (Sinrogal%) – sintetičkog analoga prostaglandina R2 alfa. Kloprostenol se koristi za sinhronizaciju estrusa domaćih životinja. Mutageni potencijal ove supstance ispitan je na humanim limfocitima periferne krvi (in vitro) i na limfocitima intramuskularno tretiranih mužjaka miševa soja BALB/cAnNCR (in vivo). Utvrđene su numeričke i strukturne hromazomske aberacije i određivan je mutageni indeks (procenat ćelija u mitozima). Kloprostenol nije ispoljio mutagene aktivnosti u okviru eksperimentalnih koncentracija i doza upotrebljenih u ovom istraživanju.