

UNIVERSITY OF BELGRADE FACULTY OF AGRICULTURE SERBIA

WITH A SUPPORT OF
EUROPEAN AQUACULTURE SOCIETY, EAS
AND
PANHELLENIC SOCIETY OF TECHNOLOGISTS ICHTHYOLOGISTS,
P.A.S.T.I.

**VI INTERNATIONAL CONFERENCE
“WATER & FISH”**

CONFERENCE PROCEEDINGS

**Faculty of Agriculture, Belgrade-Zemun, Serbia
June, 12 – 14. 2013.**

PUBLISHER

University of Belgrade, Faculty of Agriculture

FOR THE PUBLISHER:

Prof. Dr. Milica Petrović

EDITOR IN CHIEF

Prof. Dr. Vesna Poleksić

EDITORIAL BOARD

Prof. dr Zoran Marković, Prof. dr Vesna Poleksić, Dr Sadasivam Sashi Kaushik, Alistair Lane, Prof. dr Jana Pickova, Prof. dr Zdenek Adamek, Prof. dr Tomas Policar, Dr Fabrice Teletchea, Dr Zsigmond Jeney, Dr Tania Hubenova, Dr Aleksandar Joksimović, Doc. dr Nebojša Savić, Doc. dr Zorka Dulić, Doc. dr Ivana Živić, Doc. dr Renata Relić, Dr Aurelija Spirić, Dr Mirjana Lenhardt, Dr Svetlana Jeremić, Prof. dr Miroslav Ćirković

COMPUTER DESIGN

Nikša Tarle

TECHNICAL PREPARATION OF A COVER

Nikša Tarle

PRINTED BY

MONT PRESS, Novi Sad

NUMBER OF COPIES PRINTED

400 COPIES

ORGANIZATION OF THE CONFERENCE SUPPORTED BY:

Ministry of Education, Science and Technological Development of the Republic of Serbia

GENERAL SPONSOR

«Veterinary Institute, Subotica»

UNIVERZITET U BEOGRADU, POLJOPRIVREDNI FAKULTET,

**UZ PODRŠKU
EVROPSKOG DRUŠTVA ZA AKVAKULTURU, EAS
I
PANHELENSKOG DRUŠTVA TEHNOLOGA IHTIOLOGA**

**VI MEĐUNARODNA KONFERENCIJA
„VODA I RIBA“**

ZBORNİK PREDAVANJA

**Poljoprivredni fakultet, Beograd– Zemun, Srbija
12. – 14. jun, 2013. godine**

IZDAVAČ
Univerzitet u Beogradu, Poljoprivredni fakultet

ZA IZDAVAČA:
Prof. dr Milica Petrović

GLAVNI I ODGOVORNI UREDNIK
Prof. dr Vesna Poleksić

NAUČNI ODBOR
Prof. dr Zoran Marković, Prof. dr Vesna Poleksić, Dr Sadasivam Sashi Kaushik, Alistair Lane, Prof. dr Jana Pickova, Prof. dr Zdenek Adamek, Prof. dr Tomas Policar, Dr Fabrice Teletchea, Dr Zsigmond Jeney, Dr Tania Hubenova, Dr Aleksandar Joksimović, Doc. dr Nebojša Savić, Doc. dr Zorka Dulić, Doc. dr Ivana Živić, Doc. dr Renata Relić, Dr Aurelija Spirić, Dr Mirjana Lenhardt, Dr Svetlana Jeremić, Prof. dr Miroslav Ćirković

RAČUNARSKA OBRADA
Nikša Tarle

TEHNIČKO UREĐENJE KORICA
Nikša Tarle

ŠTAMPA
MONT PRESS, Novi Sad

TIRAŽ
400 primeraka

ODRŽAVANJE SKUPA POMOGLO:
Ministarstvo prosvete, nauke i tehnološkog razvoja Republike Srbije

GENERALNI SPONZOR
„Veterinarski zavod Subotica“

CONTENTS:

Kaushik, S.: FISH NUTRITION: IMPROVING NUTRIENT EFFICIENCY AND WATER USE	21
Lane, A.: THE EUROPEAN AQUACULTURE SOCIETY (EAS)	24
Pickova, J., Mraz, J., Zajic, T.: COMMON CARP AND AQUACULTURE ..	29
Marković, Z., Stanković, M., Dulić, Z., Rašković, B., Živić, I., Spasić, M., Vukojević, D., Relić, R., Poleksić, V.: CARP PRODUCTION IN SERVICE OF REINFORCEMENT OF SERBIA AGRICULTURE	33
Adámek, Z.: BIOMANIPULATION MEASURES ON CZECH RESERVOIRS AND THEIR PAST AND CURRENT SUCCESSFULNESS.....	39
Pucher, J., Focken, U.: INTENSIFICATION OF COMMON CARP CULTURE IN RURAL AREAS OF NORTHERN VIETNAM BY PLANT-BASED FEEDS WITH PROTEIN SOURCES OF DIFFERENT QUALITIES	44
Jovanović, R., Ivanov, D., Čolović, R., Vukmirović, Đ., Palić, D., Lević, J., Đuragić, O.: INFLUENCE OF FATTY ACID COMPOSITION OF FISH FEED ON FATTY ACID COMPOSITION OF CARP MEAT	50
Trbović, Z., Marković, Z., Petronijević, R., Milijašević, M., Spirić, D., Vranić, D., Spirić, A.: MULTIVARIATE ANALYSIS OF FATTY ACID PROFILES OF CARP MEAT DURING SEMI – INTENSIVE FARMING	56
Linhart, O., Kocour, M., Kašpar, V.: EDUCATION AND TRAINING IN AQUACULTURE AND WATER PROTECTION AT FACULTY OF FISHERIES AND PROTECTION OF WATERS, UNIVERSITY OF SOUTH BOHEMIA IN ČESKÉ BUDĚJOVICE, THE CZECH REPUBLIC	63
Poleksić, V., Dulić, Z., Stanković, M., Rašković, B., Spasić, M., Vukojević, D., Marković, Z.: HIGHER EDUCATION FOR AQUACULTURE/FISHERY AT THE FACULTY OF AGRICULTURE UNIVERSITY OF BELGRADE: IMPLEMENTATION OF THE FIRST BOLOGNA REFORM – LINKING THEORY AND PRACTICE	68
Teletchea, F., Fontaine, P.: BEYOND THE LEVELS OF DOMESTICATION IN FISH: MUST ALL FARMED SPECIES BE DOMESTICATED?	74

Żarski, D., Krejszeff, S., Palińska-Żarska, K., Kucharczyk, D.: CONTROLLED REPRODUCTION OF PIKEPERCH, SANDER LUCIOPERCA (L.) – ACHIEVEMENTS AND PROSPECTS	82
Polcar, T., Blecha, M., Kristan, J., Stejskal, V., Blaha, M.: COMBINATION OF INTENSIVE (RAS) AND EXTENSIVE (POND) AQUACULTURE FOR JUVENILE PRODUCTION IN PIKEPERCH (SANDER LUCIOPERCA).....	88
Cosson, J., Prokopchuk, G., Dzyuba, V., Fedorov, P.: FISH SPERMATOOZOA, PHYSICAL AND BIO-ENERGETIC INTERACTIONS WITH THEIR SURROUNDING MEDIA	92
Buza, E., Kolics, B., Kovács, B., Demény, F., Horváth, Á., Urbányi, B., Sándor, S., Müller, T.: ARTIFICIAL PROPAGATION AND REVEALED REPRODUCTION FEATURES OF WEATHERFISH (<i>MISGURNUS FOSSILIS</i>)	98
Mamedov, Ch., Sadiqbeyli, L., Akhundov, M., Hajiyev, R.: EVALUATION OF BIOLOGICAL INDICATORS OF YOUNG PERSIAN STURGEON (<i>ACIPENSER PERSICUS</i>) GROWN AT POND METHOD WITH DIFFERENT ARTIFICIAL FEEDS	104
Savić, N., Mikavica, D., Rogić, B.: THE GROWTH OF WEIGHT AND BODY LENGTH OF YOUNG RAINBOW TROUT (<i>ONCORHYNCHUS MYKISS</i> WAL.) ORIGINATING FROM DIFFERENT BROODSTOCK.....	111
Altinok, I., Kahraman, U., Capkin, E., Boran, H., Ozturk, R.: DEVELOPMENT OF LIVE VACCINES FROM YERSINIA RUCKERI BY aroA AND aroC GENE MUTATIONS.....	116
Ćirković, M., Novakov, N., Ljubojević, D., Bjelić-Čabrilo, O., Davidov, I.: FINDING OF THE NEMATODA LARVAE IN PREDATORY FISH SPECIES.....	118
Radosavljević, V., Jeremić, S., Jakić-Dimić, D.: SURVEY AND DIAGNOSTICS OF FISH DISEASES IN THE REPUBLIC OF SERBIA DURING THE PERIOD 2011-2012	124
Baltić, Ž.M., Bošković, M., Đorđević, V., Marković, R., Dimitrijević, M., Pavličević, N.: FISH-BORNE PARASITIC ZOOSES WITH SPECIAL REFERENCE TO ANTHROPOGENIC IMPACT	129
Wagenaar, G.M., Barnhoorn, I.E.J.: THE HEALTH STATUS AND EDIBILITY OF FISH FROM THREE HYPERTROPHIC IMPOUNDMENTS IN SOUTH AFRICA	136

- Relić, R., Lakić, N., Dulić, Z., Grubišić, M., Mladenović, V., Marković, Z., Poleksić, V.:** CONSUMERS' OPINION ABOUT EFFECTS OF REARING CONDITIONS AND STRESS ON FISH MEAT QUALITY 141
- Islami, H. R., Arab, N.:** EFFECT OF DIETARY VITAMIN C ON SURVIVAL AND GROWTH PERFORMANCE OF CASPIAN BROWN TROUT (*SALMO TRUTTA CASPIUS*) FINGERLINGS 148
- Skorić, S., Smederevac-Lalić, M., Višnjic-Jeftić, Ž., Hegediš, A., Mićković, B.:** RELATIONSHIPS OF OTOLITH SIZE TO TOTAL LENGTH OF THE BURBOT (*LOTA LOTA*) FROM THE DANUBE RIVER 158
- Obradović, S., Šarčević, B., Šekler, M., Đekić, V., Dekić, R., Veljović, N., Marković, M.:** OPTIMIZATION MODEL OF FISH GUARD SERVICE IN ORDER TO PROTECT AQUATIC SYSTEMS 164
- Nathanailides, C., Tsoumani, M., Kakali, F., Logothetis, P., Beza, P., Mayraganis, Th., Kanlis, G., Delis, G., Tiligadas, I., Chatziefstathiou, M.:** A CORRELATION BETWEEN ALKALINE PHOSPHATASE AND PHOSPHATE LEVELS WITH THE BIOMASS OF TROUT FARM EFFLUENTS 170
- Joksimović, A., Vrgoč, N., Krstulović Šifner, S., Isajlović, I., Ikica, Z., Marković, O.:** INTERNATIONAL COOPERATION PROJECTS IN THE RESEARCH OF THE ADRIATIC SEA 176
- Chatziefstathiou, M., Spilanis, I.:** TOWARDS THE IMPLEMENTATION OF EUROPEAN UNION'S NEW INTEGRATED MARITIME POLICY IN GREECE: BLUE GROWTH THROUGH MARINE AQUACULTURE FOR THE SUSTAINABLE DEVELOPMENT OF THE ISLANDS 185
- Fernandez-Jover, D., Sanchez-Jerez, P.:** USE OF ARTIFICIAL HABITATS AS RECRUITMENT SITES BY JUVENILE WILD FISH: EFFECTS OF FISH FARMS ON FISH DIET AND OTOLITH GROWTH 193
- Pešić, A., Regner, S., Mandić, M., Ikica, Z., Đurović, M., Joksimović, A., Marković, O.:** BIOLOGICAL CHARACTERISTICS OF ANCHOVY (*ENGRAULIS ENCRASICOLUS*) IN BOKAKOTORSKA BAY (MONTENEGRO) 197
- Vrgoč, N., Joksimović, A., Krstulović Šifner, S., Isajlović, I., Pešić, A.:** STATE OF BENTHIC (DEMERSAL) RESOURCES IN THE EASTERN COAST OF THE ADRIATIC SEA 204

Ikica, Z., Krstulović Šifner, S., Joksimović, A., Marković, O., Pešić, A., Isajlović, I., Vrgoč, N.: DISTRIBUTION OF MUSKY OCTOPUS (<i>ELEDONE MOSCHATA LAMARCK, 1798</i>) (CEPHALOPODA: OCTOPODA) IN THE SOUTH-EASTERN ADRIATIC	210
Jović, M., Marković, J., Stanković, S.: EVALUATING THE HEALTH RISKS ASSOCIATED WITH CONSUMING OF CULTIVATED / WILD MUSSELS FROM BOKA KOTORSKA BAY	217
Ljubojević, D., Ćirković, M., Jovanović, R., Novakov, N., Đorđević, V., Trbović, D., Spirić, A.: EFFECT OF RAPESEED OIL IN FISH FEED ON MEAT QUALITY OF COMMON CARP	222
Stanković, M., Ljubobratović, U., Lakić, N., Dulić, Z., Spasić, M., Vukojević, D., Gruberinić, D., Marković, M., Marković, Z.: POSSIBLE REPLACEMENT OF FISH MEAL BY SOY CONCENTRATE IN FEED FOR TROUT	227
Marković, G., Ćirković, M., Lujić, J., Pantović, J.: COMPARISON OF THE NUTRITIVE VALUE OF WHEAT AND TRITICALE IN FISH DIET	233
Radovanović, V., Demin, M., Žarković, B., Stikić, R., Milovanović, M.: APPLICATION OF A NEW CROP-QUINOA FOR FISH FEED	238
Pavličević, M., Stanojević, S., Barac, M., Pešić, M., Vučelić-Radović, B.: POTENTIAL OF SECONDARY RAW MATERIAL – SOYBEAN OKARA FOR USE AS FISH FEED	243
Palińska-Żarska, K., Żarski, D., Krejszeff, S., Kucharczyk, D.: CURRENT STATUS AND PROSPECTS IN BURBOT, <i>LOTA LOTA L.</i> , AQUACULTURE	249
Özgür E.M., Bayir, İ.: THE EFFECTS OF DIFFERENT STRIPPING TERMS ON SPERM QUALITY PARAMETERS OF RAINBOW TROUT (<i>ONCORHYNCHUS MYKISS</i>)	255
Ljubobratović, U., Kucska, B., Feledi, T., Poleksić, V., Marković, Z., Lenhardt, M., Rónyai, A.: COMBINED METHODS FOR ARTIFICIAL REPRODUCTION OF PIKEPERCH, SANDER LUCIOPERCA	261
Mahsamohamadizadeh K.H., Mehdi Shamsaie M., Abdollahtabar, Y.: EFFECTS OF WATER CURRENT ON SOME GROWTH FACTORS AND WATER QUALITY IN A CLOSED TROUT CULTURE SYSTEM	267

- Ghiasi, F. Rizbandi, T.:** DETERMINATION OF VHS VIRUS TITER IN EXPERIMENTALLY INFECTED RAINBOW TROUT, *ONCORYHINCHUS MYKISS*..... 276
- Karagiannis, P., Grigorakis, K., Nathanailides, C., Fountoulaki, E., Basilaki, A., Lopez-Albors, O., Petridis, D.:** THE INFLUENCE OF DIFFERENT THERMAL CONDITIONS ON THE PROXIMATE COMPOSITION OF RAINBOW TROUT FLESH FROM TWO FISH FARMS IN NW GREECE 291
- Milošković, A., Pavlović, M., Kovačević, S., Radojković, N., Simić, S., Simić, V.:** THE PRESENCE OF ZINC IN MUSCLE TISSUE OF PRUSSIAN CARP AND BREAM IN THE GRUŽA AND BOVAN RESERVOIRS 283
- Skorić, S., Đikanović, V., Marković, G., Hegediš, A.:** CONCENTRATIONS OF 16 ELEMENTS IN TISSUES (LIVER, MUSCLE, SCALES) OF PRUSSIAN CARP (*CARASSIUS GIBELIO*, BLOCH, 1782) IN MEDJUVRŠJE RESERVOIR, SEASONAL ASPECT 288
- Obradović, S., Ivanc A., Đekić, V., Šekler, M., Živkov-Baloš, M., Veljović, N., Šarčević, B.:** CONTENT AND DISTRIBUTION OF HEAVY METALS IN ORGANS AND TISSUES OF FRESHWATER FISH IN THE ZAPADNA MORAVA RIVER 295
- Jović, M., Onjia, A., Stanković, S.:** METAL POLLUTION INDEX AS A TOOL FOR ASSESSING WATER QUALITY OF BOKA KOTORSKA BAY 300
- Ősz, Á., Kovács, B., Kánainé Sipos, D., Kaczkó, D., Herzinyák, Z., Snoj, A., Bogataj, K., Urbányi, B., Horváth, Á.:** ORIGIN AND GENETIC STRUCTURE OF SEVERAL HUNGARIAN WILD AND DOMESTICATED BROWN TROUT POPULATIONS BASED ON PCR-RFLP AND MICROSATELLITE MARKERS 305
- Muhamedagić, S., Korjenić, E., Hamzić, A., Drešković, N., Mustafčić, A., Kapo, I., Lepar, D., Habibović, E.:** ICTHYOFAUNA OF KRIVAJA RIVER CATCHMENT AREA..... 310
- Bećiraj, A., Šarić-Kadić, I., Ičanović, I., Halimović, S., Đekić, R., Galijašević, E.:** FISH BIODIVERSITY IN THE RIVER KRUŠNICA 321
- Tomljanović, T., Piria, M., Šprem, N., Matulić, D., Zanella, D.:** FINDING OF STERLET (*ACIPENSER RUTHENUS*) IN THE SAVA RIVER NEAR ZAGREB 326

- Zekić Žujo, D.:** AGE STRUCTURE OF POPULATION OF ENDEMIC SPECIES CHONDROSTOMA PHOXINUS HECKEL, 1843 (TELEOSTEI: OSTARYOPHYSI, CYPRINIDAE) FROM BUSKO LAKE 331
- Krizmanić, J., Subakov Simić, G., Predojević, D.:** ALGAE AS WATER QUALITY BIONDICATORS OF THE RIVER DJETINJA..... 342
- Bjelanović, K., Živić, I., Dulić, Z., Živić, M., Đorđević, J., Marinković, S., Marković, Z.:** WATER QUALITY ASSESSMENT IN THE RAŠKA RIVER BASED ON ZOOBENTHOS AND ZOOPLANKTON ORGANISMS AS BIOIDICTORS 349
- Dulić, Z., Grubisić, M., Stanković, M., Živić, M., Arsić, L., Marković, Z.:** EFFECTS OF DIFFERENT ORGANIC MANURES ON BODY SIZE AND PRODUCTION OF *DAPHNIA MAGNA*..... 358
- Živić, I., Bjelanović, K., Marinković, S., Jovanović, J., Marković, Z.:** PRODUCTION OF MACROZOOBENTHOS IN THE RAČA RIVER UPSTREAM AND DOWN STREAM FROM TROUT FARM ... 364
- Terzi, E., Gedik, K., Verep, B., Yandi, I.:** ASSESSMENT OF MARINE WATER QUALITY: EFFECT OF AQUACULTURE AND DOMESTIC SEWAGE DISCHARGE 372
- Isajlović, I., Vrgoč, N., Krstulović Šifner, S., Marković, O., Pešić, A.:** ASSESSMENT OF NORWAY LOBSTER - NEPHROPS NORVEGICUS (LINNAEUS, 1758) POPULATIONS IN THE ADRIATIC SEA USING ALTERNATIVE METHODS. 374
- Krstulović Šifner, S., Ikica, Z., Đurović, M., Vrgoč, N., Isajlović, I., Joksimović, A.:** DISTRIBUTION, ABUNDANCE AND POPULATION STRUCTURE OF THE COMMON CUTTLEFISH, *SEPIA OFFICINALIS* LINNAEUS, 1758, IN THE ADRIATIC SEA 380
- Lenas, S. D., Triantafyllou, J. D., Logothetis, P., Kakali, F., Nathanailides, C.:** THE INFLUENCE OF DIFFERENT ENVIRONMENTAL CONDITIONS ON FLESH QUALITY INDICES OF FARMED SEA BASS, IN NW GREECE 386
- Mandić, M., Drakulović, D., Huter, A., Petović, S., Mandić, S.:** TEHNOLOGY OF MUSSEL (*MYTILUS GALLOPROVINCIALIS*) AND OYSTER (*OSTREA EDULIS*) FARMING IN BOKA KOTORSKA BAY 388
- Barišić, J., Rathman, M., Čož-Rakovac, R., Strunjak-Perović, I., Topić Popović, N., Sauerborn-Klobučar, R., Jendriš Škrliak, E., Mihaldinec, D.:** MAGNETIC RESONANCE IMAGING - A NEW APPROACH IN INVESTIGATIONS OF ENDANGERED MARINE BIVALVES..... 393

Hajdarević, E., Hasković, E., Hercegovac, A., Adrović, A., Skenderović, I.: LEUKOCYTES OF COMMON BREAM (<i>ABRAMIS BRAMA</i> , LINNEAUS, 1758) FROM THE ARTIFICIAL MODRAC LAKE (BOSNIA & HERZEGOVINA) IN VARIOUS SEASONS.....	395
Dikanović, V., Skorić, S., Cakić, P.: REPRESENTATIVES OF TAPEWORMS (CESTODA) OF FISHES IN BELGRADE SECTION OF THE DANUBE RIVER	402
Dekić, R., Ivanc, A., Savić, N., Manojlović, M., Četković, D., Obradović, S.: HEMATOLOGICAL EVALUATION OF RAINBOW TROUT (<i>ONCORHYNCHUS MYKISS</i>) FINGERLINGS FROM DIFFERENT HATCHERIES	409
Novakov, N., Ćirković, M., Ljubojević, D., Babić, R., Aleksić, N.: PATHOLOGICAL CHANGES WHICH <i>THELOHANELLUS NIKOLSKII</i> CAUSE ON THE FINS, SCALES AND SKIN OF COMMON CARP.....	414
Capkin, E., Terzi, E., Boran, H., Altinok, İ., Başçınar, N.: BACTERIAL DISEASE SCREENING OF CULTURED HORSE MACKEREL (<i>TRACHURUS MEDITERRANEUS</i>) IN SEA-CAGES IN SOUTHERN BLACK SEA REGION OF TURKEY.....	420
Terzi, E., Capkin, E., Altinok, İ.: DISTRIBUTION OF SULFONAMIDE RESISTANCE GENES IN BACTERIA ISOLATED FROM RAINBOW TROUT IN TURKEY	422
Tiligadas, I., Moutopoulos, K. D., Chatziefstathiou, M., Tsoumani, M.M.: HELLENIC MARICULTURE SAFETY AND HEALTH: INITIAL REPORT ON WORK RELATED INJURIES AND ILL HEALTH.....	424
Aleksić-Agelidis, A., Ljubojević, D., Babić, J., Lujić, J., Novakov, N., Marković, M.: DRESSING PERCENTAGE OF 3-YEAR OLD CARP FROM CAGE PRODUCTION SYSTEM.....	429
Pavličević, N., Đorđević, V., Dimitrijević, M., Bošković, M., Marković, R., Baltić, Ž.M.: LACTIC ACID BACTERIA: EFFECT ON THE QUALITY AND SAFETY OF FISHERY PRODUCTS	436
Bianchini, L. M., Pigliarieni, E., Rambaldi, E., Argenti, L., Menesatti, P., Costa, C.: A NON INVASIVE METHOD FOR TRACING ORGANIC FISH.....	443

- Zohreh, M., Ali, M.T., Amirreza, S.:** THE SURVEY OF SOME BACTERIAL CONTAMINATION IN FROZEN SEMI-COOKED READY TO EAT SEA FOOD PRODUCTS IN ALBORZ PROVINCE- IRAN 452
- Smederevac-Lalić, M., Zarić, V., Hegediš, A., Lenhardt, M., Mićković, B., Višnjić-Jeftić, Ž., Pucar, M., Cvijanović, G.:** THE MARKETING CHANNELS OF FISH CAUGHT IN LARGE SERBIAN RIVERS..... 457
- Rašković, B., Vukojević, D., Spasić, M., Živić, I., Stanković, M., Marković, Z., Poleksić, Z.:** EFFECT OF STRESS ON PRESENCE OF DEVELOPING NEPHRONS OF COMMON CARP REARED IN SEMIINTENSIVE SYSTEM..... 463
- Grubišić, M., Dulić, Z., Đorđević, J., Bjelanović, K., Spasojević, I., Relić, R., Marković, Z.:** SEASONAL VARIATIONS AND DIVERSITY OF CLADOCERA (CRUSTACEA) IN NATURAL, DOMESTIC WASTEWATER TREATMENT LAGOONS..... 467
- Spasojević, I., Dulić, Z., Grubišić, M., Đorđević, J., Ljubobratović, U., Raičević, V., Marković, Z.:** TOTAL COLIFORM AND FECAL COLIFORM COMMUNITY DYNAMICS IN NEWLY BUILT WASTEWATER TREATMENT LAGOONS..... 473
- Relić, R., Lakić, N., Mladenović, V., Dulić, Z., Grubišić, M., Poleksić, V.:** FISH WELFARE FROM PERSPECTIVE OF CONSUMERS IN SERBIA (PRELIMINARY RESULTS) 479
- Amiri, M, Hosseini, H., Islami, R. H.:** IDENTIFICATION AND DISTRIBUTION OF SEA CUCUMBERS IN THE CHABAHAH BAY, IRAN..... 487

FISH-BORNE PARASITIC ZOOSES WITH SPECIAL REFERENCE TO ANTHROPOGENIC IMPACT

MILAN Ž. BALTIĆ¹, MARIJA BOŠKOVIĆ¹, VESNA ĐORĐEVIĆ², RADMILA MARKOVIĆ¹, MIRJANA DIMITRIJEVIĆ¹, NATAŠA PAVLIČEVIĆ³

¹*Faculty of Veterinary Medicine, Bulevar oslobođenja 18, 11 000 Belgrade, Serbia*

²*Institute of Meat Hygiene and Technology, Kačanskog 13, 11 000 Belgrade, Serbia*

³*Veterinary Institute „Subotica“, Segedinski put 88, 24 000 Subotica, Serbia*

ZOONOTSKE PARAZITAZE RIBA SA POSEBNIM OSVRTOM NA ANTROPOGENI UTICAJ

Apstrakt

Parazitska oboljenja koja se prenose mesom ribe dugo vremena nisu dobijala zasluženu pažnju, pre svega jer su ove zoonoze bile vezane za zemlje tropskog i suptropskog pojasa, naročito azijske zemlje u kojima riba predstavlja veoma značajan izvor proteina za stanovništvo ruralnih sredina. Međutim, kao posledica globalnog zagrevanja dolazi do promena u ekosistemima voda i širenja geografskih granica u kojima se određeni paraziti pojavljuju. Širenju parazita doprinosi i fenomen globalizacije, razvijene saobraćajne mreže, komercijalni internacionalni hipermarketi, ali i demografske promene stanovništva od kojih najveći značaj imaju migracije. Na povećanje populacije pod rizikom utiču i promene kulturoloških navika i kulinarske prakse, naročito sve češća konzumacija sirove, marinirane, sušene ili na drugi način temperaturno nedovoljno obrađene ribe i proizvoda od ribe. Tokom poslednje decenija zabeležen je progresivan rast broja ljudi koji se hrane u restoranima, kioscima brze hrane ili kod uličnih prodavaca, koji često ne poštuju pravila o bezbednosti hrane. Ovakve navike posledica su brzog tempa života i sve većeg opterećenja današnjeg čoveka, ali i popularizacija raznih kuhinja, naročito azijske kuhinje, u svetu, ali i u našoj zemlji.

Prilagođavajući prirodu svojim potrebama, čovek, kako na direktan, tako i na indirektan način, utiče na rasprostranjenost samih parazita. Izmena strujanja vode, koje nastaje pregrađivanjem reka branama prilikom izgradnje hidroelektrana i povišenje temperature vode, usled izgradnje termoelektrana, dovodi do supstitucije jednih vrsta fitoplanktona, zooplanktona i riba drugim vrstama. Promene flore i faune vodenog ekosistema ima direktan uticaj i na ptice koje se njima hrane, a koje su često pravi ili prelazni domaćini parazita. Ne retko, zbog narušavanja ekosistema i introdukcije novih vrsta riba javljaju se novi paraziti koji nisu karakteristični za neko područje. Zagađenje reka teškim meta-

lima je jedan od značajnih antropogenih faktora koji utiču na prisustvo parazita, ali i na stepen akumulacije teških metala u mišićima riba. Naime, paraziti ne naseljavaju vodu zagađenu metil živom, ali istraživanja su pokazala da ukoliko su ribe već invadirane parazitima u trenutku zagađanja, akumuliraju živu u svom organizmu u manjim koncentracijama od one ribe koja nije inficirana.

U epidemiologiji zoonotskih parazitoza koje se prenose mesom riba najveći značaj imaju helminti, pre svega klase *Trematoda* vrste *Clonorchiosis sinensis* i *Opisthorchis spp.* *C. sinensis* ima izuzetan socialno-ekonomski značaj u određenim delovima Azije i predstavlja veliki zdravstveni problem. Procenjuje se da je oko 35 miliona ljudi u toku poslednje decenije inficirano ovim parazitom na globalnom nivou. *O. felineus* se nalazi u Kazakstanu, Rusiji, Nemačkoj, Italiji, Španiji, Grčkoj, Makedoniji i Albaniji, dok je *O. viverrini* karakteristična za područje Kambodže, Laosa, Vijetnama i Tajlanda. Prema podacima WHO iz 1995. godine preko 18 miliona svetskog stanovništva je obolelo od parazitoza izazvanim trematodama koje su prenete mesom riba, a njihova procena je da je pod rizikom od inficiranja više od pola milijarde stanovnika. Difilobotrijaza je parazitoza izazvana helmintima iz klase *Cestoda*, roda *Diphyllobothrium*, od kojih *Diphyllobothrium latum* predstavlja najčešći uzročnik infekcija ljudi. Ovi paraziti rasprostranjeni su u Evropi, Severnoj i Južnoj Americi i Aziji. U Evropi *D. latum* je rasprostranjen u oblastima oko alpskih jezera, a najveći broj slučajeva prijavljen je u Francuskoj, Švajcarskoj i Italiji. Od nematoda najveći značaj imaju kosmopolitski rasprostranjene *Anisakis simplex* i *Pseudoterranova decipiens*. Visok procenat parazitskih infekcija protiče subklinički ili asimptomatski, a manji broj se manifestuje kroz simptome poremećaja digestivnog trakta, dok se difilobotrijaza manifestuje hipovitaminozom B12 vitamina. Zbog velikog broja neprijavljenih slučajeva, ne uvek jasno izraženih kliničkih simptoma i manjka metoda za otkrivanje uzročnika, ove parazitoze se, uglavnom, ne dijagnostifikuju, naročito u delovima sveta za koji nisu karakteristične.

Ključne reči: helminti, ekosistem, klimatske promene, zagađenje vode
Keywords: helminths, ecosystem, climate changes, water pollution

INTRODUCTION

Compared with other well-studied parasitic diseases, fish-borne parasitic zoonoses have been public health orphans in the world of research funding, especially because these zoonoses have been limited for the most part to populations living in low- and middle-income countries (Chai et al., 2005). Zoonotic parasites are a significant food safety problem, particularly in Asia, because fish are a very important source of protein for people living in rural areas (Thien et al., 2009). In an attempt to change the nature, so they can adjust it for their needs, humans, directly or indirectly, affect parasites occurrence. As a result of global warming it is expected for some species to become locally extirpated and experience range contractions, while introductions of others, including potentially harmful pathogens, will occur in both freshwater and marine systems (Marcogliese, 2001.). Geographical barriers are slowly being breached by international travel developing into a major industry, by improved refrigerated food transport, by growing international markets, by increased tourism and by demographic changes such as migration (Chai et al., 2005; Dorny et al., 2009). Populations at risk are expanding also as a result of changes in cultural habits and culinary practices, such

as the increase of eating raw, marinated, smoked, salted, pickled, airdried or undercooked fish meat and fish products. Eating sushi, sashimi, koi-pla, kinilaw and ceviche is becoming increasingly fashionable in many countries (Macpherson, 2005). In the last decade it has been noted increasing in the number of people eating meals prepared in restaurants, canteens and fast food outlets as well as from street food vendors who do not always respect food safety (Dorny et al., 2009). All of these has led to a dramatic rise in the incidence of a large number of fish-borne zoonotic parasitic infections in previously uninfected ethnic groups (Macpherson, 2005). The role of some factors like ages, malnutrition, HIV infection and other underlying medical conditions should not be neglected in etiopathogenesis of fish-borne parasitic zoonoses (Dorny et al., 2009).

Fish parasites Trematodiasis

Trematodiasis is the infection of humans caused by trematode parasites, known also as a liver flukes. The ones that cause the most common infection in humans are *Clonorchis sinensis*, *Opisthorchis viverrini* and *Opisthorchis felineus*. The life cycles of these trematode species are similar. Eggs are shed with the feces of the definitive host (man or other piscivorous animals). If the eggs reach water they develop into miracidia which are ingested by a snail, the first intermediate host, in which they develop and ultimately shed into the water as motile cercariae which penetrate into the muscle tissue of a fish, the second intermediate host. Human infection takes place through the consumption of under-processed fish containing the infective stage of the parasite (Santos and Howgate, 2011). *Clonorchis sinensis* (Chinese liver fluke) is widely distributed in East Asia and in this region present the most important species of fish-borne zoonotic parasite. *Opisthorchis viverrini* is highly prevalent in Southeast Asia including Thailand, Laos, Cambodia and Vietnam, and there are about 9 million people who are infected globally (Chai et al., 2005). *Opisthorchis felineus* is prevalent in Spain, Italy, Albania, Greece, France, Macedonia, Switzerland, Germany, Poland, Russia, Turkey, and in parts of the former Soviet Union (Chai et al., 2005, Macpherson, 2005). The major clinical problems in infected humans are cholangitis, choledocholithiasis, pancreatitis and cholangiocarcinoma and they are associated with the long chronic infections, but severe infections can cause jaundice, portal hypertension, ascites, gastrointestinal bleeding and formation of gallstones (Chai et al., 2005; Dorny et al., 2009).

Cestodiasis

Diphyllobothriasis is the most important fish-borne zoonosis caused by a cestode parasite. At least 13 of about 50 species of *Diphyllobothrium* have been reported from humans, but most important is *Diphyllobothrium latum*. The egg hatches in water, the motile embryo coracidium is ingested by a copepod, and develops to the proceroid. When an infected copepod is ingested by the second intermediate host (freshwater, anadromous or catadromous fish), the larva is released, enters the tissues of the host and develops to the plerocercoid which is infective for the final host (Chai et al., 2005). These parasites are found in Europe, North and South America and Asia. About 20 million people are infected worldwide (Dorny et al., 2009). In Europe, Scandinavian countries present 'hot spot', and Switzerland, Sweden, Finland and Estonia report more than 10 cases per year, while Lithuania, Poland, Hungary, Italy and France average 2–10 cases annually (Chai et al., 2005). Most of the cases in Switzerland, France and Italy occur in the Alpine Lakes region. Diphyllobothriasis in humans may often be asymptomatic,

but in some cases it may cause diarrhea, abdominal pain and megaloblastic anaemia as a result of B12 deficiency (Macpherson, 2005; Dorny et al., 2009).

Nematodiasis

Of all nematode parasites most commonly involved in human infections is *Anisakis simplex*, while *Pseudoterranova decipiens* is less frequent. Infections with the *Anisakis physeteris* and *Contracaecum spp.* are sporadic (Dorny et al., 2009). The life cycle of *Anisakidae* involve small crustaceans as the first intermediate host, fishes and cephalopods as the second host, and marine mammals as the definitive host. Humans are accidental hosts that become infected by eating larvae contained within the paratenic host (Lima dos Santos and Howgate, 2011). *A. simplex* and *Pseudoterranova spp.* are found in the northern, north east and north west Atlantic, northern Pacific, in the Mediterranean Sea, Norwegian Sea and Barents Sea. Anisakiasis is prevalent in north Asia and western Europe (Netherlands, Germany, France and Spain) (Chai et al., 2005). *A. simplex* causes an acute or chronic infection followed by abdominal pain, nausea, vomiting, and diarrhea and some patients develop syndromes exhibiting clinical manifestations of allergy (Lima dos Santos and Howgate, 2011).

Gnathostomiasis is primarily a disease of the skin caused by migration of the larvae of a *Gnathostoma* nematode and it is known to occur mainly in South East Asia, but has been reported as a cause of human infection in Peru, Ecuador and Mexico. It is often reported in tourists returning from endemic areas where they have eaten undercooked freshwater fish (Nawa et al., 2005; Dorny et al., 2009).

Anthropogenic impact

Climate change

Climate change can occur as a result of natural and anthropogenic causes. Temperature change, alterations in water levels, stratification, ice cover, acidification, ultraviolet radiation are some of the factors that affect parasites of aquatic organisms, their life cycles and distribution. Climate change has been predicted to be one of the major drivers of biodiversity change, especially in lakes (Marcogliese 2001). Global climate change with rising sea-levels and water temperatures that may result in changes in ocean circulation and a decrease in salinity may also cause measurable effects on fish parasite composition and biogeography. Climate warming affect interactions between host and parasites by increasing pathogen development rates, transmission and number of generation times per year, raising the overwinter survival rate of the pathogen and increasing the host susceptibility to thermal stressors. Also, increasing temperature alter the seasonality and biogeographical range of many species, including the hosts and the parasites (Palm, 2011).

Aquatic ecosystem pollution

Aquatic ecosystems present the most sensitive systems on earth (Palm, 2011).

Pollution can affect parasitism in fish in two different ways. It can increase parasitism if host defence mechanisms are negatively affected by increasing host susceptibility, or by simply increasing the population densities of suitable intermediate or final hosts. Some studies have focused on parasite-induced stress that was measured as an increased stress hormone concentration and it often present the first step in a series of physiological reactions such as a depressed immune response or other fundamental physiological

reactions. Pollution can also decrease parasitism provided that infected hosts suffer more from environmental exposure than do uninfected hosts or pollution drives the necessary intermediate and final hosts to become extinct (Marcogliese, 2001). If the fish is living within a polluted environment where some pollutants also enter the fish, the concentration of these substances in the immediate surroundings of the parasite can also increase and accumulate in the parasite's tissue, proving that fish parasites closely interact with the metabolisms of their host. Parasites are more susceptible to the particular pollutant, such as heavy metals, than their host and some fish parasites can accumulate pollutants in a much higher concentration (to several thousand-fold higher levels) as their host organisms. That is one of the reasons why parasites, especially helminths, can serve as biological indicators of water pollution (Palm, 2011; Marcogliese, 2001).

Industrialisation, technology and transport development

The creation of large freshwater reservoirs for hydroelectrical, irrigation, flood control or drinking water purposes present one of the most dramatic and widespread anthropogenic impacts on the natural environment (Morley, 2007). In Republic of Serbia the biggest hydroelectric power station and the biggest reservoir at the same time is Djerdap, Gate I, on the Danube River. Dams construction destabilizes the aquatic environment for a number of years which has serious implications for the aquatic area. There is a replacement of rheophilic fauna with limnophilic fauna. This changes affect not only phytoplankton and zooplankton but also a fish, especially because the dams make it difficult for migratory fish to enter the reservoir which are reflected to changes in their parasite fauna such as elimination, reduction or emergence of species which are not endemic for that area (Morley, 2007). There are anthropogenic impacts that may directly affect the physical parameters of the water and consequently parasites. For example, thermal power station Nikola Tesla (TENT) on the Sava River near Obrenovac, affect water temperature. As a result of cooling water from TENT A and B water temperature in Sava River is for 2 °C higher than water in Danube River. Because of higher temperatures there is no ice cover on Sava River. Missing of ice cover has an indirect influence on parasitic fish fauna. In these conditions fish don't reduce feeding and migratory birds may remain present during all seasons, creating the potential for parasite transmission throughout the year. Also, on the higher temperature duration of parasite life cycle is shorter (Marcogliese, 2001).

Demographic and tourism habits changes

As a result of globalisation, new technology and improved transport, in the last few decades it has been noted increasing number of immigrants from developing parts of the world to developed countries in search of opportunities and better life. This mobility implies also importing different cultures, health beliefs, food preferences that can present a risk factors not only for immigrants but for public health (Broglia and Kapel, 2011). An interesting variation on the potential risks associated with the immigration is the report of *O. felineus* infections in a family of Russian immigrants residing in Israel. One of the members of the family brought back from a visit to the family's former home, in an endemic area of Siberia, an infected fish which was then consumed locally (Chai et al., 2005).

Some changes of the twentieth century, like tourism and international travel, have a great role in increasing number of not only fish-borne parasitic zoonoses, but of all parasitic diseases. In 2008, the World Tourism Organization reported that international

tourist arrivals reached 924 million, and this number was expected to increase to 1.6 billion by the year 2020. Far and exotic destination that once was available only for high class of society, nowadays present popular places for vacation. Travellers dining in street restaurants can be expected to have much higher risks of infections with various parasites, especially in Asia (Broglia and Kapel, 2011).

Changes in cultural habits, culinary practices and life style

Human behaviour and changes in live style may affect public health on many ways. Until recent eating of raw fish was part of traditions, cultural and religious beliefs, but nowadays it presents part of everyday life, not only in Asia but all over the world. Eating sushi, sashimi (traditional Japanese dishes), koi-pla, kinilaw and ceviche is becoming increasingly fashionable in many countries as a result of globalisation and media, which play a significant role in popularisation of exotic foods (Broglia and Kapel, 2011; Macpherson, 2005). People more often consume raw or undercooked foods in the so called street-food restaurants or take-away, that do not always respect food safety standards (WHO, 2002). But it is not always a fast food products that present a risk factor for fish borne parasitic zoonoses. The increased awareness of healthy dietary regimes to prevent cardiovascular disease and cancer has raised fish consumption, including fish oil, omega-3 fatty acids, which has led to more aquaculture products from Asian markets being now globally traded and consumed (Broglia and Kapel, 2011).

Control Measures

A number of control measures are used to protect fish products from parasites (Orlandi et al., 2002). Evisceration of fish soon after catching prevents the larvae from migrating from the intestinal tract to the muscle tissue of the fish (Baltić et al., 2005). Moreover, if there are parasites in fish tissue they can be killed by using physical or chemical treatments. Salting and marinating are the chemical treatments most commonly used to inactivate viable helminth larvae, but their efficiency depends on concentration of salt and duration of treatment period. Vegetable products such as shogaol and gingerol extracted from *Zingiber officinale* are able to kill *A. simplex*, but only under specific conditions (EFSA, 2010). Freezing (-35°C for 15 h or -15°C for at least 96 h) and cooking (60°C for at least 1 min.-core temperature) remain the reference processes guaranteeing the destruction of larvae. Other physical treatments which can be used for killing helminth larvae are high hydrostatic pressure, irradiation (for some parasites) and hot smoking (EFSA, 2010; Orlandi et al., 2002). Low voltage current is patented in Spain in 2005, but this method is not currently available and requires further research (EFSA, 2010).

CONCLUSIONS

Some natural changes like global warming, but more important anthropogenic impacts such as pollution, dams construction or human behaviour affect parasites life cycles, distribution, and prevalence, making the multiple routes of transmission more complicate and difficult to understand. As a result of these impacts, fish borne parasitic zoonoses present global increasing, not only health but also economic problem.

ACKNOWLEDGEMENTS

This paper was supported by the Ministry of Education and Science, Republic of Serbia, Project TR 31011.

REFERENCES

Baltić Ž. M., Kilibarda Nataša, Teodorović V., Dimitrijević Mirjana, Karabasil N., (2005): Paraziti riba i zdravlje ljudi. II International conference „Fishery“ Faculty of agriculture University of Belgrade, 10-12 February, 155-166.

Broglia A., Kapel C., (2011): Changing dietary habits in a changing world: Emerging drivers for the transmission of foodborne parasitic zoonoses, *Veterinary Parasitology* 182, 2– 13.

Chai Jong-Yil, Murrell K. Darwin, Lymberyc J. Alan, (2005): Fish-borne parasitic zoonoses: Status and issues, *International Journal for Parasitology*, 35, 1233–1254.

Dorny P., Praet N., Deckers N., Gabriel S. (2009): Emerging food-borne parasites, *Veterinary Parasitology* 163, 196–206.

European Food Safety Authority (EFSA), (2010): Scientific opinion on risk assessment of parasites in fishery products and EFSA Panel on Biological Hazards (BIOHAZ), *EFSA Journal*, 8,4,1543.

Lima dos Santos, C.A.M., Howgate, P., (2011): Fishborne zoonotic parasites and aquaculture: A review, *Aquaculture*, doi:10.1016/j.aquaculture.2011.05.046

Macpherson N.L. Calum, (2005): Human behaviour and the epidemiology of parasitic zoonoses, *International Journal for Parasitology*, 35, 1319–1331.

Marcogliese J. David, (2001): Implications of climate change for parasitism of animals in the aquatic environment, *Can. J. Zool.* 79, 1331–1352.

Morley J. Neil (2007): Anthropogenic Effects of Reservoir Construction on the Parasite Fauna of Aquatic Wildlife, *EcoHealth* 4, 374–383.

Nawa Yukifumi, Christoph Hatz, Johannes Blum, (2005): Sushi Delights and Parasites: The Risk of Fishborne and Foodborne Parasitic Zoonoses in Asia, *Clinical Infectious Diseases*, 41, 1297–1303.

Orlandi, P. A., Chu, D. M. T., Bier, J. W., Jackson, G. J., (2002): Parasites and the food supply, *Food technology*, 56, 4, 72-79.

Palm, Harry W., (2011): “Fish Parasites as Biological Indicators in a Changing World: Can We Monitor Environmental Impact and Climate Change?.” *Progress in Parasitology*. Springer Berlin Heidelberg, 223-250.

Thien Cu Pham, Anders Dalsgaard, Nguyen Thanh Nhan, Annette Olsen, K. Darwin Murrell, (2009): Prevalence of zoonotic trematode parasites in fish fry and juveniles in fish farms of the Mekong Delta, Vietnam, *Aquaculture*, 295, 1–5.

WHO, (2002): Foodborne Diseases, Emerging. Fact sheet no. 124.

CIP - Каталогизација у публикацији
Народна библиотека Србије, Београд

639.2/.3(082)

INTERNATIONAL Conference „Water & Fish“ (6 ;
2013 ; Beograd)

Conference Proceedings / VI International
Conference „Water & Fish“, June, 12-14.2013.
; [editor in chief Vesna Poleksić]. -
Belgrade : Faculty of Agriculture, 2013 (Novi
Sad : Mont Press). - 455 str. : ilustr. ; 30
cm

Na spor. nasl. str.: Zbornik predavanja. -
Tiraž 400. - Bibliografija uz svaki rad.

ISBN 978-86-7834-155-7

а) Рибарство - Зборници
COBISS.SR-ID 198755596

ISBN 978-86-7834-155-7



9 788678 341557