

UNIVERSITY OF BELGRADE FACULTY OF AGRICULTURE SERBIA

WITH A SUPPORT OF
EUROPEAN AQUACULTURE SOCIETY, EAS
AND
PANHELLENIC SOCIETY OF TECHNOLOGISTS ICHTHYOLOGISTS,
P.A.S.T.I.

**VI INTERNATIONAL CONFERENCE
“WATER & FISH”**

CONFERENCE PROCEEDINGS

**Faculty of Agriculture, Belgrade-Zemun, Serbia
June, 12 – 14. 2013.**

PUBLISHER

University of Belgrade, Faculty of Agriculture

FOR THE PUBLISHER:

Prof. Dr. Milica Petrović

EDITOR IN CHIEF

Prof. Dr. Vesna Poleksić

EDITORIAL BOARD

Prof. dr Zoran Marković, Prof. dr Vesna Poleksić, Dr Sadasivam Sashi Kaushik, Alistair Lane, Prof. dr Jana Pickova, Prof. dr Zdenek Adamek, Prof. dr Tomas Policar, Dr Fabrice Teletchea, Dr Zsigmond Jeney, Dr Tania Hubenova, Dr Aleksandar Joksimović, Doc. dr Nebojša Savić, Doc. dr Zorka Dulić, Doc. dr Ivana Živić, Doc. dr Renata Relić, Dr Aurelija Spirić, Dr Mirjana Lenhardt, Dr Svetlana Jeremić, Prof. dr Miroslav Ćirković

COMPUTER DESIGN

Nikša Tarle

TECHNICAL PREPARATION OF A COVER

Nikša Tarle

PRINTED BY

MONT PRESS, Novi Sad

NUMBER OF COPIES PRINTED

400 COPIES

ORGANIZATION OF THE CONFERENCE SUPPORTED BY:

Ministry of Education, Science and Technological Development of the Republic of Serbia

GENERAL SPONSOR

«Veterinary Institute, Subotica»

UNIVERZITET U BEOGRADU, POLJOPRIVREDNI FAKULTET,

**UZ PODRŠKU
EVROPSKOG DRUŠTVA ZA AKVAKULTURU, EAS
I
PANHELENSKOG DRUŠTVA TEHNOLOGA IHTIOLOGA**

**VI MEĐUNARODNA KONFERENCIJA
„VODA I RIBA“**

ZBORNİK PREDAVANJA

**Poljoprivredni fakultet, Beograd– Zemun, Srbija
12. – 14. jun, 2013. godine**

IZDAVAČ
Univerzitet u Beogradu, Poljoprivredni fakultet

ZA IZDAVAČA:
Prof. dr Milica Petrović

GLAVNI I ODGOVORNI UREDNIK
Prof. dr Vesna Poleksić

NAUČNI ODBOR
Prof. dr Zoran Marković, Prof. dr Vesna Poleksić, Dr Sadasivam Sashi Kaushik, Alistair Lane, Prof. dr Jana Pickova, Prof. dr Zdenek Adamek, Prof. dr Tomas Policar, Dr Fabrice Teletchea, Dr Zsigmond Jeney, Dr Tania Hubenova, Dr Aleksandar Joksimović, Doc. dr Nebojša Savić, Doc. dr Zorka Dulić, Doc. dr Ivana Živić, Doc. dr Renata Relić, Dr Aurelija Spirić, Dr Mirjana Lenhardt, Dr Svetlana Jeremić, Prof. dr Miroslav Ćirković

RAČUNARSKA OBRADA
Nikša Tarle

TEHNIČKO UREĐENJE KORICA
Nikša Tarle

ŠTAMPA
MONT PRESS, Novi Sad

TIRAŽ
400 primeraka

ODRŽAVANJE SKUPA POMOGLO:
Ministarstvo prosvete, nauke i tehnološkog razvoja Republike Srbije

GENERALNI SPONZOR
„Veterinarski zavod Subotica“

CONTENTS:

Kaushik, S.: FISH NUTRITION: IMPROVING NUTRIENT EFFICIENCY AND WATER USE	21
Lane, A.: THE EUROPEAN AQUACULTURE SOCIETY (EAS)	24
Pickova, J., Mraz, J., Zajic, T.: COMMON CARP AND AQUACULTURE ..	29
Marković, Z., Stanković, M., Dulić, Z., Rašković, B., Živić, I., Spasić, M., Vukojević, D., Relić, R., Poleksić, V.: CARP PRODUCTION IN SERVICE OF REINFORCEMENT OF SERBIA AGRICULTURE	33
Adámek, Z.: BIOMANIPULATION MEASURES ON CZECH RESERVOIRS AND THEIR PAST AND CURRENT SUCCESSFULNESS.....	39
Pucher, J., Focken, U.: INTENSIFICATION OF COMMON CARP CULTURE IN RURAL AREAS OF NORTHERN VIETNAM BY PLANT-BASED FEEDS WITH PROTEIN SOURCES OF DIFFERENT QUALITIES	44
Jovanović, R., Ivanov, D., Čolović, R., Vukmirović, Đ., Palić, D., Lević, J., Đuragić, O.: INFLUENCE OF FATTY ACID COMPOSITION OF FISH FEED ON FATTY ACID COMPOSITION OF CARP MEAT	50
Trbović, Z., Marković, Z., Petronijević, R., Milijašević, M., Spirić, D., Vranić, D., Spirić, A.: MULTIVARIATE ANALYSIS OF FATTY ACID PROFILES OF CARP MEAT DURING SEMI – INTENSIVE FARMING	56
Linhart, O., Kocour, M., Kašpar, V.: EDUCATION AND TRAINING IN AQUACULTURE AND WATER PROTECTION AT FACULTY OF FISHERIES AND PROTECTION OF WATERS, UNIVERSITY OF SOUTH BOHEMIA IN ČESKÉ BUDĚJOVICE, THE CZECH REPUBLIC	63
Poleksić, V., Dulić, Z., Stanković, M., Rašković, B., Spasić, M., Vukojević, D., Marković, Z.: HIGHER EDUCATION FOR AQUACULTURE/FISHERY AT THE FACULTY OF AGRICULTURE UNIVERSITY OF BELGRADE: IMPLEMENTATION OF THE FIRST BOLOGNA REFORM – LINKING THEORY AND PRACTICE	68
Teletchea, F., Fontaine, P.: BEYOND THE LEVELS OF DOMESTICATION IN FISH: MUST ALL FARMED SPECIES BE DOMESTICATED?	74

Żarski, D., Krejszeff, S., Palińska-Żarska, K., Kucharczyk, D.: CONTROLLED REPRODUCTION OF PIKEPERCH, SANDER LUCIOPERCA (L.) – ACHIEVEMENTS AND PROSPECTS	82
Polcar, T., Blecha, M., Kristan, J., Stejskal, V., Blaha, M.: COMBINATION OF INTENSIVE (RAS) AND EXTENSIVE (POND) AQUACULTURE FOR JUVENILE PRODUCTION IN PIKEPERCH (SANDER LUCIOPERCA).....	88
Cosson, J., Prokopchuk, G., Dzyuba, V., Fedorov, P.: FISH SPERMATOOZOA, PHYSICAL AND BIO-ENERGETIC INTERACTIONS WITH THEIR SURROUNDING MEDIA	92
Buza, E., Kolics, B., Kovács, B., Demény, F., Horváth, Á., Urbányi, B., Sándor, S., Müller, T.: ARTIFICIAL PROPAGATION AND REVEALED REPRODUCTION FEATURES OF WEATHERFISH (<i>MISGURNUS FOSSILIS</i>)	98
Mamedov, Ch., Sadiqbeyli, L., Akhundov, M., Hajiyev, R.: EVALUATION OF BIOLOGICAL INDICATORS OF YOUNG PERSIAN STURGEON (<i>ACIPENSER PERSICUS</i>) GROWN AT POND METHOD WITH DIFFERENT ARTIFICIAL FEEDS	104
Savić, N., Mikavica, D., Rogić, B.: THE GROWTH OF WEIGHT AND BODY LENGTH OF YOUNG RAINBOW TROUT (<i>ONCORHYNCHUS MYKISS</i> WAL.) ORIGINATING FROM DIFFERENT BROODSTOCK.....	111
Altinok, I., Kahraman, U., Capkin, E., Boran, H., Ozturk, R.: DEVELOPMENT OF LIVE VACCINES FROM YERSINIA RUCKERI BY aroA AND aroC GENE MUTATIONS	116
Ćirković, M., Novakov, N., Ljubojević, D., Bjelić-Čabrilo, O., Davidov, I.: FINDING OF THE NEMATODA LARVAE IN PREDATORY FISH SPECIES.....	118
Radosavljević, V., Jeremić, S., Jakić-Dimić, D.: SURVEY AND DIAGNOSTICS OF FISH DISEASES IN THE REPUBLIC OF SERBIA DURING THE PERIOD 2011-2012	124
Baltić, Ž.M., Bošković, M., Đorđević, V., Marković, R., Dimitrijević, M., Pavličević, N.: FISH-BORNE PARASITIC ZOOSES WITH SPECIAL REFERENCE TO ANTHROPOGENIC IMPACT	129
Wagenaar, G.M., Barnhoorn, I.E.J.: THE HEALTH STATUS AND EDIBILITY OF FISH FROM THREE HYPERTROPHIC IMPOUNDMENTS IN SOUTH AFRICA	136

- Ghiasi, F. Rizbandi, T.:** DETERMINATION OF VHS VIRUS TITER IN EXPERIMENTALLY INFECTED RAINBOW TROUT, *ONCORYHINCHUS MYKISS*..... 276
- Karagiannis, P., Grigorakis, K., Nathanailides, C., Fountoulaki, E., Basilaki, A., Lopez-Albors, O., Petridis, D.:** THE INFLUENCE OF DIFFERENT THERMAL CONDITIONS ON THE PROXIMATE COMPOSITION OF RAINBOW TROUT FLESH FROM TWO FISH FARMS IN NW GREECE 291
- Milošković, A., Pavlović, M., Kovačević, S., Radojković, N., Simić, S., Simić, V.:** THE PRESENCE OF ZINC IN MUSCLE TISSUE OF PRUSSIAN CARP AND BREAM IN THE GRUŽA AND BOVAN RESERVOIRS 283
- Skorić, S., Đikanović, V., Marković, G., Hegediš, A.:** CONCENTRATIONS OF 16 ELEMENTS IN TISSUES (LIVER, MUSCLE, SCALES) OF PRUSSIAN CARP (*CARASSIUS GIBELIO*, BLOCH, 1782) IN MEDJUVRŠJE RESERVOIR, SEASONAL ASPECT 288
- Obradović, S., Ivanc A., Đekić, V., Šekler, M., Živkov-Baloš, M., Veljović, N., Šarčević, B.:** CONTENT AND DISTRIBUTION OF HEAVY METALS IN ORGANS AND TISSUES OF FRESHWATER FISH IN THE ZAPADNA MORAVA RIVER 295
- Jović, M., Onjia, A., Stanković, S.:** METAL POLLUTION INDEX AS A TOOL FOR ASSESSING WATER QUALITY OF BOKA KOTORSKA BAY 300
- Ősz, Á., Kovács, B., Kánainé Sipos, D., Kaczkó, D., Herzinyák, Z., Snoj, A., Bogataj, K., Urbányi, B., Horváth, Á.:** ORIGIN AND GENETIC STRUCTURE OF SEVERAL HUNGARIAN WILD AND DOMESTICATED BROWN TROUT POPULATIONS BASED ON PCR-RFLP AND MICROSATELLITE MARKERS 305
- Muhamedagić, S., Korjenić, E., Hamzić, A., Drešković, N., Mustafčić, A., Kapo, I., Lepar, D., Habibović, E.:** ICHTYOFAUNA OF KRIVAJA RIVER CATCHMENT AREA..... 310
- Bećiraj, A., Šarić-Kadić, I., Ičanović, I., Halimović, S., Đekić, R., Galijašević, E.:** FISH BIODIVERSITY IN THE RIVER KRUŠNICA 321
- Tomljanović, T., Piria, M., Šprem, N., Matulić, D., Zanella, D.:** FINDING OF STERLET (*ACIPENSER RUTHENUS*) IN THE SAVA RIVER NEAR ZAGREB 326

- Zekić Žujo, D.:** AGE STRUCTURE OF POPULATION OF ENDEMIC SPECIES CHONDROSTOMA PHOXINUS HECKEL, 1843 (TELEOSTEI: OSTARYOPHYSI, CYPRINIDAE) FROM BUSKO LAKE 331
- Krizmanić, J., Subakov Simić, G., Predojević, D.:** ALGAE AS WATER QUALITY BIONDICATORS OF THE RIVER DJETINJA..... 342
- Bjelanović, K., Živić, I., Dulić, Z., Živić, M., Đorđević, J., Marinković, S., Marković, Z.:** WATER QUALITY ASSESSMENT IN THE RAŠKA RIVER BASED ON ZOOBENTHOS AND ZOOPLANKTON ORGANISMS AS BIOIDICTORS 349
- Dulić, Z., Grubisić, M., Stanković, M., Živić, M., Arsić, L., Marković, Z.:** EFFECTS OF DIFFERENT ORGANIC MANURES ON BODY SIZE AND PRODUCTION OF *DAPHNIA MAGNA*..... 358
- Živić, I., Bjelanović, K., Marinković, S., Jovanović, J., Marković, Z.:** PRODUCTION OF MACROZOOBENTHOS IN THE RAČA RIVER UPSTREAM AND DOWN STREAM FROM TROUT FARM ... 364
- Terzi, E., Gedik, K., Verep, B., Yandi, I.:** ASSESSMENT OF MARINE WATER QUALITY: EFFECT OF AQUACULTURE AND DOMESTIC SEWAGE DISCHARGE 372
- Isajlović, I., Vrgoč, N., Krstulović Šifner, S., Marković, O., Pešić, A.:** ASSESSMENT OF NORWAY LOBSTER - NEPHROPS NORVEGICUS (LINNAEUS, 1758) POPULATIONS IN THE ADRIATIC SEA USING ALTERNATIVE METHODS. 374
- Krstulović Šifner, S., Ikica, Z., Đurović, M., Vrgoč, N., Isajlović, I., Joksimović, A.:** DISTRIBUTION, ABUNDANCE AND POPULATION STRUCTURE OF THE COMMON CUTTLEFISH, *SEPIA OFFICINALIS* LINNAEUS, 1758, IN THE ADRIATIC SEA 380
- Lenas, S. D., Triantafyllou, J. D., Logothetis, P., Kakali, F., Nathanailides, C.:** THE INFLUENCE OF DIFFERENT ENVIRONMENTAL CONDITIONS ON FLESH QUALITY INDICES OF FARMED SEA BASS, IN NW GREECE 386
- Mandić, M., Drakulović, D., Huter, A., Petović, S., Mandić, S.:** TEHNOLOGY OF MUSSEL (*MYTILUS GALLOPROVINCIALIS*) AND OYSTER (*OSTREA EDULIS*) FARMING IN BOKA KOTORSKA BAY 388
- Barišić, J., Rathman, M., Čož-Rakovac, R., Strunjak-Perović, I., Topić Popović, N., Sauerborn-Klobučar, R., Jendriš Škrliak, E., Mihaldinec, D.:** MAGNETIC RESONANCE IMAGING - A NEW APPROACH IN INVESTIGATIONS OF ENDANGERED MARINE BIVALVES..... 393

Hajdarević, E., Hasković, E., Hercegovac, A., Adrović, A., Skenderović, I.: LEUKOCYTES OF COMMON BREAM (<i>ABRAMIS BRAMA</i> , LINNEAUS, 1758) FROM THE ARTIFICIAL MODRAC LAKE (BOSNIA & HERZEGOVINA) IN VARIOUS SEASONS.....	395
Dikanović, V., Skorić, S., Cakić, P.: REPRESENTATIVES OF TAPEWORMS (CESTODA) OF FISHES IN BELGRADE SECTION OF THE DANUBE RIVER	402
Dekić, R., Ivanc, A., Savić, N., Manojlović, M., Četković, D., Obradović, S.: HEMATOLOGICAL EVALUATION OF RAINBOW TROUT (<i>ONCORHYNCHUS MYKISS</i>) FINGERLINGS FROM DIFFERENT HATCHERIES	409
Novakov, N., Ćirković, M., Ljubojević, D., Babić, R., Aleksić, N.: PATHOLOGICAL CHANGES WHICH <i>THELOHANELLUS NIKOLSKII</i> CAUSE ON THE FINS, SCALES AND SKIN OF COMMON CARP.....	414
Capkin, E., Terzi, E., Boran, H., Altinok, İ., Başçınar, N.: BACTERIAL DISEASE SCREENING OF CULTURED HORSE MACKEREL (<i>TRACHURUS MEDITERRANEUS</i>) IN SEA-CAGES IN SOUTHERN BLACK SEA REGION OF TURKEY.....	420
Terzi, E., Capkin, E., Altinok, İ.: DISTRIBUTION OF SULFONAMIDE RESISTANCE GENES IN BACTERIA ISOLATED FROM RAINBOW TROUT IN TURKEY	422
Tiligadas, I., Moutopoulos, K. D., Chatziefstathiou, M., Tsoumani, M.M.: HELLENIC MARICULTURE SAFETY AND HEALTH: INITIAL REPORT ON WORK RELATED INJURIES AND ILL HEALTH.....	424
Aleksić-Agelidis, A., Ljubojević, D., Babić, J., Lujić, J., Novakov, N., Marković, M.: DRESSING PERCENTAGE OF 3-YEAR OLD CARP FROM CAGE PRODUCTION SYSTEM.....	429
Pavličević, N., Đorđević, V., Dimitrijević, M., Bošković, M., Marković, R., Baltić, Ž.M.: LACTIC ACID BACTERIA: EFFECT ON THE QUALITY AND SAFETY OF FISHERY PRODUCTS	436
Bianchini, L. M., Pigliarieni, E., Rambaldi, E., Argenti, L., Menesatti, P., Costa, C.: A NON INVASIVE METHOD FOR TRACING ORGANIC FISH.....	443

Zohreh, M., Ali, M.T., Amirreza, S.: THE SURVEY OF SOME BACTERIAL CONTAMINATION IN FROZEN SEMI-COOKED READY TO EAT SEA FOOD PRODUCTS IN ALBORZ PROVINCE- IRAN	452
Smederevac-Lalić, M., Zarić, V., Hegediš, A., Lenhardt, M., Mićković, B., Višnjić-Jeftić, Ž., Pucar, M., Cvijanović, G: THE MARKETING CHANNELS OF FISH CAUGHT IN LARGE SERBIAN RIVERS.....	457
Rašković, B., Vukojević, D., Spasić, M., Živić, I., Stanković, M., Marković, Z., Poleksić, Z.: EFFECT OF STRESS ON PRESENCE OF DEVELOPING NEPHRONS OF COMMON CARP REARED IN SEMIINTENSIVE SYSTEM.....	463
Grubišić, M., Dulić, Z., Đorđević, J., Bjelanović, K., Spasojević, I., Relić, R., Marković, Z.: SEASONAL VARIATIONS AND DIVERSITY OF CLADOCERA (CRUSTACEA) IN NATURAL, DOMESTIC WASTEWATER TREATMENT LAGOONS.....	467
Spasojević, I., Dulić, Z., Grubišić, M., Đorđević, J., Ljubobratović, U., Raičević, V., Marković, Z.: TOTAL COLIFORM AND FECAL COLIFORM COMMUNITY DYNAMICS IN NEWLY BUILT WASTEWATER TREATMENT LAGOONS.....	473
Relić, R., Lakić, N., Mladenović, V., Dulić, Z., Grubišić, M., Poleksić, V.: FISH WELFARE FROM PERSPECTIVE OF CONSUMERS IN SERBIA (PRELIMINARY RESULTS)	479
Amiri, M, Hosseini, H., Islami, R. H.: IDENTIFICATION AND DISTRIBUTION OF SEA CUCUMBERS IN THE CHABAHAH BAY, IRAN.....	487

DRESSING PERCENTAGE OF 3-YEAR OLD CARP FROM CAGE PRODUCTION SYSTEM

ALEKSANDRA ALEKSIĆ-AGELIDIS¹, DRAGANA LJUBOJEVIĆ², JELENA BABIĆ³, JELENA LUJIĆ⁴, NIKOLINA NOVAKOV³, MAJA MARKOVIĆ¹

¹Faculty of Veterinary Medicine, Bulevar oslobođenja 18, 11 000 Belgrade, Serbia

^{2,5}University of Novi Sad, Faculty of Agriculture, Trg Dositeja Obradovića 8, 21000 Novi Sad, Serbia

³Institute of Meat Hygiene and Technology, Kačanskog 13, 11 000 Belgrade, Serbia

⁴University of Novi Sad, Faculty of Sciences, Trg Dositeja Obradovića 3, 21000 Novi Sad, Serbia

RANDMAN TROGODIŠNJEG ŠARANA IZ KAVEZNOG SISTEMA UZGOJA

Apstrakt

Uzgoj šarana (*Cyprinus carpio* L.) u kaveznom sistemu predstavlja posebnu vrstu intenzivne proizvodnje. Odlikuje se malim početnim ulaganjima u izgradnju kaveznog ribnjaka, ne zahteva značajno angažovanje radne snage i obezbeđuje veliku proizvodnju po jedinici zapremine.

Izbor lokacije je od velikog značaja jer utiče na ekonomsku održivost, proizvodnju i mortalitet šarana. U odnosu na druge uzgojne sisteme, utvrđeno je da u kaveznom sistemu morbiditet i mortalitet mogu značajno varirati i u slučaju odstupanja vrednosti parametara kvaliteta vode od optimalnih, gubici mogu nastupiti brzo i biti dramatični.

Nedostatak kaveznog sistema gajenja ogleda se u olakšanoj transmisiji bolesti i zagađenju vodenog ekosistema u kome je kavezni sistem postavljen, što se može sprečiti uvođenjem zaštitnih sistema. Uz poštovanje načela dobre proizvođačke prakse, održavanjem dobrog kvaliteta vode i optimalno izbalansiranu hranu primerenu starosnoj kategoriji uzgajanog šarana, očekivan mortalitet je 1-5%.

Zahtevi savremenog tržišta su sve više usmereni ka obrađenoj ribi, posebno filetima. Uklanjanjem kože i odstranjivanjem unutrašnjih organa i intramuskularnih kostiju, fileti šarana i ostalih ciprinida koje se gaje u polikulturi sa šaranom, postaju visokovredni obroci, koji se lako i brzo spremaju. Randman riba je ekonomski parametar čija je vrednost značajna pri svim tehnološkim operacijama vezanim za preradu ribe. Iskazuje se kao odnos primarno obrađenog trupa u odnosu na masu žive ribe.

Odnos jestivog i nejestivog dela trupa ribe može značajno da varira u zavisnosti od vrste, mase i veličine ribe i sezone ulova). Od njega direktno zavisi ekonomičnost proizvodnje i neopravdano je zanemaren kao parametar u istraživanjima.

Razlika u randmanu između različitih linija i njihovih hibrida javlja se zbog različitog oblika, dužine glave, debljine i širine tela. Randman riba značajno je povoljniji od randmana kod ostalih vrsta životinja. Delovi koji se odbacuju mogu sadržati sve ili samo neke delove glave, krljušti, kože, creva, gonada i peraja.

Cilj istraživanja bio je određivanje randmana kod trogodišnjeg šarana poreklom iz kaveznog sistema uzgoja. Uzorkovanje je obavljeno iz dva kavezna sistema na Tikveškom jezeru (Makedonija) (41°20'51"N 21°57'58"E) i iz jednog na Bilečkom jezeru (Bosna i Hercegovina) (42°49'31"N 18°26'17"E). Ishrana riba vršena je kompletnim krmnim smešama različitih proizvođača. Iz svakog kaveznog sistema uzorkovano je po 8 jedinki muškog i 8 jedinki ženskog pola (ukupno po 24 jedinke svakog pola).

Morfometrijske osobine riba utvrđene su sa ciljem da se ispita uticaj pola na randman trogodišnjeg šarana gajenog u kaveznom sistemu. Nakon što su izlovljeni, šarani su držani na ledu. Krljušt je skinuta ručno nazubljenim nožem. Glava je odsečena cirkularnim rezom ispred pojasa pektoralnog peraja tako da je peraje ostalo na trupu. Peraja su odsečena na početku perajnih žbica. Egzenteracija organa zajedno sa gonadama obavljena je ručno. Dobijeni su obrađen trup, koji podrazumeva trup ribe bez krljušti, peraja, unutrašnjih organa i glave. Posle odstranjivanja krljušti, unutrašnji oragni, gonade, glava, peraja i obrađen trup su izmereni. Rezultati su obrađeni t-testom kako bi se utvrdio uticaj pola na randman trogodišnjeg šarana gajenog u kaveznom sistemu proizvodnje.

U odnosu na pol (ženke naspram mužjaka šarana) pažnja je usmerena na parametre koji su važni sa aspekta gajenja i što povoljnijeg randmana.

U našem istraživanju randman je određen kao odnos mase cele ribe i mase trupa bez glave, krljušti, peraja i unutrašnjih organa. Rezultati dobijeni u našem istraživanju pokazuju veću težinu ženskih u odnosu na muške jedinke (FW), ali nemaju statističku značajnost.

Ukupna dužina (TL), standardna dužina (SL) i dužina trupa (CL) ženki u odnosu na mužjake imaju statistički značajnu razliku ($p < 0,01$), dok razlike u telesnoj masi ženki i mužjaka nisu statistički značajne. Rezultati dobijeni merenjem težine trupa (CW) i težine unutrašnjih organa (VW) ženki u odnosu na iste parametre kod mužjaka imaju statistički značajnu razliku ($p < 0,01$). Vrednosti dobijene merenjem težine glave (HW), težine gonada (GW) mužjaka, kao i vrednosti gonadosomatskog indeksa (GSI) u odnosu na vrednosti istih parametara kod ženki imaju statistički značaj ($p < 0,01$). Razlike u dužini glave (HL), visini tela (BH), masi peraja (FW), masi jedinki (FW) i obrađenih fileta (FilletW) mužjaka i ženki, nemaju statistički značaj. Vrednosti dobijene izračunavanjem relativne težine fileta i randmana uzorkovanih trogodišnjih jedinki šarana muškog i ženskog pola nemaju statistički značajnu razliku.

Ključne reči: Šaran, kavezni sistem, pol, randman
Keywords: carp, female, male, dressing percentage

INTRODUCTION

Carp (*Cyprinus carpio* L.) is domesticated in every continent except Antarctica, owing to its ability in adapting to different values of water quality parameters, including tolerance to very low dissolved oxygen concentrations. Considering the benefits that are reflected by the extensive reproductive abilities, breeding and prime selection potentials, and high resistance to various diseases, it is not surprising that carp represents

the dominant species in all counties with prerequisites for extensive, semi-intensive and intensive freshwater fish production. The carp growth in different production systems depends on many factors, such as varietal, density, welfare and fish feed with proper nutrients along with geographic location.

Cage production system for carp is a special type of intensive production. It is characterized by lower initial investments in cage construction and minimal labor engagement, ensuring high production yield per volume capacity of water contained in the constructed units. Modern cages are constructed from metal and synthetic polymers (Mihailovic et al., 2007). Cage has to be robust enough to endure associated hazards such as winds and waves, uncomplicated servicing and maintenance. Location choice is critical as it affects the production and mortality of the produce (Carp) directly impacting the profitability of the production unit.

Below listed are three-selection criterion considered for selection of sites for cage culture (Bogut et al., 2007).

1. Physical and chemical parameters including temperature, dissolved oxygen concentration, water flow, pollution and algal blooms.
2. Factors relating to the choice of location taking into account weather conditions, shelter, substrate, water currents and the degree of blur. Additionally the shape and size of the cage and the depth to which they are installed.
3. Profitability of the farm, and includes legal aspects, accessibility, facilities, security, economic and social conditions.

Superior production practices, quality of water and optimally balanced diet appropriate for the cultured carp category, expected mortality rate is 1-5%. Compared to other breeding systems, it was found that the cage system morbidity and mortality could vary significantly in the case of deviation in the values of water quality parameters from the optimum, the losses can occur quickly and be dramatic. (Orajić et al., 2007)

Shortcoming of cage production system is reflected through allowing of disease transmission and aquatic ecosystems contamination where the cage system is set up, this can be prevented by protective systems introduction.

Modern market demands are increasingly directed towards processed fish, especially fillets (Hough, 1993; Vallod, 1995). By removing the skin, internal organs and intramuscular bones, fillets of carp and other cyprinids grown in polyculture along with the carp, become high value meals that are easily and quickly prepared (Lin et al., 1989).

Dressing percentage of fish is an economic parameter with significant value attached to it, in all technological operations related to fish processing. It is expressed as the ratio of the primary processed carcass (dressed) relative to carcass weight of live fish.

The ratio of edible and non-edible part of the fish carcass can vary significantly depending on the species, weight, size of the fish and season catch (Baltic and Teodorović, 1997). Since the dressing percentage directly influences the economy (profitability) of production, it is unjustly neglected as the parameter in scientific research/studies (Lju-bojević et al, 2012).

The difference in yield of different lines and their hybrids occur due to different body shape, head length, width and fat of the body. Dressing percentage of fish is significantly higher comparatively than the yield of other animal species (Ćirković et al., 2002). Parts that are rejected may include all or some parts of the head, scales, skin, intestine, gonads and fins.

According to Dunhamu et al. (1983) fish carcass primary handling involves removal of the head, scales and internal organs, while Lovell (1981) reference states that primary

treatment involves the removal of skin, dorsal and pectoral fins, head and abdominal organs cavity, with the tail present. Naumovski (1991) and Tumbas (1976) define fish carcass primary handling as a process which include removal of the tail fin.

In this study, yield was determined as the ratio of the fish weight and the carcass without the head, scales, fins and internal organs.

MATERIALS AND METHODS

The specimens of 3-year old carp from cage production system were sampled in the Tikveš lake in Macedonia (41°20'51"N 21°57'58"E) (two groups) and from the Bilećko Lake in Bosnia and Herzegovina (42°49'31"N 18°26'17"E) (one group). Fish were fed with a completed feed mixture from different manufacturers. From every cage system, 8 male and 8 female samples are taken (24 individuals of each sex). For the statistical analysis of the results, software package Microsoft Office Excel 2010 and Data Analysis ToolPack, were used. Statistical t-test was used for mean values determination. Morphometric characteristics determination, (described by Baltić and Teodorović, 1997) of 3-year old carp specimens from cage production system is conducted in order to evaluate the dressing percentage in relation to sex. After being harvested, the carp were kept on ice. The scales are removed with a serrated knife. The head is cut off with a circular incision in front of the pectoral fins (Gela and Linhart, 2000). The internal organs along with the gonads were removed manually and a processed carcass (without scales, fins, internal organs and head) is obtained. After dissection, the internal organs, gonads, head, fins and carcass were measured. The obtained results were used to determine the relative percentage ratio in relation to the initially measured fish body weight.

RESULTS

The biometric parameters of 3-year old male and female carp specimen are presented in the Table 1: (TL-total length, SL-standard length, CL-body length, HL-head length and BH-body height), and weight parameters (FW-fish weight, CW-carcass weight, FW- fillet weight, HW- head weight, FW-fin weight, GW-gonad weight, VW- viscera weight). After dissection acquired data were used to calculate the percentage of fillets, head, fins, viscera and waste per fish weight, as well as important indices such as dressing percentage (DP), gonadosomatic index (GSI) and relative fillet weight. The obtained results were used to determine the relative percentage ratio of each parameter in relation to the initially measured fish body weight. For the statistical analysis of the results, software package Microsoft Office Excel 2010 and Data Analysis ToolPack, were used. Statistical t-test was used for mean values determination.

Table 1. The values of biometric parameters in carp (*Cyprinus carpio*) in relation to sex (females, males) Values in the table are the mean values \pm SD (n = 24), values in the same row with different letters in superscript differ significantly at $p < 0,01$.

Parameter	Unit	Female n=24 mean \pm S.D.	Male n=24 Mean \pm S.D.
Biometric parameters			
Total length (TL)	mm	457,5 \pm 13,72 ^a	438 \pm 8,18 ^b
Standard length (SL)	mm	393,88 \pm 16,76 ^a	369,87 \pm 21,83 ^b
Carcass length (CL)	mm	322,5 \pm 13,72 ^a	308 \pm 8,18 ^b
Head length (HL)	mm	87,92 \pm 2,61	87,7 \pm 2,8
Body height (BH)	mm	126 \pm 4,8	125,54 \pm 6,51
Weight parameters			
Fish weight (FW)	g	1550,23 \pm 55,59	1495,47 \pm 109,09
Carcass weight (CW)	g	1010,12 \pm 19,57 ^a	947 \pm 5 \pm 72,68 ^b
Fillet weight (filletw)	g	750,42 \pm 33,81	727,5 \pm 51,29
Head weight (HW)	g	235,96 \pm 30,98 ^b	263,89 \pm 27,35 ^a
Fins weight (FW)	g	40,05 \pm 4,59	38,88 \pm 5,4
Viscera weight (VW)	g	267,18 \pm 57,13 ^a	222,14 \pm 24,64 ^b
Gonad weight (GW)	g	56,57 \pm 3,21 ^b	59,61 \pm 2,89 ^a
Calculated parameters			
GSI		3,66 \pm 0,26 ^b	4,01 \pm 0,4 ^a
Dressing percentage	%	65,57 \pm 2,25	63,57 \pm 5,49
Fillet weight relative	%	48,41 \pm 1,29	48,6 \pm 2,02

The results obtained in our study showed slightly higher body height (BH) and head length (HL) values in the females compared to the males. Weight parameters obtained (fish weight, fin weight and fillet weight) had no statistically significant difference ($p > 0.05$) Total length (TL), standard length (SL), and carcass length (CL) of females compared to males had a statistically significant difference ($p < 0.01$). In female carp, significantly higher carcass weight (CW) and viscera weight (VW) were found ($p < 0.01$). Significant differences ($p < 0.01$) were found in head weight (HW) and gonad weight (GW) as well in gonadosomatic index value (GSI) in the favor of male carps.

DISCUSSION

Carp (*Cyprinus carpio* L.) belongs to the species that have expressed a high degree of sexual dimorphism. The results of research conducted by Bay et al. (2006) showed almost the same growth rate of male and female 3-year carp from four different sub-groups examined. The faster growth of female carp was confirmed by research of Cherfasa et al. (1996). Analysis of the relation between sexual maturity and growth rate of the Chinese and European carp conducted by Hulata et al. (1995) showed that the beginning of sexual maturity has a negative impact on the physical growth of many species of fish. The production of sex hormones during sexual maturation, especially their accumula-

tion, inhibits growth as a result of the reaction between sex hormones and growth hormones. Due to the different time of reaching sexual maturity amongst different sexes, the results of this study are consistent with the conclusion that males attribute of reaching sexual maturity earlier than females, utilize more nutrients for development of the gonads, while sexually immature female specimen(s) during the maturation period go through intensive physical growth.

CONCLUSION

The results of this study demonstrate greater weighing females compared to males (FW), but of no statistical significance.

The results gonad weight (GW) of males and values of the gonad somatic index (GSI) of males had a statistically significant difference compared to the values of the same parameters in females ($p < 0.01$). Total length (TL), standard length (SL) and body length (CL) of females compared to males had a statistically significant difference ($p < 0.01$), while the difference in weight of females and males were statistically of little significance. Results obtained by measurements of carcass weight (CW) and the weight of the internal organs (VW) females compared to males with the same parameters demonstrated statistically significant difference ($p < 0.01$). The difference in values obtained by measuring the weight of the head (HW), gonad weight (GW) of males, and gonad somatic index GSI values compared to the values of the same parameters in females demonstrated variance of statistical significance ($p < 0.01$).

Analytical Differences in the head length (HL), body height (BH), body weight of individuals (FW) and fin weight (FW) of males and females demonstrated variance of no statistical significance.

Values obtained by calculating the relative weights and fillet yield of three-year sampled carp male and female demonstrated no variance of statistical difference. Production of 3-year old carp in cage production system is economically unprofitable compared to the production of a 2-year old carp, due to utilization of more nutrients and energy received in feed during sexual maturation for development of the gonads.

ACKNOWLEDGEMENT

The presented results stemmed from the work on the project no. TR31011 "Influence of dietary components in cyprinids meat quality, cost of production, and economic losses" research program in the field of technological development funded by the Ministry of Education and Science, Republic of Serbia.

REFERENCES

Baltić, M., Teodorović, V. (1997): Higijena mesa riba, rakova i školjki. Veterinarski fakultet. Beograd.

Buchtova, H., Svobodova, Z., Kocour, M., Velišek, J. (2006): Evaluation of the Dressing Percentage of 3-year-old Experimental Scaly Crossbreds of the Common Carp (*Cyprinus carpio*, Linnaeus 1758) in Relation to Sex. Acta Vet., Brno, 75:123-132.

Buchtova, H., Svobodova, Z., Kocour, M., Velišek, J. (2006): Evaluation of growth and dressing out parameters of experimental scaly crossbreds in 3-year-old common carp (*Cyprinus carpio*, Linnaeus 1758). Aquaculture Research 37, 466-471 pp.

Bogut, I., Magovac, R., Sabo, D., Bodakoš, D., Galović, D., Arežina, M., Rajković, V. (2007): Cage fattening results of Common Carp (*Cyprinus carpio*) in hydroaccumulation Grabovo near Vukovar. *Krmiva* 49, Zagreb, 4, 207-214 pp.

Ćirković, M., Trbović, D., Ljubojević, D., Đorđević, V. (2011): Meat quality of fish farmed in polyculture in carp ponds in Republic of Serbia. *Tehnologija mesa*, vol. 52, br. 1, str. 106-121.

Gela, D., Linhart, O. (2000): Evaluation of slaughtering value of common carp from diallel crossing. *Czech Journal of Animal Science* 45, 53-58 pp.

Gross, R. (1997): Dressing percentage in market-size Common Carp: Effect of strain, year-class, sex, body size and shape. *Aquaculture sponsored symposium on the Carp*. Budapest, September 6-9, poster.

Ljubojević, D., Ćirković, M., Novakov, N., Babić, J., Lujić, J., Marković, T. (2012): Faktori koji utiču na randman šaranskih riba. *Tehnologija mesa*, vol. 53, br 1, str. 14-19.

Orajić, D., Zrnčić, S. (2009): Bolesti šarana-opasnosti u kaveznom uzgoju, Uzgoj slatkovodne ribe, stanje i perspective. *Hrvatska gospodarska komora. Zbornik radova*, 87-94 pp.

Lovell, R.T., Menghe, Li. (1992): Comparison of Feed Conversion, Dressing Yield and Muscle Composition for Second-and Third-Year Chanell Catfish. *The Progressive Fish Culturist* 54, 171-173 pp.

Ljubojević, D., Ćirković, M., Novakov, N., Babić, J., Lujić, J., Marković, T., (2012): Uticaj različitih čimbenika na randman komercijalno uzgajanih ciprinidnih vrsta riba na šaranskim ribnjacima u Srbiji. *Croatian Journal of Fisheries* 70, 89-97 pp.

Mihailović, P., Marković, Z. (2007): Projektovanje kaveznih sistema za gajenje riba. III Međunarodna konferencija Ribarstvo. *Zbornik radova*, 317-324 pp.

Safner, R., Treer, T., Aničić, I., Kolak, A. (2001): Dressing percentage of four croatian common carp (*Cyprinus carpio* L.) populations. *Ribarstvo* 59 (4), 131-141pp.

Safner, R., Treer, T., Aničić, I., Kolak, A., Lovrinov, M., (1997): The Assessment of Phenotypic Correlation for the Common Carp (*Cyprinus carpio*). *Dressing Percentage Agriculturae conspectus scientificus*, vol. 63, No 1-2, 79-86 pp.

Vallois D., (1995): Carp processing and market analysis: a case study in France. *Aquaculture*, 129, 475-478 pp.

CIP - Каталогизација у публикацији
Народна библиотека Србије, Београд

639.2/.3(082)

INTERNATIONAL Conference „Water & Fish“ (6 ;
2013 ; Beograd)

Conference Proceedings / VI International
Conference „Water & Fish“, June, 12-14.2013.
; [editor in chief Vesna Poleksić]. -
Belgrade : Faculty of Agriculture, 2013 (Novi
Sad : Mont Press). - 455 str. : ilustr. ; 30
cm

Na spor. nasl. str.: Zbornik predavanja. -
Tiraž 400. - Bibliografija uz svaki rad.

ISBN 978-86-7834-155-7

а) Рибарство - Зборници
COBISS.SR-ID 198755596

ISBN 978-86-7834-155-7



9 788678 341557