



International Journal of Botany

ISSN: 1811-9700

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Bryophyte Diversity Within Urban Areas: Case Study of the City of Belgrade (Serbia)

¹M. Sabovljević and ²S. Grdović

¹Institute of Botany, Faculty of Biology, University of Belgrade, Takovska 43, 11000, Belgrade, Serbia

²Faculty of Veterinary Medicine, University of Belgrade, Bulevar Oslobođenja 18, 11000, Belgrade, Serbia

Abstract: The bryophyte diversity and urban bryophyte flora of the city of Belgrade was studied. In total 210 taxa were recorded, with 23 hepatics and 187 mosses. Comparing to some other urban areas of Europe, bryophyte diversity of the Belgrade metropolitan belongs among the richest in Europe, however bryophyte are not abundant. Among bryophytes of Belgrade city, there are 14 nationally red-listed species and two internationally threatened. According to the frequency of records, the Belgrade bryophytes are classified to rare, common and spread. Rare species within the urban area are 94, 69 are common to find and only 24 are spread and easy to record in Belgrade wide area. Urban metropolitan areas are different from native but gives various condition in small shelters for rich diversity of small organisms like bryophytes.

Key words: Diversity, bryophytes, mosses, hepatics, urban flora, Belgrade

INTRODUCTION

Urban areas comprise many ecological situations which differs among each other. Considering the small size of the microhabitats, it can not offer potentially many resources for the proper development of huge plants. However, small plants like bryophytes i.e., liverworts and mosses are widely adapted to such sites and even more appreciate the absence of other competitive plants. Therefore, bryophyte flora of urban areas represent significant part of the city vegetation.

The urban bryophyte vegetation is interesting not only by its floristic composition, but to presence of rare and significant taxa, vegetation relationships and from phytogeographical point of view.

The city of Belgrade (serb. Beograd) is the capital of Serbia having about 2 million inhabitants (ca. 21% of the Serbian population). It is located in the south-eastern of Europe, where, the Pannonian plain meets the Balkan Peninsula, at the confluence of the Sava and Danube rivers ($44^{\circ}49'14''$ North, $20^{\circ}27'44''$ East). Belgrade lies 116.75 m (383 ft) above sea level. The city has an urban area of 360 km² (139 sq miles), while together with its metropolitan area it covers 3,223 km² (1,244.4 sq miles). Central Belgrade has hilly terrain, while the highest point of Belgrade proper is Torlak Hill at 303 m (994 ft). The mountains of Avala (511 m (1,677 ft)) and Kosmaj (628 m (2,060 ft)) lie south of the city. Across the Sava and Danube, the land is mostly flat, consisting of alluvial plains and loess cliffs and plateaus.

It is one of the oldest cities in Europe and since ancient times it has been an important traffic focal point,

an intersection of the roads of Eastern and Western Europe. Its territory is divided into 17 municipalities and covers 3.6% of the territory of Serbia.

Belgrade has a moderate continental climate with a year-round average temperature of 11.7°C (53.1°F) and four well defined seasons.

The hottest month is July, with an average temperature of +22.1°C (71.8°F). There are, on average, 31 days a year when the temperature is above 30°C and 95 days when the temperature is above 25°C. Belgrade receives about 700 mm (27.56 inches) of precipitation a year. The average annual number of sunny hours is 2,096. The sunniest months are July and August, with an average of about 10 sunny hours a day, while December and January are the gloomiest, with an average of 2-2.3 sunny hours a day. The highest ever recorded temperature in Belgrade was +43.1°C, while on the other end, the lowest temperature was -21°C. Mean atmospheric pressure in Belgrade is 1,001 millibars and mean relative humidity is 69.5%.

Autumn is longer than spring, with longer sunny and warm periods - the so-called Indian summer. Winter is not so severe, with an average of 21 days with temperature below zero. January is the coldest month, with average temperature of 0.4°C. Spring is short and rainy. Summer arrives abruptly and is hot and dry (Jovanović, 1994) (Fig. 1).

Hydrological network of Belgrade is well developed. Geology is mainly limestone, but loess, sandstone and even some serpentine fragments can be found and substrate over are very diverse.




Fig. 1: Climate diagram of Belgrade (Beograd)

Earlier, there are a few sporadic bryophyte records from the wide area of Belgrade (Soška, 1949; Pavletić, 1955). Grdović and Stevanović (2006) gave contribution to the urban bryoflora of downtown area in Belgrade citing 58 species.

In this study and extensive list of species recorded in wide city area of Belgrade is given as well as the urban bryophyte ecology. The study represent the first extensive investigation of the urban bryophytes within South Eastern Europe.

Up to date studies on bryophytes of urban areas are known for only a few huge urban areas e.g., Berlin and Brandenburg (Schaepe, 1986; Benkert *et al.*, 1995), Brussels (Vanderpoorten, 1997), Wien (Hohenwallner and Zechmeister, 2001a, b; Zechmeister *et al.*, 2001; Hohenwallner, 2000a, b).

Among Southern European urban areas an extensive data on urban bryophyte floras of Western Mediterranean towns in Spain (Ballesteros Segura and Ron, 1985; Casas and Saiz, 1982; Esteve *et al.*, 1977; Fiol, 1983; Heras and Soria, 1990; Lara and Mazimpaka, 1989; Lara *et al.*, 1991; Mazimpaka *et al.*, 1988, 1993; Soria and Ron, 1990, 1995; Soria *et al.*, 1992; Vicente *et al.*, 1986; Viera and Ron, 1986), Portugal (Bento-Pereira and Sérgio, 1983; Sérgio, 1981; Sérgio and Bento-Pereira, 1981) and Italy (Carcano, 2004, 1989; Cortini Pedrotti, 1989; Aleffi, 1991; Aleffi and Taruschio, 1996; Dia and Not, 1991; Lo Giudice, 1992; Giudice *et al.*, 1997; Pokorny *et al.* 2006) are well documented.

A few, but very important studies on urban bryophyte ecology give insights into the biological peculiarities and significance of these plants in urban environment (Gilbert, 1968, 1970, 1989; Nickl-Navratil, 1960; Vareschi, 1936; Brandes, 1983; Franzen, 2001; Humer-Hochwimmer and Zechmeister, 2001; Solga *et al.*, 2005, 2006 a, b; Solga and Frahm, 2006; Sabovljević *et al.*,

2005, 2007; Vukojević *et al.*, 2005, 2006; Frahm and Sabovljević, 2007).

MATERIALS AND METHODS

During 2000-2007 an extensive bryophyte collection were made in the metropolitan region of the Belgrade city, which includes territories of all municipalities of Belgrade (Stari Grad, Vracař, Savski Venac, Zvezdara, Cukarica, Palilula, Novi Beograd, Zemun, Surčin, Voždovac, Rakovica, Obrenovac, Grocka and partly (northern areas of) Barajevo, Sopot and Mladenovac). A random sampling in different habitats and microhabitats within the urban area including larger green surface were used to collect as much as possible data on urban bryophyte flora and discover the its diversity within the city of Belgrade.

Nomenclature follows Sabovljević and Natcheva (2006) and Ros *et al.* (2007) for hepaticas and Hill *et al.* (2006) for mosses. The exception are made within *Hypnum cupressiforme* complex and *Syntrichia ruraliformis* (following Sabovljević *et al.*, 2008).

All the bryophyte collected are assigned to one of four main habitat types (following classification of habitat types proposed by Ron *et al.* (1987): (1) reophilic (species adapted to water flow and inhabited water proximity); (2) rupestrian and chasmophytic (species growing on concrete and mortar, tiles and bricks, wet, shaded, sunny and dry limestone or acidic rocks and walls); (3) terrestrial (sunny, dry, humid and nitrified soils); (4) epiphytic (tree bases, trunks). The city is for the purpose of easier comparison divided to five zones: (A) old city; (B) modern city; (C) inner parks and green surfaces; (D) outer parks, gardens and yards and (E) the rivers Danube and Sava sides with attachments through the city. The bryophytes collected are classified as: R) Rare (one or two record(s)); K) common (3-7 records) and S) Spread (more than 7 records, within studied area).

For the comparison with bryoflora richness of other cities we randomly chose some western, central and south European cities: Cologne (Sabovljević and Sabovljević, in press) and Rolandseck (Frahm, 2006) in Germany; Trento in Italy (Pokorny *et al.*, 2006), Vienna in Austria (Hohenwallner and Zechmeister, 2001a, b), Madrid (Mazimpaka *et al.*, 1988), Granada (Esteve *et al.*, 1977) and Murcia (Rams *et al.*, 2000) in Spain, Szczecin (Fudali, 1994) in Poland and Bruxelles (Vanderpoorten, 1997) in Belgium.

Bryophyte catalogue: For the each taxa found in the city of Belgrade (wide area), information of the city zones, they lived and habitat, they were growing, are given. Hepaticas and mosses are listed separately and alphabetically. Specimens are deposited in Belgrade University Herbarium (BEOU) and the herbarium of Faculty of Veterinary Medicine.

RESULTS

The bryophyte flora of the Belgrade city counts 210 species of which 187 belongs to mosses and 23 to hepatics. They are listed below (see the chapter Material and Methods for the abbreviations):

Belgrade bryophyte species

Hepatics

1. *Calypogeia azurea* Stotler et Crotz. D4 - (R)
2. *Chiloscyphus polyanthos* (L.) Corda D4 - (R).
3. *Cololejeunea rossettiana* (C. Massal.) Schiffn. - D2, D3 - (R)
4. *Conocephalum conicum* (L.) Dumort. B2, C2, C3, D1 - (K)
5. *Fossombronia wondraczekii* (Corda) Lindb. D3 - (R)
6. *Frillania dilatata* (L.) Dumort. D4 - (K)
7. *Jungermannia* sp. - D3 - (R)
8. *Lejeunea cavifolia* (Ehrh.) Lindb. D4 - (R)
9. *Lophocolea bidentata* (L.) Dumort. D4 - (K)
10. *Lophocolea heterophylla* (Schrad.) Dumort. D4 - (K)
11. *Lunularia cruciata* (L.) Lindb. A3, B3 - (R)
12. *Marchantia polymorpha* L. D1 - (K)
13. *Metzgeria furcata* (L.) Dumort. D2, D4 - (K)
14. *Pellia endiviifolia* (Dicks.) Dumort. - D1, D3 - (K)
15. *Pellia epiphylla* (L.) Corda D4 - (K)
16. *Plagiochila porelloides* (Torrey ex Nees) Lindenb. D3 - (K)
17. *Porella platyphylla* (L.) Pfeiff. D2, D4 - (K)
18. *Radula complanata* (L.) Dumort. D4 - (K)
19. *Radula lindbergiana* Gottsche ex C. Hartm. D4 - (R)
20. *Riccia fluitans* L. D1, D3 - (R)
21. *Riccia sorocarpa* Bisch. D3 - (R)
22. *Ricciocarpus natans* (L.) Corda - E1 (R)
23. *Scapania nemorea* (L.) Grolle - D3 - (R)

Mosses

1. *Aloina aloides* (Koch ex Schultz) Kindb. A3, B3 - (K)
2. *Aloina ambigua* (Bruch and Schimp.) Limpr. A3, B3 - (K)
3. *Aloina rigida* (Hedw.) Limpr. A3, B3 - (K)
4. *Amblystegium confervoides* (Brid.) Schimp. D2, D4, E4 - (K)
5. *Amblystegium serpens* (Hedw.) Schimp. A1, A2, A3, A4, B3, C2, C3, C4, D1, D2, D3, D4, E2, E4 - (S)
6. *Amblystegium subtile* (Hedw.) Schimp. C4, D2, D3, D4, E4 - (S)
7. *Anomodon attenuatus* (Hedw.) Huebener D1, D4 - (R)
8. *Anomodon longifolius* (Schleich. ex Brid.) Hartm. C4 - (R)

9. *Anomodon viticulosus* (Hedw.) Hook. and Taylor D3, D4 - (K)
10. *Atrichum undulatum* (Hedw.) P. Beauv. D3 - (K)
11. *Barbula convoluta* Hedw. B2, C2, D2, D4, E2 - (K)
12. *Barbula unguiculata* Hedw. A2, A3, C2, D2, D4, E2 - (K)
13. *Brachytheciastrum velutinum* (Hedw.) Ignatov and Huttunen A2, A3, C2, D1, D3, D4 - (K)
14. *Brachythecium albicans* (Hedw.) Schimp. D3 - (R)
15. *Brachythecium campestre* (Mull.Hall.) Schimp. D2 - (R)
16. *Brachythecium glareosum* (Bruch ex Spruce) Schimp. D2, D3, D4 - (K)
17. *Brachythecium rivulare* Schimp. C2, D2, D3 - (R)
18. *Brachythecium rutabulum* (Hedw.) Schimp. A2, A3, B3, C2, C3, D2, D3, D4 - (S)
19. *Brachythecium salebrosum* (Hoffm. Ex F. Weber and D. Mohr) Schimp. A3, B3, C2, D2, D4 - (S)
20. *Bryoerythrophyllum recurvirostre* (Hedw.) P. C. Chen A3, B3, C3 - (K)
21. *Bryum algovicum* Sendtn.ex Mull.Hal. C2, D2 - (R)
22. *Bryum archangelicum* Bruch and Schimp. E2 - (R)
23. *Bryum argenteum* Hedw. A2, A3, A4, B2, C2, C3, C4, D2, D4, E2 - (S)
24. *Bryum atrovirens* agg. A3, C2 - (K)
25. *Bryum caespiticium* Hedw. A2, A3, C2, D2, D3, E2, E4 - (S)
26. *Bryum caespiticium* Hedw. aff. *Bryum badium* (Brid.) Schimp. A2 - (R)
27. *Bryum capillare* Hedw. A2, B3, C2, C4, D2, D4, E2 - (S)
28. *Bryum creberrimum* Taylor C3, D2, E4 - (R)
29. *Bryum dichotomum* Hedw. A2, A3, C2, D2, E2, E4 (K)
30. *Bryum intermedium* (Brid.) Blandow - B2, C2, C3, D2 - (K)
31. *Bryum kunzei* Hornsch. A2, A3 - (K)
32. *Bryum moravicum* Podp. D4 - (K)
33. *Bryum pallens* Sw. B2, D2, D4, E2 - (K)
34. *Bryum palescens* Schleicht. ex Schwagr. B2, C2, D2, D4 - (K)
35. *Bryum pseudotriquetrum* (Hedw.) P. Gaertn. et al. D2, E2, E4 - (K)
36. *Bryum subapiculatum* Hampe C3 - (R)
37. *Bryum turbinatum* (Hedw.) Turner D2 - (R)
38. *Callicladium haldanianum* (Grew.) H. A. Crum D4 - (R)
39. *Calliergonella cuspidata* (Hedw.) Loeske D3 - (R)
40. *Campyliadelphus chrysophyllus* (Brid.) R. S. Chopra D3, E4 - (R)
41. *Campylophyllum calcareum* (Crundw. and Nyholm) Hedenas D4 - (R)
42. *Ceratodon purpureus* (Hedw.) Brid. A2, A3, A4, C2, C3, C4, D2, D3, E2 - (S)
43. *Cirriphyllum piliferum* (Hedw.) Grout. C2, D3 - (R)

44. *Cratoneuron filicinum* (Hedw.) Spruce D1 - (R)
 45. *Crossidium crassinerve* (De Not.) Jur. A3, D3 - (R)
 46. *Crossidium laxofilamentosum* W. Fray and Kurschner A3, D3 - (R)
 47. *Crossidium squamiferum* (Viv.) Jur. A2 - (R)
 48. *Ctenidium molluscum* (Hedw.) Mitt. C2 - (R)
 49. *Cynodontium polycarpon* (Hedw.) Schimp. D2 - (R)
 50. *Dichodontium pellucidum* (Hedw.) Schimp. D2, D3 - (K)
 51. *Dicranella heteromalla* (Hedw.) Schimp. D3 - (K)
 52. *Dicranella varia* (Hedw.) Schimp. D2 - (K)
 53. *Dicranum scoparium* Hedw. B3, D3 - (R)
 54. *Didymodon cordatus* Jur. A3, C3, D3 - (R)
 55. *Didymodon fallax* (Hedw.) R. H. Zander A3, C2, D4 - (K)
 56. *Didymodon insulanus* (De Not.) M.O. Hill. A3, B3, D2, D3, E2 - (K)
 57. *Didymodon luridus* Hornsch. A2, B3, C2 - (K)
 58. *Didymodon rigidulus* Hedw. A2, B2, B3, C2 - (S)
 59. *Didymodon spadiceus* (Mitt.) Limpr. D1, D2 - (R)
 60. *Didymodon tophaceus* (Brid.) Lisa D1, D2 - (R)
 61. *Didymodon vinealis* (Brid.) R. H. Zander A2, A3, B2, B3, C2, D2, D3, D4, E4 - (S)
 62. *Ditrichum flexicaule* (Schwagr.) Hampe B3, D3, D4 - (K)
 63. *Ditrichum heteromallum* (Hedw.) E. Britton B3, D3, D4 - (K)
 64. *Ditrichum pallidum* (Hedw.) Hampe B3, D4 - (R)
 65. *Ditrichum pussillum* (Hedw.) Hampe D2, D4 - (R)
 66. *Drepanocladus polygamus* (Schimp.) Hedenas C1 - (R)
 67. *Encalypta streptocarpa* Hedw. D2, D3 - (K)
 68. *Encalypta vulgaris* Hedw. B3 - (R)
 69. *Enthostodon* sp. B1, D1 - (R)
 70. *Entodon concinnus* (De Not.) Paris D1 1
 71. *Eucladium verticillatum* (With.) Bruch and Schimp. D2 - (R)
 72. *Eurhynchiastrum pulchellum* (Hedw.) Ignatov and Huttunen A2, C3, D2, D3 - (K)
 73. *Eurhynchium striatum* (Hedw.) Schimp. C2, C3, D4 - (K)
 74. *Fissidens adianthoides* Hedw. C3, D3 - (K)
 75. *Fissidens bryoides* Hedw. D3 - (R)
 76. *Fissidens dubius* P. Beauv. C3, D2 - (K)
 77. *Fissidens taxifolius* Hedw. C3, D3 - (K)
 78. *Funaria hygrometrica* Hedw. A2, A3, B3, C3 - (S)
 79. *Grimmia orbicularis* Bruch ex Wilson D2 - (R)
 80. *Grimmia pulvinata* (Hedw.) Sm. A2, B2, C2, D2, D4, E2, E4 - (S)
 81. *Grimmia trichophylla* Grev. B3, D1, E2 - (K)
 82. *Gymnostomum aeruginosum* Sm. D1, D2 - (R)
 83. *Gymnostomum calcareum* Nees and Hornsch. A2, B2 - (R)
 84. *Herzogiella seligeri* (Brid.) Z. Iwats. D4, E4 - (K)
 85. *Hilpertia velenovskyi* (Schiffn.) R. H. Zander - A3 - (R)
 86. *Homalia trichomanoides* (Hedw.) Brid. D4 - (R)
 87. *Homalothecium aureum* (Spruce) H. Rob. A3 - (R)
 88. *Homalothecium lutescens* (Hedw.) H. Rob. D1, D2, D3, D4, C2, C4, E2, E4 - (S)
 89. *Homalothecium phillippeanum* (Spruce) Schimp. C2, C4, D2, D4 - (K)
 90. *Homalothecium sericeum* (Hedw.) Schimp. B2, B3, C2, D2, D3, D4 - (S)
 91. *Homomalium incurvatum* (Schrad. ex Brid.) Loeske D4, E4 - (K)
 92. *Hygroamblystegium tenax* (Hedw.) Jenn. A1, C2, D1, D2, D4 - (K)
 93. *Hygroamblystegium varium* (Hedw.) Monk. A3, C2, C4, D2, D4, E2, E4 - (S)
 94. *Hygrohypnum luridum* (Hedw.) Jenn. D1 - (R)
 95. *Hypnum andoi* A.J.E.Sm. D4 - (R)
 96. *Hypnum cupressiforme* Hedw. var *cupressiforme* A2, A4, B2, C2, C4, D2, D3, D4, E2, E4 - (S)
 97. *Hypnum jutlandicum* Holmen and E.Warncke D4 - (R)
 98. *Hypnum lacuinoisum* (Brid.) Hoffm. ex Brid. C4, D4 - (K)
 99. *Hypnum resupinatum* Taylor D4 - (K)
 100. *Hypnum tectorum* Brid. - D4 - (R)
 101. *Isopterygyopsis muelleriana* (Schimp.) Z. Iwats. C4, D2, D4 - (K)
 102. *Isopterygyopsis pulchella* (Hedw.) Z. Iwats. D4 - (R)
 103. *Isothecium alopecuroides* (Lam. ex Dubois) Isov. C2, D2 - (K)
 104. *Isothecium myosuroides* Brid. C2, D2, D3, D4 - (S)
 105. *Kindbergia praelonga* (Hedw.) Ochyra A3, C3, D3, D4 - (K)
 106. *Leptodyctium riparium* (Hedw.) Warnst. A2, B3, C2, C4, D1 - (K)
 107. *Lescurea mutabilis* (Brid.) Lindb. ex I. Hagen D2, D3, D4 - (R)
 108. *Leskeia polycarpa* Hedw. A2, B2, C4, D1, D2, D3, D4, E2, E4 - (S)
 109. *Leucodon sciuroides* (Hedw.) Schwagr. D4 - (R)
 110. *Mnium stellare* Hedw. D3 - (R)
 111. *Orthothecium intricatum* (Hartm.) Schimp. C4 - (R)
 112. *Orthothecium rufescens* (Dicks. ex Brid.) Schimp. D2 - (R)
 113. *Orthotrichum affine* Schrad. ex Brid. C2, D1, D4, E2 - (K)
 114. *Orthotrichum anomalum* Hedw. A2, B2, C2, D2, D4 - (S)
 115. *Orthotrichum cupulatum* Hoffm. ex Brid. C2, E4 - (R)
 116. *Orthotrichum diaphanum* Schrad. ex Brid. A2, B2, C2, C4, D1, D2, D4, E2, E4 - (S)

117. *Orthotrichum lyellii* Hook. and Taylor D4, E4 - (K)
118. *Orthotrichum patens* Bruch ex Brid. D4 - (R)
119. *Orthotrichum pumilum* Sw. D2, E4 - (R)
120. *Orthotrichum rupestre* Schleich. ex Schwagr. D2, D4 - (K)
121. *Orthotrichum speciosum* Nees E4 - (R)
122. *Orthotrichum stramineum* Hornsch. ex Brid. E4 - (R)
123. *Orthotrichum striatum* Hedw. D2, D3, E4 - (R)
124. *Orthotrichum tenellum* Bruch ex Brid. D2 - (R)
125. *Oxyrrhynchium hians* (Hedw.) Loeske B3, C2, C3, D2, D3 - (S)
126. *Oxyrrhynchium speciosum* (Brid.) Warnst. C3, D1, D2 - (K)
127. *Oxystegus tenuirostris* (Hook. and Taylor) A. J. E. Sm. D2 - (R)
128. *Palustriella commutata* (Hedw.) Ochyra D1 - (R)
129. *Phascum cuspidatum* Hedw. A3, B3, C3, D3 - (K)
130. *Physcomitrella patens* (Hedw.) Bruch and Schimp. - B1, B3, D3 - (R)
131. *Physcomitrium pyriforme* (Hedw.) Bruch and Schimp. - D1, D3 - (R)
132. *Plagiommium affine* (Blandow ex Funck) T.J. Kop. E2 - (R)
133. *Plagiommium cuspidatum* (Hedw.) T. Kop. C3 - (R)
134. *Plagiommium elatum* (Bruch and Schimp.) T.J. Kop. D3, D4 - (R)
135. *Plagiommium medium* (Bruch and Schimp.) T.J. Kop. D3 - (R)
136. *Plagiommium rostratum* (Schrad.) T. J. Kop. E2 - (R)
137. *Plagiommium undulatum* (Hedw.) T. J. Kop. B3, C3, D3 - (K)
138. *Plagiothecium cavifolium* (Brid.) Z. Iwats. D3 - (R)
139. *Plasteurhynchium meridionale* (Schimp.) M.Fleisch. A2 - (R)
140. *Platyhypnidium ripariooides* (Hedw.) Dixon C3, D3 - (K)
141. *Pleuridium acuminatum* Lindb. D3 - (R)
142. *Pleurochaete squarrosa* (Brid.) Lindb. - C3, D3 - (K)
143. *Pohlia cruda* (Hedw.) Lindb. D2 - (R)
144. *Pohlia melanodon* (Brid.) A. J. Shaw. - B1, B3 - (R)
145. *Polytrichum commune* Hedw. D3 - (R)
146. *Pseudocrossidium hornschuchianum* (Schultz) R. H. Zander C3, D3 - (K)
147. *Pseudoleskella nervosa* (Brid.) Nyholm D2, D4 - (K)
148. *Pterygynandrum filiforme* Hedw. D2, D4 - (K)
149. *Pylaisia polyantha* (Hedw.) Schimp. C4, D2, D3, D4, E4 - (S)
150. *Rhynchostegiella tenella* (Dicks.) Limpr. C1, C2, D2 - (K)
151. *Rhynchostegiella teneriffae* (Mont.) Dirkse and Bouman C2 - (R)
152. *Rhynchostegium confertum* (Dicks.) Schimp. A2, A3, D3, D4, E3 - (R)
153. *Rhynchostegium megapolitanum* (Blandow ex F. Weber and D. Mohr) Schimp. E2 - (R)
154. *Schistidium apocarpum* (Hedw.) Bruch and Schimp. A2, B2, C2, D2, E2 - (S)
155. *Schistidium crassipilum* H. H. Blom A2, B2, C2, D2 - (K)
156. *Schistidium elegantulum* H. H. Blom D2 - (R)
157. *Schistidium helveticum* (Schkuhr) Deguchi A2, B2, C2, D2 - (K)
158. *Sciuro-hypnum flotowianum* (Sendtn.) Ignatov and Huttunen C3 - (R)
159. *Sciuro-hypnum plumosum* (Hedw.) Ignatov and Huttunen C3, D2, D4 - (R)
160. *Sciuro-hypnum populeum* (Hedw.) Ignatov and Huttunen C2, C3, D4 - (K)
161. *Scleropodium touretii* (Brid.) L.F. Koch C2, D3 - (R)
162. *Syntrichia calcicola* J. J. Amann B2, C3, D2 (K)
163. *Syntrichia laevipila* Brid. D2, D4, E2 - (K)
164. *Syntrichia montana* Nees D2 - (R)
165. *Syntrichia papillosa* (Wilson) Jur. D4, E4 - (K)
166. *Syntrichia princeps* (De Not.) Mitt. C2 - (K)
167. *Syntrichia ruraliformis* (Besch.) Cardot - C3, D2 - (R)
168. *Syntrichia ruralis* (Hedw.) F. Weber and D. Mohr A2, D2, D4, E2 - (S)
169. *Syntrichia virescens* (De Not.) Ochyra D4, E2 - (R)
170. *Taxiphyllum wissgrillii* (Garov.) Wijk and Margad. D4 - (R)
171. *Thuidium tamariscinum* (Hedw.) Schimp. D3 - (R)
172. *Tortella fragilis* (Hook. and Wilson) Limpr. B3 - (R)
173. *Tortella tortuosa* (Hedw.) Limpr. B3, D3 - (R)
174. *Tortula inermis* (Brid.) Mont. C2 - (R)
175. *Tortula lanceola* R. H. Zander C3, D3 - (R)
176. *Tortula muralis* Hedw. var. *aestiva* Hedw. A3, C2, C4 - (R)
177. *Tortula muralis* Hedw. var. *muralis* A2, A3, B3, C2, C4, D2, E2 - (S)
178. *Tortula schimperi* M.J. Cano, O. Werner and J. Guerra - D3 - (R)
179. *Tortula subulata* Hedw. B3, D2, D3 - (K)
180. *Tortula truncata* (Hedw.) Mitt. C3, D4 - (K)
181. *Trichodon cylindricus* (Hedw.) Schimp. D3 - (R)
182. *Trichostomum brachydontium* Bruch A3, C2, C4, D2, E4 - (K)
183. *Trichostomum crispulum* Bruch A2, A3, D3 - (K)
184. *Ulota crispa* (Hedw.) Brid. D2, D4 - (R)
185. *Warnstorffia exannulata* (Schimp.) Loeske C4 - (R)
186. *Weissia brachycarpa* (Nees and Hornsch.) Jur. D2, D4 - (R)
187. *Weissia controversa* Hedw. D3 - (R)

DISCUSSION

Belgrade has rich bryophyte flora. However, hepatics are not abundant and any species does not enter the category of spread (S) within the investigated area. This gives us the idea on Belgrade climate where the very dry long summers combined with disturbed and non adequate habitats prevent the large presence of hepatics.

Among 187 moss taxa recorded species found up to two times which means rare (R) are the most presented in Belgrade (94 taxa). The 24 spread taxa (S) were recorded more than 7 times within Belgrade city area. Common moss taxa to record in Belgrade city (recorded by us 3-7 times) are 69.

According to the data obtained, Belgrade has many small microhabitats and contains many significant populations of some threatened Serbian and European mosses.

Three species (*Callicladium haldanianum* (EN), *Hilpertia velenovskyi* (EN) and *Hypnum andoi* (VU) are nationally and/or internationally threatened taxa (Sabovljević *et al.*, 2004; ECCB, 1995). Further, eleven species are of lower risk in Serbia: *Crossidium crassinerve*, *C. laxofilamentosum*, *Ditrichum pallidum*, *D. pussilum*, *Hygroamblystegium tenax*, *Isopterygyopsis pulchella*, *Physcomitrella patens*, *Physcomitrium pyriforme*, *Rhynchostegiella teneriffae*, *Scleropodium touretii* and *Syntrichia papillosa*. One data deficient species was recorded as well: *Anomodon longifolius*.

Especially interesting habitats are Belgrade fragile loess cliffs sites in Zemun, Bežanija, Donji Kalemegdan and Višnjička Kosa with its specific cryptogamic flora and rare and endangered species like *Hilpertia velenovskyi* and *Crossidium laxofilamentosum*. The fragile ecology are already discussed by Sabovljević (2003a, 2004) and Pócs *et al.* (2004).

The second record for Serbia of *Schistidium elegantulum*, known previously only from the road edge in Obrež, Srem (Sabovljević, 2003b) was made, along the Sava on the concrete.

The bryologically richest site in wider Belgrade is Avala Mt. Data covers wider area of Avala Mt. can be found in Sabovljević and Cvetić (2003).

Comparing Belgrade bryophyte diversity to the other European cities, it is one of the highest (Fig. 2). Eventhough, the sizes and the geographical positions of the cities compared are not took into consideration the urban flora of Belgrade is very rich and interesting not only from the diversity point but also from the biogeographical, ecological and taxonomic point of view.

Further investigations in distribution and vegetational relationships among urban bryophytes are needed.




Fig. 2: Comparison of bryophyte diversity richness in some European cities (hepa-hepatic, moss-mosses, bryo - bryophytes)

ACKNOWLEDGMENT

We acknowledge Grant No. 143015 of Serbian Ministry of Science.

REFERENCES

- Aleffi, M., 1991. Bryophyte flora and air quality in the town of Jesi (Marche, Central Italy). Arch. Bot. Ital., 67: 128-140.
- Aleffi, M. and E. Taruschio, 1996. Bryophyte flora and air quality in the town of Treia (Marche, Central Italy). Arch. Geobot., 2: 47-58.
- Ballesteros Segura, T. and M.E. Ron, 1985. Contribution to bryophyte flora of the city of Toledo. Anales Jard. Bot., 42: 87-91.
- Benkert, D., P. Erzberger, J. Klawitter, W. Linder, C. Linke, A. Schaepe, M. Steinland and W. Wiegle, 1995. The moss list of Brandenburg and Berlin including the threat levels. Verh. Bot., 128: 1-68.
- Bento-Pereira, F. and C. Sérgio, 1983. Lichens and bryophytes as bioindicators of atmospheric pollution. Rev. Biologia, 12: 297-313.
- Brandes, D., 1983. Flora and vegetation of the railways of Middle Europe. Phytocoenologia, 11: 31-115.
- Carcano, L., 1989. Moss inventory of the urban area of Rome (Italy). Braun Blanquetia, 3: 147-150.
- Casas, C. and C. Saiz, 1982. The bryophytes of the cathedral of Sevilla. Collect. Bot., 13: 163-175.
- Cortini Pedrotti, C., 1989. Bryophyte flora of the Camerino town (Central Italy). Braun Blanquetia, 3: 75-86.
- Dia, M.G. and R. Not, 1991. Biodetrmental agents of monumental in historical center of the Palermo city. Quad. Bot. Ambientale Appl., 2: 3-10.
- ECCB, 1995. Red Data Book of European Bryophytes. Trondheim.

- Esteve, F., J. Varo and M.L. Zafra, 1977. Bryological studies of the Granada city. Trab. Dep. Bot. Univ. Granada, 4: 45-71.
- Fiol, A., 1983. Bryophytes of urban habitats of Palma de Mallorca. Bol. Soc. Hist. Nat. Balears, 27: 65-76.
- Frahm, J.P., 2006. Notulae bryologicae rhenanae 7. Moss flora of Rolandseck. Arch. Bryology, 12: 1-7.
- Frahm, J.P. and M. Sabovljević, 2007. Reduction of microdust by mosses. Immissionsschutz, 4: 152-156.
- Franzen, I., 2001. Epiphytic mosses and lichens as bioindicators of air quality in west edge of Ruhr region. Limprichtia, 16: 1-23.
- Fudali, E., 1994. Species diversity and spatial distribution of bryophytes in urban areas-A case study of the city of Szczecin. Fragm. Fl. Geobot., 39: 563-570.
- Gilbert, O.L., 1968. Bryophytes as indicators of air pollution in the Tyne Valley. New Phytologist, 67: 15-30.
- Gilbert, O.L., 1970. Urban bryophyte communities in North-East England. Trans. Br. Bryological Soc., 6: 306-316.
- Gilbert, O.L., 1989. The Ecology of Urban Habitats. Chapman and Hall.
- Giudice, R.L., V. Mazimpaka and F. Lara, 1997. The urban bryophyte flora of the city of Enna (Sicily, Italy). Nova Hedw., 64: 249-265.
- Grdović, S. and V. Stevanović, 2006. The moss flora of central urban area of Belgrade. Arch. Biol. Sci., 58: 55-59.
- Heras, P. and A. Soria, 1990. Urban mosses and hepaticas of the city of Vitoria-Gasteiz. Soc. Est. Vascos, Secc. Ci. Nat., 7: 75-116.
- Hill, M.O., N. Bell, M.A. Bruggeman-Nannenga, M. Brugues and M.J. Cano *et al.*, 2006. An annotated checklist of the mosses of Europe and Macaronesia. J. Bryol., 28: 198-267.
- Hohenwallner, D., 2000a. Bioindication with mosses in city area of Vienna. Limprichtia, 15: 1-88.
- Hohenwallner, D., 2000b. Viennas' Chinese Wall. Dérive, 1: 15-15.
- Hohenwallner, D. and H.G. Zechmeister, 2001a. Noteworthy moss records in the Vienna downtown Linzer Biologische Beiträge, 33: 295-298.
- Hohenwallner D. and H.G. Zechmeister, 2001b. Factors influencing bryophyte species richness and populations in urban environments: A case study. Nova Hedwigia, 73: 87-96.
- Humer-Hochwimmer, K. and H.G. Zechmeister, 2001. Epiphytic mosses in Vienna forest within the city of Vienna and its significance for bioindication of atmospheric pollutants. Limprichtia, 18: 1-99.
- Jovanović, S., 1994. Ecological study of the Ruderal flora and Vegetation of Belgrade. 1st Edn., Univezitet u Beogradu, Biološki fakultet.
- Lara, F. and V. Mazimpaka, 1989. Contribution to the bryophyte flora of the city of Segovia. Madrid, Anales Jard. Bot., 46: 481-485.
- Lara, F., C. Lopez and V. Mazimpaka, 1991. Bryophyte ecology in the town of Segovia, Spain [Ecología de los briofitos urbanos en la ciudad de Segovia (España)] Cryptogamie. Bryologie et Lichenologie, 12: 425-439.
- Lo Giudice, R., 1992. Contribution to the urban bryophyte flora of Catania. Quad. Bot. Ambientale Appl., 3: 3-10.
- Mazimpaka, V., J. Vicente and E. Ron, 1988. Contribution to the urban bryophyte flora of the city of Madrid. Anales Jard. Bot., 45: 61-73.
- Mazimpaka, V., F. Lara and C. Lopez-Garcia, 1993. Ecological data on the bryophyte flora of the town of Cuenca (Spain). Nova Hedwigia, 56: 113-129.
- Nickl-Navrátil, H., 1960. Small moss communities of towns. Nova Hedwigia, 2: 425-462.
- Pavletić, Z., 1955. Prodromus Flore Briofita Jugoslavije. 1st Edn., JAZU, Zagreb.
- Pokorný, L., F. Lara and V. Mazimpaka, 2006. The bryophyte flora of the city of Trento (North Italy). Cryptogamie Bryol., 27: 265-284.
- Pócs, T., M. Sabovljević, F. Puche, J.G. Segarra Moragues, C. Gimeno and H. Kürschner, 2004. Crossidium laxofilamentosum Frey and Kürschner (Pottiaceae), new to Europe and to North Africa. Studies on the cryptogamic vegetation on loess cliffs, VII. J. Bryol., 26: 113-124.
- Rams, S., R.M. Ros, M.J. Cano and J. Guerra, 2000. Urban bryophyte flora of the city of Murcia. Boletín de la Sociedad Espanola de Briología, 7: 9-18.
- Ron, E., V. Mazimpaka, J. Vicente and I. Granzow de la Cerda, 1987. Urban bryophytes in Spanish towns. Symp. Biol. Hung., 35: 727-753.
- Ros, R.M., V. Mazimpaka, U. Abou-Salma, M. Aleffi and T.L. Blockeel *et al.*, 2007. Hepaticas and Anthocerotes of the Mediterranean, an annotated checklist. Cryptogamie Bryol., 28: 351-437.
- Sabovljević, M., 2003a. Bryophyte flora of South Banat (Vojvodina, Yugoslavia). Cryptogamie Bryol., 24: 241-252.
- Sabovljević, M., 2003b. The data on presence and chorology of taxa from the genus *Schistidium* Bruch and *Schimpfer* (Grimmiaceae) in the Federal Republic of Yugoslavia. Bocconeia, 16: 991-999.
- Sabovljević, M. and T. Cvetic, 2003. Bryophyte flora of Avala Mt. (C. Serbia, Yugoslavia). Lindbergia, 28: 90-96.

- Sabovljević, M., 2004. Life strategies of bryophytes on loess cliffs in Vojvodina (Serbia). *Arch. Biol. Sci.*, 56: 127-130.
- Sabovljević, M., T. Cvetic and V. Stenevanovic, 2004. Bryophyte red list of Serbia and Montenegro. *Biodiversity and Conservat.*, 13: 1781-1790.
- Sabovljević, M., V. Vukojević, N. Mihailović, G. Dražić and Ž. Vučinić, 2005. Determination of heavy metal deposition in the county of Obrenovac (Serbia) using mosses as bioindicators. I: Aluminium (Al), Arsenic (As) and Boron (B). *Arch. Biol. Sci.*, 57: 205-212.
- Sabovljević, M. and R. Natceva, 2006. Check list of the liverworts and hornworts of South-East Europe. *Phytol. Balc.*, 12: 169-180.
- Sabovljević, M., V. Vukojević, A. Sabovljević, N. Mihailović, G. Dražić and Ž. Vučinić, 2007. Determination of heavy metal deposition in the county of Obrenovac (Serbia) using mosses as bioindicators III. Copper (Cu), Iron (Fe) and Mercury (Hg). *Arch. Biol. Sci.*, 59: 351-361.
- Sabovljević, M., R. Natceva, G. Dihoru, E. Tsakiri, S. Dragičević, A. Erdag and B. Papp, 2008. Check list of the mosses of South-East Europe. *Phytol. Balc.*, 14: 207-244.
- Sabovljević, M. and A. Sabovljević, 2009. Biodiversity within urban areas: A case study on bryophytes of the city of Cologne (NRW, Germany). *Plant Biosystems*.
- Schaepe, A., 1986. Changes of the mossflora of Berlin (West). *Bryophytorum Bibliotheca*, 33: 1-392.
- Soška, T., 1949. Survey of the mosses and lichens of the Belgrade surroundings. *Glasnik muzeja Srpske zemlje*, Ser. B, 1-2: 93-112.
- Sérgio, C., 1981. Changes in epiphytic bryophyte flora in urban area of Lisbon in last 140 years. *Bol. Soc. Brot. Sér.*, 2: 291-303.
- Sérgio, C. and F. Bento-Pereira, 1981. Lichens and bryophytes as bioindicators of atmospheric pollution. I. *Bol. Soc. Brot. Sér.*, 2: 313-331.
- Solga, A., J. Burkhardt, H.G. Zechmeister and J.P. Frahm, 2005. Nitrogen content, ^{15}N natural abundance and biomass of the two pleurocarpous mosses *Pleurozium schreberi* (Brid.) Mitt. and *Scleropodium purum* (Hedw.) Limpr. in relation to atmospheric nitrogen deposition. *Environ. Pollut.*, 134: 465-473.
- Solga, A. and J.P. Frahm, 2006. Nitrogen accumulation of six pleurocarpous moss species and their suitability for monitoring nitrogen deposition. *J. Bryol.*, 28: 46-52.
- Solga, A., T. Eichert and J.P. Frahm, 2006a. Historical alteration in the nitrogen content and ^{15}N natural abundance of mosses in western Germany: Indication for regionally varying changes in atmospheric nitrogen deposition within the last 150 years. *Atmos. Environ.*, 40: 8044-8055.
- Solga, A., J. Burkhardt and J.P. Frahm, 2006b. A new approach to assess atmospheric nitrogen deposition by way of standardized exposition of mosses. *Environ. Monit. Assess.*, 116: 399-417.
- Soria, A. and M.E. Ron, 1990. Contribution to the urban bryophyte flora of the city of Logrono. Madrid, *Anales Jard. Bot.*, 46: 427-432.
- Soria, A., M.E. Ron and P. Heras, 1992. Comparative analysis of the urban bryoflora of Vitoria-Gasteiz with the others Spanish cities. *Actes Simposi Internac. Bot. Pius Font I. Quer.*, pp: 271-276.
- Soria, A. and M.E. Ron, 1995. Contributions to the spanish urban bryophyte flora. *Cryptogamie Bryol. Lichénol.*, 16: 285-299.
- Vanderpoorten, A., 1997. A bryological survey of the Brussels capital region (Belgium). *Scripta Botanica Belgica*, 14: 1-40.
- Vareschi, V., 1936. Epiphytic vegetation of Zurich. *Ber. Schweiz. Bot. Ges.*, 46: 445-488.
- Vicente, J., I.G. De La Cerda, V. Mazimpaka and M.E. Ron, 1986. Contribution to the urban bryoflora of the city of Avila. *Trab. Dep. Bot.*, 13: 45-49.
- Viera, C. and M.E. Ron, 1986. Contribution to the urban bryoflora of the city of Badajoz. *Trap. Dep. Bot.*, 13: 45-49.
- Vukojević, V., M. Sabovljević and S. Jovanović, 2005. Mosses accumulate heavy metals from the substracta of coal ash. *Arch. Biol. Sci.*, 57: 101-106.
- Vukojević, V., M. Sabovljević, A. Sabovljević, N. Mihailović, G. Dražić and Ž. Vučinić, 2006. Determination of heavy metal deposition in the county of Obrenovac (Serbia) using mosses as bioindicators II: Cadmium (Cd), Cobalt (Co) and Chromium (Cr). *Arch. Biol. Sci.*, 58: 95-104.
- Zechmeister, H.G., D. Hohenwallner and K. Humer-Hochwimmer, 2001. The researches of the mossflora of Vienna. *Berichte der Reinhold-Tüxen-Gesellschaft*, 13: 291-295.